
Zhihan Wang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1571019/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Unsaturated fatty-acid based HPLC fingerprints in combination with quantitative analysis of multi-components by single-marker for the classification of <i>Rana chensinensis</i> ovum. New Journal of Chemistry, 2022, 46, 10441-10450.	2.8	1
2	Antioxidant Activity Evaluation of Oviductus Ranae Protein Hydrolyzed by Different Proteases. Molecules, 2021, 26, 1625.	3.8	6
3	Crystal engineering construction of caffeic acid derivatives with potential applications in pharmaceuticals and degradable polymeric materials. CrystEngComm, 2020, 22, 7847-7857.	2.6	2
4	Synthesis and characterization of BPA-free polyesters by incorporating a semi-rigid cyclobutanediol monomer. Polymer Chemistry, 2020, 11, 6081-6090.	3.9	11
5	Renewable Cyclobutane-1,3-dicarboxylic Acid (CBDA) Building Block Synthesized from Furfural via Photocyclization. ACS Sustainable Chemistry and Engineering, 2020, 8, 8909-8917.	6.7	22
6	Antitussive and Anti-inflammatory Dual-active Agents Developed from Natural Product Lead Compound 1-Methylhydantoin. Molecules, 2019, 24, 2355.	3.8	18
7	Quality Evaluation of Oviductus Ranae Based on PUFAs Using HPLC Fingerprint Techniques Combined with Chemometric Methods. Foods, 2019, 8, 322.	4.3	12
8	Scalable preparation and property investigation of a cis-cyclobutane-1,2-dicarboxylic acid from β-trans-cinnamic acid. Chemical Communications, 2019, 55, 214-217.	4.1	21
9	Protein-Based Fingerprint Analysis for the Identification of Ranae Oviductus Using RP-HPLC. Molecules, 2019, 24, 1687.	3.8	18
10	Environmental influences on quality features of <i>Oviductus Ranae</i> in the Changbai Mountains. RSC Advances, 2019, 9, 36050-36057.	3.6	7
11	Furfural-Derived Diacid Prepared by Photoreaction for Sustainable Materials Synthesis. ACS Sustainable Chemistry and Engineering, 2018, 6, 8136-8141.	6.7	25
12	Cyclobutane-1,3-Diacid (CBDA): A Semi-Rigid Building Block Prepared by [2+2] Photocyclization for Polymeric Materials. Scientific Reports, 2017, 7, 13704.	3.3	18
13	Polyladderane Constructed from a Gemini Monomer through Photoreaction. Angewandte Chemie, 2017, 129, 12323-12327.	2.0	9
14	Polyladderane Constructed from a Gemini Monomer through Photoreaction. Angewandte Chemie - International Edition, 2017, 56, 12155-12159.	13.8	27
15	Lighting the Way to Greener Chemistry: Incandescent Floodlights as a Facile UV Light Source for Classic and Cutting-Edge Photoreactions. ACS Sustainable Chemistry and Engineering, 2016, 4, 5053-5059.	6.7	20
16	Stereoregular Two-Dimensional Polymers Constructed by Topochemical Polymerization. Macromolecules, 2015, 48, 2894-2900.	4.8	46
17	Linear polyester synthesized from furfural-based monomer by photoreaction in sunlight. Green Chemistry, 2015, 17, 4720-4724.	9.0	40
18	A two-dimensional hydrogen bonded organic framework self-assembled from a three-fold symmetric carbamate. Chemical Communications, 2014, 50, 5209-5211.	4.1	24

ZHIHAN WANG

#	Article	IF	CITATIONS
19	An unsaturated hydrogen bonded network generated from three-fold symmetric carbamates. CrystEngComm, 2014, 16, 7176-7179.	2.6	5
20	Synthesis of polymeric ladders by topochemical polymerization. Chemical Communications, 2014, 50, 1218-1220.	4.1	24
21	Synthesis of 5-bromo-6-methyl imidazopyrazine, 5-bromo and 5-chloro-6-methyl imidazopyridine using electron density surface maps to guide synthetic strategy. Tetrahedron, 2011, 67, 9063-9066.	1.9	10
22	Scalable Synthesis of 4-Substituted 5-Bromo-6-methylpyrimidines. Synthesis, 2011, 2011, 1529-1531.	2.3	1
23	Isolation and identification of two steroid compounds from <i>Oviductus Ranae</i> . Natural Product Research, 2010, 24, 1518-1522.	1.8	14