

Philip W Askenase

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1602586/publications.pdf>

Version: 2024-02-01

53
papers

2,730
citations

201575

27
h-index

175177

52
g-index

58
all docs

58
docs citations

58
times ranked

2587
citing authors

#	ARTICLE	IF	CITATIONS
1	Exosome Carrier Effects; Resistance to Digestion in Phagolysosomes May Assist Transfers to Targeted Cells; II Transfers of miRNAs Are Better Analyzed via Systems Approach as They Do Not Fit Conventional Reductionist Stoichiometric Concepts. <i>International Journal of Molecular Sciences</i> , 2022, 23, 6192.	1.8	5
2	Ancient Evolutionary Origin and Properties of Universally Produced Natural Exosomes Contribute to Their Therapeutic Superiority Compared to Artificial Nanoparticles. <i>International Journal of Molecular Sciences</i> , 2021, 22, 1429.	1.8	18
3	Exosomes provide unappreciated carrier effects that assist transfers of their miRNAs to targeted cells; I. They are "The Elephant in the Room"™. <i>RNA Biology</i> , 2021, 18, 1-16.	1.5	8
4	Antibodies Enhance the Suppressive Activity of Extracellular Vesicles in Mouse Delayed-Type Hypersensitivity. <i>Pharmaceuticals</i> , 2021, 14, 734.	1.7	5
5	Small extracellular vesicles released by infused mesenchymal stromal cells target M2 macrophages and promote TGF-β ² upregulation, microvascular stabilization and functional recovery in a rodent model of severe spinal cord injury. <i>Journal of Extracellular Vesicles</i> , 2021, 10, e12137.	5.5	71
6	Rare Skin Reactions after mRNA Vaccination, Similar to Jones' "Mote Basophil Responses. <i>New England Journal of Medicine</i> , 2021, 385, 1720-1721.	13.9	6
7	Orally Administered Exosomes Suppress Mouse Delayed-Type Hypersensitivity by Delivering miRNA-150 to Antigen-Primed Macrophage APC Targeted by Exosome-Surface Anti-Peptide Antibody Light Chains. <i>International Journal of Molecular Sciences</i> , 2020, 21, 5540.	1.8	22
8	COVID-19 therapy with mesenchymal stromal cells (MSC) and convalescent plasma must consider exosome involvement: Do the exosomes in convalescent plasma antagonize the weak immune antibodies?. <i>Journal of Extracellular Vesicles</i> , 2020, 10, e12004.	5.5	43
9	Artificial nanoparticles are not as good as the real thing. <i>Nature</i> , 2020, 582, S5-S5.	13.7	2
10	Syngeneic red blood cell-induced extracellular vesicles suppress delayed-type hypersensitivity to self-antigens in mice. <i>Clinical and Experimental Allergy</i> , 2019, 49, 1487-1499.	1.4	15
11	Delayed-Type Hypersensitivity Underlying Casein Allergy Is Suppressed by Extracellular Vesicles Carrying miRNA-150. <i>Nutrients</i> , 2019, 11, 907.	1.7	23
12	Intravenously administered contact allergens coupled to syngeneic erythrocytes induce in mice tolerance rather than effector immune response. <i>Folia Medica Cracoviensia</i> , 2019, 59, 61-73.	0.3	2
13	Antibody Light Chains Dictate the Specificity of Contact Hypersensitivity Effector Cell Suppression Mediated by Exosomes. <i>International Journal of Molecular Sciences</i> , 2018, 19, 2656.	1.8	15
14	Intravenously delivered mesenchymal stem cell-derived exosomes target M2-type macrophages in the injured spinal cord. <i>PLoS ONE</i> , 2018, 13, e0190358.	1.1	164
15	The cationic small molecule GW4869 is cytotoxic to high phosphatidylserine-expressing myeloma cells. <i>British Journal of Haematology</i> , 2017, 177, 423-440.	1.2	24
16	Expression of activation-induced cytidine deaminase enhances the clearance of pneumococcal pneumonia: evidence of a subpopulation of protective anti-pneumococcal B1a cells. <i>Immunology</i> , 2016, 147, 97-113.	2.0	19
17	Functions of Exosomes and Microbial Extracellular Vesicles in Allergy and Contact and Delayed-Type Hypersensitivity. <i>International Archives of Allergy and Immunology</i> , 2016, 171, 1-26.	0.9	39
18	Epicutaneous immunization with ovalbumin and CpG induces TH1/TH17 cytokines, which regulate IgE and IgG2a production. <i>Journal of Allergy and Clinical Immunology</i> , 2016, 138, 262-273.e6.	1.5	21

#	ARTICLE	IF	CITATIONS
19	A subset of AID-dependent B α 1a cells initiates hypersensitivity and pneumococcal pneumonia resistance. <i>Annals of the New York Academy of Sciences</i> , 2015, 1362, 200-214.	1.8	21
20	From Mysterious Supernatant Entity to miRNA-150 in Antigen-Specific Exosomes: a History of Hapten-Specific T Suppressor Factor. <i>Archivum Immunologiae Et Therapiae Experimentalis</i> , 2015, 63, 345-356.	1.0	16
21	Macrophages play an essential role in antigen-specific immune suppression mediated by T _{scp} CD ₈ ⁺ cell-derived exosomes. <i>Immunology</i> , 2015, 146, 23-32.	2.0	48
22	Diagnostic and therapeutic potentials of exosomes in CNS diseases. <i>Brain Research</i> , 2015, 1617, 63-71.	1.1	120
23	Free Extracellular miRNA Functionally Targets Cells by Transfecting Exosomes from Their Companion Cells. <i>PLoS ONE</i> , 2015, 10, e0122991.	1.1	59
24	Epicutaneous immunization with phosphorylcholine conjugated to bovine serum albumin (PC-BSA) and TLR9 ligand CpG alleviates pneumococcal pneumonia in mice. <i>Pharmacological Reports</i> , 2014, 66, 570-575.	1.5	3
25	Antigen-specific, antibody-coated, exosome-like nanovesicles deliver suppressor T-cell microRNA-150 to effector T cells to inhibit contact sensitivity. <i>Journal of Allergy and Clinical Immunology</i> , 2013, 132, 170-181.e9.	1.5	187
26	Natural killer cell-mediated contact sensitivity develops rapidly and depends on interferon α , interferon β and interleukin β . <i>Immunology</i> , 2013, 140, 98-110.	2.0	71
27	Immunobiology of Antigen-Specific Immunoglobulin Free Light Chains in Chronic Inflammatory Diseases. <i>Current Pharmaceutical Design</i> , 2012, 18, 2278-2289.	0.9	12
28	Stimulatory Lipids Accumulate in the Mouse Liver within 30 min of Contact Sensitization to Facilitate the Activation of Na α -ve iNKT Cells in a CD1d-Dependent Fashion. <i>Scandinavian Journal of Immunology</i> , 2011, 74, 52-61.	1.3	19
29	Identification of Initiator B Cells, a Novel Subset of Activation-Induced Deaminase-Dependent B-1-Like Cells That Mediate Initiation of Contact Sensitivity. <i>Journal of Immunology</i> , 2008, 181, 1717-1727.	0.4	29
30	Interleukin-4-dependent innate collaboration between iNKT cells and B-1 B cells controls adaptive contact sensitivity. <i>Immunology</i> , 2006, 117, 536-547.	2.0	30
31	Invariant NKT Cells Rapidly Activated via Immunization with Diverse Contact Antigens Collaborate In Vitro with B-1 Cells to Initiate Contact Sensitivity. <i>Journal of Immunology</i> , 2006, 177, 3686-3694.	0.4	49
32	An Hour after Immunization Peritoneal B-1 Cells Are Activated to Migrate to Lymphoid Organs Where within 1 Day They Produce IgM Antibodies That Initiate Elicitation of Contact Sensitivity. <i>Journal of Immunology</i> , 2005, 175, 7170-7178.	0.4	64
33	TLR-Dependent IL-4 Production by Invariant V α 14 β 18 γ NKT Cells to Initiate Contact Sensitivity In Vivo. <i>Journal of Immunology</i> , 2005, 175, 6390-6401.	0.4	62
34	Extravascular T-cell recruitment requires initiation begun by V α 14 β NKT cells and B-1 B cells. <i>Trends in Immunology</i> , 2004, 25, 441-449.	2.9	81
35	Cutaneous Immunization Rapidly Activates Liver Invariant V α 14 NKT Cells Stimulating B-1 B Cells to Initiate T Cell Recruitment for Elicitation of Contact Sensitivity. <i>Journal of Experimental Medicine</i> , 2003, 198, 1785-1796.	4.2	154
36	B-1 B Cells Mediate Required Early T Cell Recruitment to Elicit Protein-Induced Delayed-Type Hypersensitivity. <i>Journal of Immunology</i> , 2003, 171, 6225-6235.	0.4	76

#	ARTICLE	IF	CITATIONS
37	Subunits of IgM Reconstitute Defective Contact Sensitivity in B-1 Cell-Deficient Mice: Light Chains Recruit T Cells Independent of Complement. <i>Journal of Immunology</i> , 2002, 169, 4113-4123.	0.4	30
38	B Cell-dependent T Cell Responses. <i>Journal of Experimental Medicine</i> , 2002, 196, 1277-1290.	4.2	114
39	Yes T cells, but three different T cells (Th2, Th1 and NK T cells), and also B-1 cells mediate contact sensitivity. <i>Clinical and Experimental Immunology</i> , 2001, 125, 345-350.	1.1	65
40	Topical tacrolimus and cyclosporin A differentially inhibit early and late effector phases of cutaneous delayed-type and immunoglobulin E hypersensitivity. <i>Immunology</i> , 2001, 104, 235-242.	2.0	21
41	Proposing Th2 DTH Relevant to Asthma: Cutaneous Basophil Hypersensitivity Then and Now. , 2000, 78, 112-123.		4
42	Early Local Generation of C5a Initiates the Elicitation of Contact Sensitivity by Leading to Early T Cell Recruitment. <i>Journal of Immunology</i> , 2000, 165, 1588-1598.	0.4	108
43	Required Early Complement Activation in Contact Sensitivity with Generation of Local C5-dependent Chemotactic Activity, and Late T Cell Interferon γ : A Possible Initiating Role of B Cells. <i>Journal of Experimental Medicine</i> , 1997, 186, 1015-1026.	4.2	81
44	Blockade of CD2-LFA-3 interactions protects human skin allografts in immunodeficient mouse/human chimeras. <i>Nature Biotechnology</i> , 1997, 15, 759-762.	9.4	59
45	Elicitation of Nickel Sulfate (NiSO ₄)-Specific Delayed-Type Hypersensitivity Requires Early-Occurring and Early-Acting, NiSO ₄ -Specific DTH-Initiating Cells with an Unusual Mixed Phenotype for an Antigen-Specific Cell. <i>Cellular Immunology</i> , 1995, 161, 244-255.	1.4	17
46	DNFB Contact Sensitivity (CS) In BALB/c and C3H/He Mice: Requirement for Early-Occurring,		