

Tim J Croudace

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1603358/publications.pdf>

Version: 2024-02-01

130
papers

8,686
citations

38720

50
h-index

48277

88
g-index

131
all docs

131
docs citations

131
times ranked

10474
citing authors

#	ARTICLE	IF	CITATIONS
1	Association Between Duration of Untreated Psychosis and Outcome in Cohorts of First-Episode Patients. <i>Archives of General Psychiatry</i> , 2005, 62, 975.	13.8	1,303
2	Incidence of Schizophrenia and Other Psychoses in England, 1950–2009: A Systematic Review and Meta-Analyses. <i>PLoS ONE</i> , 2012, 7, e31660.	1.1	385
3	Outcomes of conduct problems in adolescence: 40 year follow-up of national cohort. <i>BMJ: British Medical Journal</i> , 2009, 338, a2981-a2981.	2.4	277
4	The Short Mood and Feelings Questionnaire (SMFQ): A Unidimensional Item Response Theory and Categorical Data Factor Analysis of Self-Report Ratings from a Community Sample of 7-through 11-Year-Old Children. <i>Journal of Abnormal Child Psychology</i> , 2006, 34, 365-377.	3.5	271
5	Reappraising the long-term course and outcome of psychotic disorders: the AESOP-10 study. <i>Psychological Medicine</i> , 2014, 44, 2713-2726.	2.7	192
6	Mokken scale analysis of mental health and well-being questionnaire item responses: a non-parametric IRT method in empirical research for applied health researchers. <i>BMC Medical Research Methodology</i> , 2012, 12, 74.	1.4	175
7	Psychometric evaluation and predictive validity of Ryff's psychological well-being items in a UK birth cohort sample of women. <i>Health and Quality of Life Outcomes</i> , 2006, 4, 76.	1.0	169
8	Clinical and social determinants of duration of untreated psychosis in the Å†SOP first-episode psychosis study. <i>British Journal of Psychiatry</i> , 2006, 189, 446-452.	1.7	168
9	The effects of maternal postnatal depression and child sex on academic performance at age 16 years: a developmental approach. <i>Journal of Child Psychology and Psychiatry and Allied Disciplines</i> , 2010, 51, 1150-1159.	3.1	161
10	Early developmental milestones in adult schizophrenia and other psychoses. A 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Schizophrenia Research</i> , 2001, 52, 1-19.	1.1	159
11	Friendships and Family Support Reduce Subsequent Depressive Symptoms in At-Risk Adolescents. <i>PLoS ONE</i> , 2016, 11, e0153715.	1.1	151
12	A Longitudinal Typology of Symptoms of Depression and Anxiety Over the Life Course. <i>Biological Psychiatry</i> , 2007, 62, 1265-1271.	0.7	147
13	Recovering Quality of Life (ReQoL): a new generic self-reported outcome measure for use with people experiencing mental health difficulties. <i>British Journal of Psychiatry</i> , 2018, 212, 42-49.	1.7	141
14	Characteristics of Outcome in Schizophrenia at 13 Years. <i>British Journal of Psychiatry</i> , 1995, 167, 596-603.	1.7	137
15	Mood, anxiety and psychotic phenomena measure a common psychopathological factor. <i>Psychological Medicine</i> , 2015, 45, 1483-1493.	2.7	128
16	Mortality in Schizophrenia and Other Psychoses: A 10-Year Follow-up of the Å”SOP First-Episode Cohort. <i>Schizophrenia Bulletin</i> , 2015, 41, 664-673.	2.3	128
17	Forty-Year Psychiatric Outcomes Following Assessment for Internalizing Disorder in Adolescence. <i>American Journal of Psychiatry</i> , 2007, 164, 126-133.	4.0	122
18	Measuring mental health and wellbeing outcomes for children and adolescents to inform practice and policy: a review of child self-report measures. <i>Child and Adolescent Psychiatry and Mental Health</i> , 2014, 8, 14.	1.2	111

#	ARTICLE	IF	CITATIONS
19	The Course of Schizophrenia Over 13 Years. <i>British Journal of Psychiatry</i> , 1996, 169, 580-586.	1.7	110
20	Developmental Trajectories of Sex-typed Behavior in Boys and Girls: A Longitudinal General Population Study of Children Aged 2.5-8 Years. <i>Child Development</i> , 2008, 79, 1583-1593.	1.7	107
21	Cognitive behaviour therapy for improving social recovery in psychosis: a report from the ISREP MRC Trial Platform study (Improving Social Recovery in Early Psychosis). <i>Psychological Medicine</i> , 2009, 39, 1627.	2.7	101
22	Randomised trial of a parenting intervention during neonatal intensive care. <i>Archives of Disease in Childhood: Fetal and Neonatal Edition</i> , 2007, 92, F438-F443.	1.4	97
23	Elevated morning cortisol is a stratified population-level biomarker for major depression in boys only with high depressive symptoms. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 3638-3643.	3.3	97
24	An Evaluation of the Precision of Measurement of Ryff's Psychological Well-Being Scales in a Population Sample. <i>Social Indicators Research</i> , 2010, 97, 357-373.	1.4	93
25	Profiles of family-focused adverse experiences through childhood and early adolescence: The ROOTS project a community investigation of adolescent mental health. <i>BMC Psychiatry</i> , 2011, 11, 109.	1.1	92
26	Non-linear relationship between an index of social deprivation, psychiatric admission prevalence and the incidence of psychosis. <i>Psychological Medicine</i> , 2000, 30, 177-185.	2.7	91
27	Timing of menarche and depressive symptoms in adolescent girls from a UK cohort. <i>British Journal of Psychiatry</i> , 2011, 198, 17-23.	1.7	91
28	Ten-Year Outcomes of First-Episode Psychoses in the MRC A-SOP-10 Study. <i>Journal of Nervous and Mental Disease</i> , 2015, 203, 379-386.	0.5	90
29	Analysis of cluster randomized cross-over trial data: a comparison of methods. <i>Statistics in Medicine</i> , 2007, 26, 274-289.	0.8	89
30	Aggressive incidents in first-episode psychosis. <i>British Journal of Psychiatry</i> , 2001, 178, 433-440.	1.7	88
31	Serotonin transporter genotype, morning cortisol and subsequent depression in adolescents. <i>British Journal of Psychiatry</i> , 2009, 195, 39-45.	1.7	87
32	Reduction in adolescent depression after contact with mental health services: a longitudinal cohort study in the UK. <i>Lancet Psychiatry</i> , 2017, 4, 120-127.	3.7	86
33	Rumination, anxiety, depressive symptoms and subsequent depression in adolescents at risk for psychopathology: a longitudinal cohort study. <i>BMC Psychiatry</i> , 2013, 13, 250.	1.1	84
34	Developmental Typology of Trajectories to Nighttime Bladder Control: Epidemiologic Application of Longitudinal Latent Class Analysis. <i>American Journal of Epidemiology</i> , 2003, 157, 834-842.	1.6	76
35	Diagnostic change 10 years after a first episode of psychosis. <i>Psychological Medicine</i> , 2015, 45, 2757-2769.	2.7	74
36	Three-year outcome of first-episode psychoses in an established community psychiatric service. <i>British Journal of Psychiatry</i> , 2000, 176, 210-216.	1.7	73

#	ARTICLE	IF	CITATIONS
37	The predictive validity of a diagnosis of schizophrenia. <i>British Journal of Psychiatry</i> , 1997, 170, 321-327.	1.7	71
38	Impact of the ICD-10 Primary Health Care (PHC) diagnostic and management guidelines for mental disorders on detection and outcome in primary care. <i>British Journal of Psychiatry</i> , 2003, 182, 20-30.	1.7	71
39	Ethnicity and long-term course and outcome of psychotic disorders in a UK sample: The A-SOP-10 study. <i>British Journal of Psychiatry</i> , 2017, 211, 88-94.	1.7	71
40	Childhood and adolescent predictors of schizophrenia in the Northern Finland 1966 Birth Cohort - a descriptive life-span model. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2000, 250, 311-319.	1.8	63
41	The relationship between early personality and midlife psychological well-being: evidence from a UK birth cohort study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2008, 43, 679-687.	1.6	62
42	Cohort Profile: Risk patterns and processes for psychopathology emerging during adolescence: the ROOTS project. <i>International Journal of Epidemiology</i> , 2010, 39, 361-369.	0.9	62
43	Classifying Patients by Antipsychotic Adherence Patterns Using Latent Class Analysis: Characteristics of Nonadherent Groups in the California Medicaid (Medi-Cal) Program. <i>Value in Health</i> , 2008, 11, 48-56.	0.1	61
44	Polymorphisms in <i>BDNF</i> (Val66Met) and <i>5-HTTLPR</i> , morning cortisol and subsequent depression in at-risk adolescents. <i>British Journal of Psychiatry</i> , 2010, 197, 365-371.	1.7	60
45	Comparative Effectiveness of Cognitive Therapies Delivered Face-to-Face or over the Telephone: An Observational Study Using Propensity Methods. <i>PLoS ONE</i> , 2012, 7, e42916.	1.1	60
46	Association between pubertal development and depressive symptoms in girls from a UK cohort. <i>Psychological Medicine</i> , 2012, 42, 2579-2589.	2.7	57
47	Outcome of psychosis in people of African-Caribbean family origin. <i>British Journal of Psychiatry</i> , 1999, 175, 43-49.	1.7	56
48	The persistence of developmental markers in childhood and adolescence and risk for schizophrenic psychoses in adult life. A 34-year follow-up of the Northern Finland 1966 birth cohort. <i>Schizophrenia Research</i> , 2004, 71, 213-225.	1.1	55
49	Childhood adversity subtypes and depressive symptoms in early and late adolescence. <i>Development and Psychopathology</i> , 2015, 27, 885-899.	1.4	54
50	Schizophrenia and other psychotic disorders in Caribbean-born migrants and their descendants in England: systematic review and meta-analysis of incidence rates, 1950-2013. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2015, 50, 1039-1055.	1.6	54
51	Discharged from special hospital under restrictions: a comparison of the fates of psychopaths and the mentally ill. <i>Criminal Behaviour and Mental Health</i> , 1998, 8, 39-55.	0.4	53
52	Calibrating well-being, quality of life and common mental disorder items: Psychometric epidemiology in public mental health research. <i>British Journal of Psychiatry</i> , 2016, 209, 162-168.	1.7	53
53	Biased Mentalizing in Children Aged Seven to 11: Latent Class Confirmation of Response Styles to Social Scenarios and Associations with Psychopathology. <i>Social Development</i> , 2007, 16, 181-202.	0.8	52
54	Latent mixture models for multivariate and longitudinal outcomes. <i>Statistical Methods in Medical Research</i> , 2010, 19, 271-289.	0.7	52

#	ARTICLE	IF	CITATIONS
55	Health-Related Quality of Life (HRQL) and Continuous Antipsychotic Treatment: 3-year Results from the Schizophrenia Health Outcomes (SOHO) Study. <i>Value in Health</i> , 2009, 12, 536-543.	0.1	51
56	Educational consequences of mental disorders treated in hospital. A 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Psychological Medicine</i> , 2001, 31, 339-349.	2.7	50
57	Comparing growth trajectories of risk behaviors from late adolescence through young adulthood: An accelerated design. <i>Developmental Psychology</i> , 2013, 49, 1732-1738.	1.2	50
58	Trajectories of Daytime Wetting and Soiling in a United Kingdom 4 to 9-Year-Old Population Birth Cohort Study. <i>Journal of Urology</i> , 2008, 179, 1970-1975.	0.2	49
59	General and specific components of depression and anxiety in an adolescent population. <i>BMC Psychiatry</i> , 2011, 11, 191.	1.1	47
60	Clinical effectiveness and cost-effectiveness of tailored intensive liaison between primary and secondary care to identify individuals at risk of a first psychotic illness (the LEGs study): a cluster-randomised controlled trial. <i>Lancet Psychiatry</i> , 2015, 2, 984-993.	3.7	47
61	Evaluation of ethnic disparities in detection of depression and anxiety in primary care during the maternal period: Combined analysis of routine and cohort data. <i>British Journal of Psychiatry</i> , 2016, 208, 453-461.	1.7	46
62	Randomized Trial of a Parenting Intervention for Very Preterm Infants: Outcome at 2 Years. <i>Journal of Pediatrics</i> , 2009, 155, 488-494.e1.	0.9	44
63	Measuring the benefits of treatment for psychosis: validity and responsiveness of the EQ-5D. <i>British Journal of Psychiatry</i> , 2009, 195, 170-177.	1.7	41
64	5-HTTLPR and Early Childhood Adversities Moderate Cognitive and Emotional Processing in Adolescence. <i>PLoS ONE</i> , 2012, 7, e48482.	1.1	39
65	Early development, stress and depression across the life course: pathways to depression in a national British birth cohort. <i>Psychological Medicine</i> , 2014, 44, 2845-2854.	2.7	38
66	On Dimensionality, Measurement Invariance, and Suitability of Sum Scores for the PHQ-9 and the GAD-7. <i>Assessment</i> , 2022, 29, 355-366.	1.9	38
67	Birth order and risk for schizophrenia: a 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Acta Psychiatrica Scandinavica</i> , 2001, 104, 148-152.	2.2	35
68	Cohort profile: mental health following extreme trauma in a northern Ugandan cohort of War-Affected Youth Study (The WAYS Study). <i>SpringerPlus</i> , 2013, 2, 300.	1.2	35
69	Psychological Distress during Pregnancy in a Multi-Ethnic Community: Findings from the Born in Bradford Cohort Study. <i>PLoS ONE</i> , 2013, 8, e60693.	1.1	34
70	Is the incidence of psychotic disorder in decline? Epidemiological evidence from two decades of research. <i>International Journal of Epidemiology</i> , 2009, 38, 1255-1264.	0.9	33
71	Childhood cognitive function and adult psychopathology: associations with psychotic and non-psychotic symptoms in the general population. <i>British Journal of Psychiatry</i> , 2012, 201, 124-130.	1.7	33
72	Cortisol levels in response to starting school in children at increased risk for social phobia. <i>Psychoneuroendocrinology</i> , 2012, 37, 462-474.	1.3	33

#	ARTICLE	IF	CITATIONS
73	A comparison of approaches for assessing covariate effects in latent class analysis. <i>Longitudinal and Life Course Studies</i> , 2015, 6, .	0.3	33
74	Public mental health: evidenced-based priorities. <i>Lancet, The</i> , 2015, 385, 1472-1475.	6.3	31
75	Measuring patients' views: a bifactor model of distinct patient-reported outcomes in psychosis. <i>Psychological Medicine</i> , 2011, 41, 277-289.	2.7	29
76	Reliability Generalization for Childhood Autism Rating Scale. <i>Journal of Autism and Developmental Disorders</i> , 2013, 43, 2855-2865.	1.7	29
77	Cognitive behaviour therapy for improving social recovery in psychosis: Cost-effectiveness analysis. <i>Schizophrenia Research</i> , 2009, 112, 158-163.	1.1	28
78	Postwar environment and long-term mental health problems in former child soldiers in Northern Uganda: the WAYS study. <i>Journal of Epidemiology and Community Health</i> , 2014, 68, 425-430.	2.0	28
79	Factors associated with antidepressant, anxiolytic and hypnotic use over 17 years in a national cohort. <i>Journal of Affective Disorders</i> , 2008, 110, 234-240.	2.0	27
80	Protocol for the Mindful Student Study: a randomised controlled trial of the provision of a mindfulness intervention to support university students' well-being and resilience to stress. <i>BMJ Open</i> , 2016, 6, e012300.	0.8	27
81	Prevalence of and recovery from common mental disorder including psychotic experiences in the UK Primary Care Improving Access to Psychological Therapies (IAPT) Programme. <i>Journal of Affective Disorders</i> , 2020, 272, 84-90.	2.0	27
82	Factors of psychological distress: clinical value, measurement substance, and methodological artefacts. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2015, 50, 515-524.	1.6	26
83	Outcomes following first-episode psychosis – “Why we should intervene early in all ages, not only in youth. <i>Australian and New Zealand Journal of Psychiatry</i> , 2016, 50, 1055-1063.	1.3	25
84	Lifetime affect and midlife cognitive function: prospective birth cohort study. <i>British Journal of Psychiatry</i> , 2014, 204, 194-199.	1.7	24
85	A modelling strategy for the analysis of clinical trials with partly missing longitudinal data. <i>International Journal of Methods in Psychiatric Research</i> , 2003, 12, 139-150.	1.1	23
86	The validity of subjective quality of life measures in psychotic patients with severe psychopathology and cognitive deficits: an item response model analysis. <i>Quality of Life Research</i> , 2012, 21, 237-246.	1.5	23
87	Ten-year outcomes in first episode psychotic major depression patients compared with schizophrenia and bipolar patients. <i>Schizophrenia Research</i> , 2016, 176, 417-422.	1.1	23
88	Deciding who gets treatment for depression and anxiety: a study of consecutive GP attenders. <i>British Journal of General Practice</i> , 2005, 55, 846-53.	0.7	23
89	Neurological Signs at the First Psychotic Episode as Correlates of Long-Term Outcome: Results From the AESOP-10 Study. <i>Schizophrenia Bulletin</i> , 2021, 47, 118-127.	2.3	20
90	Three decades of antidepressant, anxiolytic and hypnotic use in a national population birth cohort. <i>British Journal of Psychiatry</i> , 2006, 189, 156-160.	1.7	19

#	ARTICLE	IF	CITATIONS
91	Meta-analytic approaches to determine gender differences in the age-incidence characteristics of schizophrenia and related psychoses. <i>International Journal of Methods in Psychiatric Research</i> , 2013, 22, 36-45.	1.1	19
92	The influence of different types of war experiences on depression and anxiety in a Ugandan cohort of war-affected youth: the WAYS study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2014, 49, 1783-1792.	1.6	19
93	Biological and psychosocial risk factors for psychotic major depression. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016, 51, 233-245.	1.6	19
94	Improvements in social functioning reported by a birth cohort in mid-adult life: A person-centred analysis of GHQ-28 social dysfunction items using latent class analysis. <i>Personality and Individual Differences</i> , 2007, 42, 305-316.	1.6	18
95	Adolescent self-organization predicts midlife memory in a prospective birth cohort study.. <i>Psychology and Aging</i> , 2013, 28, 958-968.	1.4	18
96	An evaluation of computerized adaptive testing for general psychological distress: combining GHQ-12 and Affectometer-2 in an item bank for public mental health research. <i>BMC Medical Research Methodology</i> , 2016, 16, 58.	1.4	17
97	Improvement and decline of cognitive function in schizophrenia over one year: a longitudinal investigation using latent growth modelling. <i>BMC Psychiatry</i> , 2007, 7, 16.	1.1	16
98	Who Do You Think Is in Control in Addiction? A Pilot Study on Drug-related Locus of Control Beliefs. <i>Addictive Disorders and Their Treatment</i> , 2012, 11, 195-205.	0.5	16
99	The catechol-O-methyltransferase gene (COMT) and cognitive function from childhood through adolescence. <i>Biological Psychology</i> , 2013, 92, 359-364.	1.1	16
100	Heterogeneity in Patient-Reported Outcomes following Low-Intensity Mental Health Interventions: A Multilevel Analysis. <i>PLoS ONE</i> , 2014, 9, e99658.	1.1	16
101	Symptom remission at 12-weeks strongly predicts long-term recovery from the first episode of psychosis. <i>Psychological Medicine</i> , 2020, 50, 1452-1462.	2.7	16
102	A randomised controlled trial assessing the severity and duration of depressive symptoms associated with a clinically significant response to sertraline versus placebo, in people presenting to primary care with depression (PANDA trial): study protocol for a randomised controlled trial. <i>Trials</i> , 2017, 18, 496.	0.7	15
103	Juvenile and early adulthood smoking and adult educational achievements – A 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Scandinavian Journal of Public Health</i> , 2001, 29, 87-95.	1.2	14
104	A longitudinal investigation of the impact of typology of urinary incontinence on quality of life during midlife: Results from a British prospective study. <i>Maturitas</i> , 2009, 64, 246-248.	1.0	14
105	Factors influencing the costs of epilepsy in adults with an intellectual disability. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2012, 21, 205-210.	0.9	14
106	Validity of single item responses to short message service texts to monitor depression: an mHealth sub-study of the UK ACUDep trial. <i>BMC Medical Research Methodology</i> , 2015, 15, 56.	1.4	14
107	Maternal psychological distress in primary care and association with child behavioural outcomes at age three. <i>European Child and Adolescent Psychiatry</i> , 2016, 25, 601-613.	2.8	13
108	Clinical services for adults with an intellectual disability and epilepsy: A comparison of management alternatives. <i>PLoS ONE</i> , 2017, 12, e0180266.	1.1	13

#	ARTICLE	IF	CITATIONS
109	Multilevel ordinal factor analysis of the Positive and Negative Syndrome Scale (PANSS). <i>International Journal of Methods in Psychiatric Research</i> , 2014, 23, 25-35.	1.1	12
110	Monoamine Oxidase A (MAOA) Gene and Personality Traits from Late Adolescence through Early Adulthood: A Latent Variable Investigation. <i>Frontiers in Psychology</i> , 2017, 8, 1736.	1.1	12
111	Depressive symptoms during adolescence: comparison between epidemiological and high risk sampling. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2012, 47, 1333-1341.	1.6	11
112	Exploring the item sets of the Recovering Quality of Life (ReQoL) measures using factor analysis. <i>Quality of Life Research</i> , 2019, 28, 1005-1015.	1.5	11
113	Psychiatric outcomes 10 years after treatment with antidepressants or anxiolytics. <i>British Journal of Psychiatry</i> , 2008, 193, 327-331.	1.7	10
114	Comparison of high and low intensity contact between secondary and primary care to detect people at ultra-high risk for psychosis: study protocol for a theory-based, cluster randomized controlled trial. <i>Trials</i> , 2013, 14, 222.	0.7	10
115	Predictors of human rotation. <i>Laterality</i> , 2013, 18, 265-281.	0.5	10
116	Adolescent Self-control Predicts Midlife Hallucinatory Experiences: 40-Year Follow-up of a National Birth Cohort. <i>Schizophrenia Bulletin</i> , 2014, 40, 1543-1551.	2.3	10
117	Item bank measurement of depression: will one dimension work?. <i>Journal of Clinical Epidemiology</i> , 2014, 67, 4-6.	2.4	9
118	Diagnostic stability and outcome after first episode psychosis. <i>Journal of Mental Health</i> , 2021, 30, 104-112.	1.0	9
119	Computerized adaptive testing of population psychological distress: simulation-based evaluation of GHQ-30. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016, 51, 895-906.	1.6	8
120	Use of the theory of planned behaviour to assess factors influencing the identification of individuals at ultra-high risk for psychosis in primary care. <i>Microbial Biotechnology</i> , 2012, 6, 265-275.	0.9	7
121	The validity of patient- and clinician-rated measures of needs and the therapeutic relationship in psychosis: A pooled analysis. <i>Psychiatry Research</i> , 2013, 209, 711-720.	1.7	7
122	Factor analysis of treatment outcomes from a UK specialist addiction service: Relationship between the Leeds Dependence Questionnaire, Social Satisfaction Questionnaire and 10-item Clinical Outcomes in Routine Evaluation. <i>Drug and Alcohol Review</i> , 2014, 33, 643-650.	1.1	7
123	Changes of hand preference in Parkinson's disease. <i>Journal of Neural Transmission</i> , 2012, 119, 693-696.	1.4	6
124	Effects of ignoring clustered data structure in confirmatory factor analysis of ordered polytomous items: a simulation study based on PANSS. <i>International Journal of Methods in Psychiatric Research</i> , 2016, 25, 205-219.	1.1	6
125	An item response theory analysis of an item pool for the recovering quality of life (ReQoL) measure. <i>Quality of Life Research</i> , 2021, 30, 267-276.	1.5	6
126	A Life Course Approach to Well-Being. , 2007, , 187-205.		6

#	ARTICLE	IF	CITATIONS
127	Antidepressant treatment with sertraline for adults with depressive symptoms in primary care: the PANDA research programme including RCT. Programme Grants for Applied Research, 2019, 7, 1-108.	0.4	6
128	Determinants of patient-reported outcome trajectories and symptomatic recovery in Improving Access to Psychological Therapies (IAPT) services. Psychological Medicine, 2022, 52, 3231-3240.	2.7	4
129	BILOG-MG, MULTILog, PARSCALE and TESTFACT4 programs on CD with single volume manual combining reference guide and users guide (program examples: Input and output). British Journal of Mathematical and Statistical Psychology, 2005, 58, 193-195.	1.0	3
130	Understanding causes of and developing effective interventions for schizophrenia and other psychoses. Programme Grants for Applied Research, 2016, 4, 1-184.	0.4	3