## Lotfi Mabrouk

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1608061/publications.pdf

Version: 2024-02-01

840776 940533 19 258 11 16 citations h-index g-index papers 19 19 19 302 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Temporal and depth distribution of microepiphytes on Posidonia oceanica (L.) Delile leaves in a meadow off Tunisia. Marine Ecology, 2011, 32, 148-161.	1.1	61
2	What factors drive seasonal variation of phytoplankton, protozoans and metazoans on leaves of Posidonia oceanica and in the water column along the coast of the Kerkennah Islands, Tunisia?. Marine Pollution Bulletin, 2013, 71, 286-298.	5.0	39
3	Variability in the structure of epiphyte assemblages on leaves and rhizomes of Posidonia oceanica in relation to human disturbances in a seagrass meadow off Tunisia. Aquatic Botany, 2013, 108, 33-40.	1.6	19
4	Fluctuating asymmetry in grass goby Zosterisessor ophiocephalus Pallas, 1811 inhabiting polluted and unpolluted area in Tunisia. Marine Pollution Bulletin, 2014, 85, 248-251.	5.0	18
5	Variability in the Structure of Phytoplankton Assemblages in relation to Human Disturbance in Southern Coast of Tunisia. Journal of Marine Biology, 2014, 2014, 1-10.	1.0	16
6	A Comparison of Abundance and Diversity of Epiphytic Microalgal Assemblages on the Leaves of the Seagrasses <i>Posidonia oceanica</i> (L.) and <i>Cymodocea nodosa</i> (Ucria) Asch in Eastern Tunisia. Journal of Marine Biology, 2014, 2014, 1-10.	1.0	14
7	Temporal and spatial zonation of macroepiphytes on <i>Posidonia oceanica</i> (L.) Delile leaves in a meadow off Tunisia. Marine Ecology, 2015, 36, 77-92.	1.1	13
8	Variability patterns of epibenthic microalgae in eastern Tunisian coasts. Scientia Marina, 2017, 81, 487.	0.6	13
9	High leaf fluctuating asymmetry in two native plants growing in heavy metal-contaminated soil: the case of Metlaoui phosphate mining basin (Gafsa, Tunisia). Environmental Monitoring and Assessment, 2020, 192, 406.	2.7	12
10	Bathymetric variation of epiphytic assemblages on Posidonia oceanica (L.) Delile leaves in relation to anthropogenic disturbance in the southeastern Mediterranean. Environmental Science and Pollution Research, 2014, 21, 13588-13601.	5.3	11
11	Spatial scale variability in shoot density and epiphytic leaves of <i>Posidonia oceanica</i> on Kerkennah Island (Tunisia) in relation to current tide effects. Marine Ecology, 2015, 36, 1311-1331.	1.1	11
12	Diversity and temporal fluctuations of epiphytes and sessile invertebrates on the rhizomes <i><scp>P</scp>osidonia oceanica</i> in a seagrass meadow off <scp>T</scp> unisia. Marine Ecology, 2014, 35, 212-220.	1.1	9
13	Comparison of spatial scale variability of shoot density and epiphytic leaf assemblages of Halophila stipulacea and Cymodocea nodosa on the Eastern Coast of Tunisia. Plant Biosystems, 2020, 154, 413-426.	1.6	8
14	Variability in the Structure of Planktonic Microalgae Assemblages in Water Column Associated withPosidonia oceanica(L.) Bed in Tunisia. Journal of Marine Biology, 2014, 2014, 1-7.	1.0	4
15	Temporal and bathymetric variation of epiphyte cover and leaf biomass in a southern <i><scp>P</scp>osidonia oceanica</i> ( <scp>L</scp> .) meadow: the case of <scp>M</scp> ahdia coast, <scp>T</scp> unisia. Marine Ecology, 2017, 38, e12394.	1.1	4
16	Variability in the structure of epiphytic microalgae assemblages on the leaves of <em>Posidonia oceanica</em> in relation to human disturbance in a meadow off Tunisia. Scientia Marina, 2014, 78, 27-39.	0.6	4
17	Spatial patterns and floristic composition of plant communities in uninhabited islets: the case of Kaboudia (Chebba, Eastern Tunisia). Plant Biosystems, 2021, 155, 251-266.	1.6	1
18	Comparison of epiphyte algal assemblages on the leaves of marine seagrasses <i>Posidonia oceanica</i> (L.) Delile, <i>Cymodocea nodosa</i> (Ucria) Asch, and the lessepsian <i>Halophila stipulacea</i> (Forssk.) Asch in Chebba (East of Tunisia). Marine Ecology, 2021, 42, e12642.	1.1	1

#	Article	IF	CITATIONS
19	Factors controlling phytoplankton dynamics in an arid reservoir in Tunisia (case of Sidi Saad dam). Environmental Monitoring and Assessment, 2021, 193, 354.	2.7	0