
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1608522/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Hepatic Arterial Infusion of Oxaliplatin, Fluorouracil, and Leucovorin Versus Transarterial Chemoembolization for Large Hepatocellular Carcinoma: A Randomized Phase III Trial. Journal of Clinical Oncology, 2022, 40, 150-160. | 0.8 | 137 |
| 2 | Tenofovir vs. entecavir on prognosis of hepatitis B virus-related hepatocellular carcinoma after curative resection. Journal of Gastroenterology, 2022, 57, 185-198. | 2.3 | 16 |
| 3 | Drug-coated balloon combined with provisional drug-eluting stent implantation for the treatment of de novo Medina 0,1,0 or 0,0,1 left main coronary bifurcation lesions: A proof-ofconcept study. , 2022, 26, 218-225. | | 2 |
| 4 | Lenvatinib, toripalimab, plus hepatic arterial infusion chemotherapy <i>versus</i> lenvatinib alone for advanced hepatocellular carcinoma. Therapeutic Advances in Medical Oncology, 2021, 13, 175883592110027. | 1.4 | 91 |
| 5 | Effect of prior cancer on survival of hepatocellular carcinoma: implications for clinical trial eligibility criteria. BMC Cancer, 2021, 21, 147. | 1.1 | 1 |
| 6 | Comparison of HBV reactivation between patients with high HBV-DNA and low HBV-DNA loads undergoing PD-1 inhibitor and concurrent antiviral prophylaxis. Cancer Immunology, Immunotherapy, 2021, 70, 3207-3216. | 2.0 | 21 |
| 7 | Cancerâ€Associated Fibroblastâ€Mediated Cellular Crosstalk Supports Hepatocellular Carcinoma Progression. Hepatology, 2021, 73, 1717-1735. | 3.6 | 147 |
| 8 | Hepatic Arterial Infusion Chemotherapy of Oxaliplatin, Fluorouracil, and Leucovorin With or Without Sorafenib as Initial Treatment for Advanced Hepatocellular Carcinoma. Frontiers in Oncology, 2021, 11, 619461. | 1.3 | 14 |
| 9 | Prognostic Values of Alpha-Fetoprotein and Des-Gamma-Carboxyprothrombin in Hepatocellular Carcinoma in China: An Analysis of 4792 Patients. Journal of Hepatocellular Carcinoma, 2021, Volume 8, 657-670. | 1.8 | 12 |
| 10 | Reductions in AFP and PIVKA-II can predict the efficiency of anti-PD-1 immunotherapy in HCC patients. BMC Cancer, 2021, 21, 775. | 1.1 | 28 |
| 11 | Development and validation of prognostic nomograms for single large and huge hepatocellular carcinoma after curative resection. European Journal of Cancer, 2021, 155, 85-96. | 1.3 | 19 |
| 12 | Comparative safety and efficacy of molecular-targeted drugs, immune checkpoint inhibitors, hepatic arterial infusion chemotherapy and their combinations in advanced hepatocellular carcinoma: findings from advances in landmark trials. Frontiers in Bioscience, 2021, 26, 873. | 0.8 | 6 |
| 13 | Angiographic quantitative flow ratio-guided coronary intervention (FAVOR III China): a multicentre, randomised, sham-controlled trial. Lancet, The, 2021, 398, 2149-2159. | 6.3 | 175 |
| 14 | Clinical outcomes in 2481 unselected real-world patients treated with a polymer-free sirolimus-eluting stent: 3Âyears results from the NANO multicenter Registry. BMC Cardiovascular Disorders, 2021, 21, 537. | 0.7 | 1 |
| 15 | Optimal subsequent treatments for patients with hepatocellular carcinoma resistant to anti-PD-1 treatment. Immunotherapy, 2021, , . | 1.0 | 0 |
| 16 | Development and Validation of a Prognostic Score for Hepatocellular Carcinoma Patients in Immune Checkpoint Inhibitors Therapies: The Hepatocellular Carcinoma Modified Gustave Roussy Immune Score. Frontiers in Pharmacology, 2021, 12, 819985. | 1.6 | 9 |
| 17 | Radiofrequency ablation versus laparoscopic hepatectomy for hepatocellular carcinoma: A real world single center study. European Journal of Surgical Oncology, 2020, 46, 548-559. | 0.5 | 23 |
| 18 | Optical coherence tomography enables more accurate detection of functionally significant intermediate non-left main coronary artery stenoses than intravascular ultrasound: A meta-analysis of 6919 patients and 7537 lesions. International Journal of Cardiology, 2020, 301, 226-234. | 0.8 | 19 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Stereotactic Body Radiotherapy as a Salvage Therapy after Incomplete Radiofrequency Ablation for Hepatocellular Carcinoma: A Retrospective Cohort Study. Journal of Oncology, 2020, 2020, 1-9. | 0.6 | 6 |
| 20 | Stereotactic Body Radiotherapy vs. Radiofrequency Ablation in the Treatment of Hepatocellular Carcinoma: A Meta-Analysis. Frontiers in Oncology, 2020, 10, 1639. | 1.3 | 22 |
| 21 | Feasibility of using deep learning to detect coronary artery disease based on facial photo. European Heart Journal, 2020, 41, 4400-4411. | 1.0 | 67 |
| 22 | <p>Baseline HBV Loads Do Not Affect the Prognosis of Patients with Hepatocellular Carcinoma Receiving Anti-Programmed Cell Death-1 Immunotherapy</p> . Journal of Hepatocellular Carcinoma, 2020, Volume 7, 337-345. | 1.8 | 6 |
| 23 | Single versus multiple port laparoscopic left lateral sectionectomy for hepatocellular carcinoma: A retrospective comparative study. International Journal of Surgery, 2020, 77, 15-21. | 1.1 | 3 |
| 24 | Tescalcin is an unfavorable prognosis factor that regulats cell proliferation and survival in hepatocellular carcinoma patients. Cancer Communications, 2020, 40, 355-369. | 3.7 | 16 |
| 25 | Oneâ€year clinical results of the NANO registry: A multicenter, prospective allâ€comers registry study in patients receiving implantation of a polymerâ€free sirolimusâ€eluting stent. Catheterization and Cardiovascular Interventions, 2020, 95, 658-664. | 0.7 | 8 |
| 26 | Intention to control low central venous pressure reduced blood loss during laparoscopic hepatectomy: A double-blind randomized clinical trial. Surgery, 2020, 167, 933-941. | 1.0 | 10 |
| 27 | <p>Can Immediately Treating Subcentimeter Hepatocellular Carcinoma Improve the Survival of Patients?</p> . Journal of Hepatocellular Carcinoma, 2020, Volume 7, 377-384. | 1.8 | 5 |
| 28 | Stereotactic Body Radiotherapy as a Salvage Therapy after Incomplete Radiofrequency Ablation for Hepatocellular Carcinoma: A Retrospective Propensity Score Matching Study. Cancers, 2019, 11, 1116. | 1.7 | 17 |
| 29 | Macrophages induce CD47 upregulation via IL-6 and correlate with poor survival in hepatocellular carcinoma patients. Oncolmmunology, 2019, 8, e1652540. | 2.1 | 55 |
| 30 | A nomogram predicting the recurrence of hepatocellular carcinoma in patients after laparoscopic hepatectomy. Cancer Communications, 2019, 39, 1-11. | 3.7 | 33 |
| 31 | Transarterial Chemoembolization Combined with Radiofrequency Ablation in the Treatment of Stage B1 Intermediate Hepatocellular Carcinoma. Journal of Oncology, 2019, 2019, 1-7. | 0.6 | 15 |
| 32 | Predictive factors for the benefit of tripleâ€drug transarterial chemoembolization for patients with unresectable hepatocellular carcinoma. Cancer Medicine, 2019, 8, 4200-4213. | 1.3 | 9 |
| 33 | 12-Month clinical results of drug-coated balloons for de novo coronary lesion in vessels exceeding 3.0Âmm. International Journal of Cardiovascular Imaging, 2019, 35, 579-586. | 0.7 | 13 |
| 34 | Vessels That Encapsulate Tumor Clusters (VETC) Pattern Is a Predictor of Sorafenib Benefit in Patients with Hepatocellular Carcinoma. Hepatology, 2019, 70, 824-839. | 3.6 | 62 |
| 35 | Robot-assisted laparoscopic partial hepatic caudate lobectomy. Minimally Invasive Therapy and Allied Technologies, 2019, 28, 292-297. | 0.6 | 7 |
| 36 | The Optimal Management for Sub-Centimeter Hepatocellular Carcinoma: Curative Treatments or Follow-Up?. Medical Science Monitor, 2019, 25, 4941-4951. | 0.5 | 5 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Comparison of new-generation drug-eluting stents versus drug-coated balloon for in-stent restenosis: a meta-analysis of randomised controlled trials. BMJ Open, 2018, 8, e017231. | 0.8 | 14 |
| 38 | Oneâ€year clinical outcomes and multislice computed tomography angiographic results following implantation of the <scp>N</scp> eo <scp>V</scp> as bioresorbable sirolimusâ€eluting scaffold in patients with single de novo coronary artery lesions. Catheterization and Cardiovascular Interventions, 2018, 91, 617-622. | 0.7 | 6 |
| 39 | Intravascular imaging in cardiovascular ageing. Experimental Gerontology, 2018, 109, 31-37. | 1.2 | 3 |
| 40 | Use of sodium nitroprusside in retrograde percutaneous coronary intervention for chronic total occlusion. Medicine (United States), 2018, 97, e11498. | 0.4 | 2 |
| 41 | Sorafenib Monotherapy Versus Sorafenib Combined with Regional Therapies for Hepatocellular Carcinoma Patients with Pulmonary Oligometastases: A Propensity Score-matched Analysis. Journal of Cancer, 2018, 9, 1745-1753. | 1.2 | 14 |
| 42 | Tumor Location Influences Oncologic Outcomes of Hepatocellular Carcinoma Patients Undergoing Radiofrequency Ablation. Cancers, 2018, 10, 378. | 1.7 | 36 |
| 43 | Implications of the local hemodynamic forces on the formation and destabilization of neoatherosclerotic lesions. International Journal of Cardiology, 2018, 272, 7-12. | 0.8 | 16 |
| 44 | Comparison of Stable and Unstable Ethiodized Oil Emulsions for Transarterial Chemoembolization of Hepatocellular Carcinoma:ÂResults of a Single-Center Double-BlindÂProspective Randomized ControlledÂTrial. Journal of Vascular and Interventional Radiology, 2018, 29, 1068-1077.e2. | 0.2 | 19 |
| 45 | Neointima and neoatherosclerotic characteristics in bare metal and first- and second-generation drug-eluting stents in patients admitted with cardiovascular events attributed to stent failure: an optical coherence tomography study. EuroIntervention, 2018, 13, e1831-e1840. | 1.4 | 13 |
| 46 | Invasive or non-invasive imaging for detecting high-risk coronary lesions?. Expert Review of Cardiovascular Therapy, 2017, 15, 165-179. | 0.6 | 15 |
| 47 | Optical coherence tomography assessment of a complex bifurcation lesion treated with double kissing Crush technique. Medicine (United States), 2017, 96, e5740. | 0.4 | 1 |
| 48 | A head to head comparison of XINSORB bioresorbable sirolimus-eluting scaffold versus metallic sirolimus-eluting stent: 180 days follow-up in a porcine model. International Journal of Cardiovascular Imaging, 2017, 33, 1473-1481. | 0.7 | 9 |
| 49 | Left jackknife position: a novel position for laparoscopic hepatectomy. Chinese Journal of Cancer, 2017, 36, 31. | 4.9 | 6 |
| 50 | Assessing Hepatic Fibrosis Using 2-D Shear Wave Elastography in Patients with Liver Tumors: A Prospective Single-Center Study. Ultrasound in Medicine and Biology, 2017, 43, 2522-2529. | 0.7 | 13 |
| 51 | Intravascular Ultrasound Classification of Plaque in Angiographic True Bifurcation Lesions of the Left Main Coronary Artery. Chinese Medical Journal, 2016, 129, 1538-1543. | 0.9 | 5 |
| 52 | Comparison between twoâ€dimensional and threeâ€dimensional quantitative coronary angiography for the prediction of functional severity in true bifurcation lesions: Insights from the randomized DKâ€CRUSH II, III, and IV trials. Catheterization and Cardiovascular Interventions, 2016, 87, 589-598. | 0.7 | 5 |
| 53 | Intravascular ultrasound-guided drug-eluting stent implantation: An updated meta-analysis of randomized control trials and observational studies. International Journal of Cardiology, 2016, 216, 133-139. | 0.8 | 73 |
| 54 | Derived neutrophil to lymphocyte ratio predicts prognosis for patients with HBV-associated hepatocellular carcinoma following transarterial chemoembolization. Oncology Letters, 2016, 11, 2987-2994. | 0.8 | 30 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Prospective, single-center cohort study analyzing the efficacy of complete laparoscopic resection on recurrent hepatocellular carcinoma. Chinese Journal of Cancer, 2016, 35, 25. | 4.9 | 37 |
| 56 | Vulnerable plaque detection: an unrealistic quest or a feasible objective with a clinical value?. Heart, 2016, 102, 581-589. | 1.2 | 27 |
| 57 | Expression of Pim-3 in colorectal cancer and its relationship with prognosis. Tumor Biology, 2016, 37, 9151-9156. | 0.8 | 14 |
| 58 | A randomized controlled trial on patients with or without adjuvant autologous cytokine-induced killer cells after curative resection for hepatocellular carcinoma. Oncolmmunology, 2016, 5, e1083671. | 2.1 | 56 |
| 59 | Noninvasive Prediction of Atherosclerotic Progression: The PROSPECT-MSCT Study. JACC: Cardiovascular Imaging, 2016, 9, 1009-1011. | 2.3 | 27 |
| 60 | The impact of dual antiplatelet therapy duration on primary composite endpoint after drug-eluting stent implantation: A meta-analysis of 10 randomized trials. International Journal of Cardiology, 2016, 202, 504-506. | 0.8 | 1 |
| 61 | Parecoxib prevents complications in hepatocellular carcinoma patients receiving hepatic transarterial chemoembolization: a prospective score-matched cohort study. Oncotarget, 2016, 7, 27938-27945. | 0.8 | 4 |
| 62 | Clinical and multimodality imaging results at 6 months of a bioresorbable sirolimus-eluting scaffold for patients with single de novo coronary artery lesions: the NeoVas first-in-man trial. EuroIntervention, 2016, 12, 1279-1287. | 1.4 | 16 |
| 63 | Evaluation of vascular healing of polymer-free sirolimus-eluting stents in native coronary artery stenosis: a serial follow-up at three and six months with optical coherence tomography imaging. EuroIntervention, 2016, 12, e574-e583. | 1.4 | 15 |
| 64 | Clinical outcomes of "complete, partially complete, and incomplete―revascularisation at five-year follow-up after percutaneous intervention of unprotected left main coronary artery disease with drug-eluting stents. EuroIntervention, 2016, 12, e957-e963. | 1.4 | 3 |
| 65 | Comparison of intravascular ultrasound guided versus angiography guided drug eluting stent implantation: a systematic review and meta-analysis. BMC Cardiovascular Disorders, 2015, 15, 153. | 0.7 | 81 |
| 66 | Impact of body mass index on longâ€ŧerm clinical outcomes after secondâ€generation drug eluting stent implantation: Insights from the international global <scp>RESOLUTE</scp> program. Catheterization and Cardiovascular Interventions, 2015, 85, 952-958. | 0.7 | 9 |
| 67 | Periprocedural myocardial infarction is associated with increased mortality in patients with coronary artery bifurcation lesions after implantation of a drugâ€eluting stent. Catheterization and Cardiovascular Interventions, 2015, 85, 696-705. | 0.7 | 14 |
| 68 | TACC3 promotes stemness and is a potential therapeutic target in hepatocellular carcinoma. Oncotarget, 2015, 6, 24163-24177. | 0.8 | 54 |
| 69 | Clinical Impact of Dual Antiplatelet Therapy Use in Patients Following Everolimus-eluting Stent Implantation. Chinese Medical Journal, 2015, 128, 714-720. | 0.9 | 3 |
| 70 | Impact of oral anti–hepatitis B therapy on the survival of patients with hepatocellular carcinoma initially treated with chemoembolization. Chinese Journal of Cancer, 2015, 34, 205-16. | 4.9 | 17 |
| 71 | Biolimus-eluting stent with biodegradable polymer improves clinical outcomes in patients with acute myocardial infarction. Heart, 2015, 101, 271-278. | 1.2 | 15 |
| 72 | Prognostic nomogram for patients with unresectable hepatocellular carcinoma after transcatheter arterial chemoembolization. Journal of Hepatology, 2015, 63, 122-130. | 1.8 | 101 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Optimal Medical Therapy Improves Clinical Outcomes in Patients Undergoing Revascularization With Percutaneous Coronary Intervention or Coronary Artery Bypass Grafting. Circulation, 2015, 131, 1269-1277. | 1.6 | 167 |
| 74 | Smoking Is Associated With Adverse Clinical Outcomes in PatientsÂUndergoing Revascularization With PCI or CABC. Journal of the American College of Cardiology, 2015, 65, 1107-1115. | 1.2 | 99 |
| 75 | Comparison of long-term in-stent vascular response between abluminal groove-filled biodegradable polymer sirolimus-eluting stent and durable polymer everolimus-eluting stent: 3-year OCT follow-up from the TARGET I trial. International Journal of Cardiovascular Imaging, 2015, 31, 1489-1496. | 0.7 | 7 |
| 76 | Long-term therapy with sorafenib is associated with pancreatic atrophy. Journal of Surgical Research, 2015, 199, 314-321. | 0.8 | 10 |
| 77 | Prognostic implications of severe coronary calcification in patients undergoing coronary artery bypass surgery: An analysis of the SYNTAX Study. Catheterization and Cardiovascular Interventions, 2015, 85, 199-206. | 0.7 | 32 |
| 78 | Early (before 6 months), late (6-12 months) and very late (after 12 months) angiographic scaffold restenosis in the ABSORB Cohort B trial. EuroIntervention, 2015, 10, 1288-1298. | 1.4 | 34 |
| 79 | Angiographic and clinical comparisons of intravascular ultrasound- versus angiography-guided drug-eluting stent implantation for patients with chronic total occlusion lesions: two-year results from a randomised AIR-CTO study. EuroIntervention, 2015, 10, 1409-1417. | 1.4 | 139 |
| 80 | Biodegradable or biocompatible polymer drug-eluting stent: a Gordian knot. EuroIntervention, 2015, 11, 250-252. | 1.4 | 2 |
| 81 | Circumferential distribution of the neointima at six-month and two-year follow-up after a bioresorbable vascular scaffold implantation: a substudy of the ABSORB Cohort B Clinical Trial. EuroIntervention, 2015, 10, 1299-1306. | 1.4 | 20 |
| 82 | Left atrial appendage closure monitoring without sedation: a pilot study using intracardiac echocardiography through the oesophageal route. EuroIntervention, 2015, 11, 936-941. | 1.4 | 29 |
| 83 | Bioresorbable vascular scaffold treatment induces the formation of neointimal cap that seals the underlying plaque without compromising the luminal dimensions: a concept based on serial optical coherence tomography data. EuroIntervention, 2015, 11, 746-756. | 1.4 | 42 |
| 84 | Comparison of one-year clinical outcomes between intravascular ultrasound-guided versus angiography-guided implantation of drug-eluting stents for left main lesions: a single-center analysis of a 1,016-patient cohort. Patient Preference and Adherence, 2014, 8, 1299. | 0.8 | 43 |
| 85 | Prognostic implications of coronary calcification in patients with obstructive coronary artery disease treated by percutaneous coronary intervention: a patient-level pooled analysis of 7 contemporary stent trials. Heart, 2014, 100, 1158-1164. | 1.2 | 216 |
| 86 | Mesenchymal stem cells with overexpression of midkine enhance cell survival and attenuate cardiac dysfunction in a rat model of myocardial infarction. Stem Cell Research and Therapy, 2014, 5, 37. | 2.4 | 35 |
| 87 | Response to Letter Regarding Article, "Quantification of Incomplete Revascularization and Its Association With Five-Year Mortality in the Synergy Between Percutaneous Coronary Intervention With Taxus and Cardiac Surgery (SYNTAX) Trial: Validation of the Residual SYNTAX Score― Circulation, 2014, 129, e355-6. | 1.6 | 1 |
| 88 | Scaffold and Edge Vascular Response Following Implantation of Everolimus-Eluting Bioresorbable Vascular Scaffold. JACC: Cardiovascular Interventions, 2014, 7, 1361-1369. | 1.1 | 23 |
| 89 | Berberine attenuates adverse left ventricular remodeling and cardiac dysfunction after acute myocardial infarction in rats: Role of autophagy. Clinical and Experimental Pharmacology and Physiology, 2014, 41, 995-1002. | 0.9 | 59 |
| 90 | The impact of everolimus versus other rapamycin derivative-eluting stents on clinical outcomes in patients with coronary artery disease: A meta-analysis of 16 randomized trials. Journal of Cardiology, 2014, 64, 185-193. | 0.8 | 15 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Short- and Long-Term Implications of a Bioresorbable Vascular Scaffold Implantation on the Local Endothelial Shear Stress Patterns. JACC: Cardiovascular Interventions, 2014, 7, 100-101. | 1.1 | 11 |
| 92 | Fusion of optical coherence tomographic and angiographic data for more accurate evaluation of the endothelial shear stress patterns and neointimal distribution after bioresorbable scaffold implantation: comparison with intravascular ultrasound-derived reconstructions. International Journal of Cardiovascular Imaging, 2014, 30, 485-494. | 0.7 | 37 |
| 93 | Implications of a bioresorbable vascular scaffold implantation on vessel wall strain of the treated and the adjacent segments. International Journal of Cardiovascular Imaging, 2014, 30, 477-484. | 0.7 | 3 |
| 94 | Effect of the Endothelial Shear Stress Patterns on Neointimal Proliferation Following Drug-Eluting Bioresorbable Vascular Scaffold Implantation. JACC: Cardiovascular Interventions, 2014, 7, 315-324. | 1.1 | 108 |
| 95 | Assessing Bioresorbable Coronary Devices. JACC: Cardiovascular Imaging, 2014, 7, 1130-1148. | 2.3 | 60 |
| 96 | Impact of the Complexity of Bifurcation Lesions Treated With Drug-Eluting Stents. JACC: Cardiovascular Interventions, 2014, 7, 1266-1276. | 1.1 | 153 |
| 97 | Prognostic Value of Site SYNTAX Score and Rationale for Combining Anatomic and Clinical Factors in Decision Making. Journal of the American College of Cardiology, 2014, 64, 423-432. | 1.2 | 48 |
| 98 | NOBORIâ,,¢ biodegradable-polymer biolimus-eluting stent versus durable-polymer drug-eluting stents: A meta-analysis. International Journal of Cardiology, 2014, 174, 151-153. | 0.8 | 2 |
| 99 | Berberine improves pressure overload-induced cardiac hypertrophy and dysfunction through enhanced autophagy. European Journal of Pharmacology, 2014, 728, 67-76. | 1.7 | 128 |
| 100 | Predicting 3-Year Mortality After Percutaneous Coronary Intervention. JACC: Cardiovascular Interventions, 2014, 7, 464-470. | 1.1 | 50 |
| 101 | Temporal Evolution of Strut Light Intensity After Implantation of Bioresorbable Polymeric Intracoronary Scaffolds in the ABSORB Cohort B Trial. Circulation Journal, 2014, 78, 1873-1881. | 0.7 | 12 |
| 102 | Predictive Performance of SYNTAX Score II in Patients With Left Main and Multivessel Coronary Artery Disease. Circulation Journal, 2014, 78, 1942-1949. | 0.7 | 64 |
| 103 | 68â€Predicting 3-Year Mortality After PCI: Revised Clinical Syntax Model Based on Patient Level Data from 7 Contemporary Stent Trials. Heart, 2014, 100, A38.2-A39. | 1.2 | Ο |
| 104 | Comparison of acute gain and late lumen loss after PCI with bioresorbable vascular scaffolds versus everolimus-eluting stents: an exploratory observational study prior to a randomised trial. EuroIntervention, 2014, 10, 672-680. | 1.4 | 30 |
| 105 | Stenting strategy for coronary artery bifurcation with drug-eluting stents: a meta-analysis of nine randomised trials and systematic review. EuroIntervention, 2014, 10, 561-569. | 1.4 | 66 |
| 106 | Bioresorbable scaffolds for coronary artery disease: current status and future prospective. Chinese Medical Journal, 2014, 127, 1141-8. | 0.9 | 1 |
| 107 | Nine-month angiographic and two-year clinical follow-up of polymer-free sirolimus-eluting stent versus durable-polymer sirolimus-eluting stent for coronary artery disease: the Nano randomized trial. Chinese Medical Journal, 2014, 127, 2153-8. | 0.9 | 14 |
| 108 | Impact of biodegradable versus durable polymer drug-eluting stents on clinical outcomes in patients with coronary artery disease: a meta-analysis of 15 randomized trials. Chinese Medical Journal, 2014, 127, 2159-66. | 0.9 | 5 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Assessment of plaque evolution in coronary bifurcations located beyond everolimus eluting scaffolds: serial intravascular ultrasound virtual histology study. Cardiovascular Ultrasound, 2013, 11, 25. | 0.5 | 7 |
| 110 | Progress in Treatment by Percutaneous Coronary Intervention: The Stent of the Future. Revista Espanola De Cardiologia (English Ed), 2013, 66, 483-496. | 0.4 | 34 |
| 111 | Short-Term and Long-Term Clinical Impact of Stent Thrombosis and Graft Occlusion in the SYNTAX Trial at 5 Years. Journal of the American College of Cardiology, 2013, 62, 2360-2369. | 1.2 | 62 |
| 112 | The Anatomic- and Clinical-Based NERS (New Risk Stratification) Score II to Predict Clinical Outcomes After Stenting Unprotected Left Main Coronary Artery Disease. JACC: Cardiovascular Interventions, 2013, 6, 1233-1241. | 1.1 | 24 |
| 113 | Intracoronary Optical Coherence Tomography and Histology of Overlapping Everolimus-Eluting Bioresorbable Vascular Scaffolds in a Porcine Coronary Artery Model. JACC: Cardiovascular Interventions, 2013, 6, 523-532. | 1.1 | 84 |
| 114 | Radial approach for patients with ST-segment elevation acute coronary syndrome: It is definitely the best access site. International Journal of Cardiology, 2013, 168, 3140-3142. | 0.8 | 0 |
| 115 | TCT-837 Everolimus-eluting versus other rapamycin derivatives-eluting stents in patients with coronary artery disease: a meta-analysis of 16 randomized trials. Journal of the American College of Cardiology, 2013, 62, B253. | 1.2 | 1 |
| 116 | Comparison of Intravascular Ultrasound Versus Angiography Guided Drug-eluting Stent Implantation: A Meta-analysis of Randomized Trials and Observational Studies Involving 28,745 Patients. American Journal of Cardiology, 2013, 111, 41B. | 0.7 | 0 |
| 117 | The Negative Impact of Incomplete Angiographic Revascularization on Clinical Outcomes and Its Association With Total Occlusions. Journal of the American College of Cardiology, 2013, 61, 282-294. | 1.2 | 257 |
| 118 | Bioresorbable scaffolds: Current knowledge, potentialities and limitations experienced during their first clinical applications. International Journal of Cardiology, 2013, 167, 11-21. | 0.8 | 56 |
| 119 | Clinical outcomes after zotarolimus and everolimus drug eluting stent implantation in coronary artery bifurcation lesions: insights from the RESOLUTE All Comers Trial. Heart, 2013, 99, 1267-1274. | 1.2 | 36 |
| 120 | Bioresorbable vascular scaffolds in the clinical setting. Interventional Cardiology, 2013, 5, 639-646. | 0.0 | 7 |
| 121 | Bioresorbable Drug-Eluting Magnesium-Alloy Scaffold for Treatment of Coronary Artery Disease. International Journal of Molecular Sciences, 2013, 14, 24492-24500. | 1.8 | 109 |
| 122 | The Ginsenoside Rg1 Prevents Transverse Aortic Constriction–Induced Left Ventricular Hypertrophy and Cardiac Dysfunction by Inhibiting Fibrosis and Enhancing Angiogenesis. Journal of Cardiovascular Pharmacology, 2013, 62, 50-57. | 0.8 | 60 |
| 123 | Quantification of Incomplete Revascularization and its Association With Five-Year Mortality in the Synergy Between Percutaneous Coronary Intervention With Taxus and Cardiac Surgery (SYNTAX) Trial Validation of the Residual SYNTAX Score. Circulation, 2013, 128, 141-151. | 1.6 | 326 |
| 124 | Intimal Flaps Detected by Optical Frequency Domain Imaging in the Proximal Segments of Native Coronary Arteries. Circulation Journal, 2013, 77, 2327-2333. | 0.7 | 10 |
| 125 | Bioresorbable scaffolds in the treatment of coronary artery disease. Medical Devices: Evidence and Research, 2013, 6, 37. | 0.4 | 17 |
| 126 | A randomised comparison of a novel abluminal groove-filled biodegradable polymer sirolimus-eluting stent with a durable polymer everolimus-eluting stent: clinical and angiographic follow-up of the TARGET I trial. EuroIntervention, 2013, 9, 75-83. | 1.4 | 60 |

YAO-JUN ZHANG

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | Percutaneous pulmonary artery denervation completely abolishes experimental pulmonary arterial hypertension in vivo. EuroIntervention, 2013, 9, 269-276. | 1.4 | 80 |
| 128 | In vivo assessment of the three-dimensional haemodynamic micro-environment following drug-eluting bioresorbable vascular scaffold implantation in a human coronary artery: fusion of frequency domain optical coherence tomography and angiography. EuroIntervention, 2013, 9, 890-890. | 1.4 | 17 |
| 129 | Revisiting: "Comparison of intravascular ultrasound versus angiography-guided drug-eluting stent implantation: a meta-analysis of one randomised trial and ten observational studies involving 19,619 patients― EuroIntervention, 2013, 9, 891-892. | 1.4 | 7 |
| 130 | Pulmonary arterial hypertension: pharmacologic therapies and potential pulmonary artery denervation treatment. EuroIntervention, 2013, 9, R149-R154. | 1.4 | 1 |
| 131 | Acute Effects of Nicardipine and Esmolol on The Cardiac Cycle, Intracardiac Hemodynamic and Endothelial Shear Stress in Patients With Unstable Angina Pectoris and Moderate Coronary Stenosis: Results From Single Center, Randomized Study. Cardiovascular Therapeutics, 2012, 30, 162-171. | 1.1 | 4 |
| 132 | TCT-78 Long-Term (4-Year) Clinical Outcomes of Total Occlusions and Completeness of Revascularisation in the Synergy between Percutaneous Coronary Intervention with Taxus and Cardiac Surgery Trial. Journal of the American College of Cardiology, 2012, 60, B25. | 1.2 | 0 |
| 133 | TCT-246 Comparison of intravascular ultrasound versus angiography guided drug-eluting stent implantation: a meta-analysis of randomized trials and observational studies involving 17,570 patients. Journal of the American College of Cardiology, 2012, 60, B70. | 1.2 | 0 |
| 134 | Bioresorbable Scaffolds: Current Evidence and Ongoing Clinical Trials. Current Cardiology Reports, 2012, 14, 626-634. | 1.3 | 33 |
| 135 | Angioscopy study from a large patient population comparing sirolimusâ€eluting stent with biodegradable versus durable polymer. Catheterization and Cardiovascular Interventions, 2012, 80, 420-428. | 0.7 | 10 |
| 136 | Five-year clinical follow-up of unprotected left main bifurcation lesion stenting: one-stent versus two-stent techniques versus double-kissing crush technique. EuroIntervention, 2012, 8, 803-814. | 1.4 | 40 |
| 137 | Comparison of intravascular ultrasound versus angiography-guided drug-eluting stent implantation: a meta-analysis of one randomised trial and ten observational studies involving 19,619 patients. EuroIntervention, 2012, 8, 855-865. | 1.4 | 163 |