Yuan Xu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1613007/publications.pdf

Version: 2024-02-01

501174 567247 44 896 15 28 citations h-index g-index papers 46 46 46 1679 docs citations citing authors all docs times ranked

#	Article	IF	CITATIONS
1	Association between decreased thyroid stimulating hormone and hyperuricemia in type 2 diabetic patients with early-stage diabetic kidney disease. BMC Endocrine Disorders, 2021, 21, 1.	2.2	39
2	A Wearable Gait Analysis System Used in Type 2 Diabetes Mellitus Patients: A Case–Control Study. Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2021, Volume 14, 1799-1808.	2.4	4
3	Abnormal apelin-ACE2 and SGLT2 signaling contribute to adverse cardiorenal injury in patients with COVID-19. International Journal of Cardiology, 2021, 336, 123-129.	1.7	11
4	Utilizing reclassification toâ€,exploreâ€,characteristics and prognosis of KDIGOSCr AKI subgroups: a retrospective analysis of a multicenter prospective cohort study. Renal Failure, 2021, 43, 1569-1576.	2.1	0
5	Search for clinical predictors of good glycemic control in patients starting or intensifying oral hypoglycemic pharmacological therapy: A multicenter prospective cohort study. Journal of Diabetes and Its Complications, 2020, 34, 107464.	2.3	4
6	Homocysteine Levels are Associated with Endothelial Function in Newly Diagnosed Type 2 Diabetes Mellitus Patients. Metabolic Syndrome and Related Disorders, 2019, 17, 323-327.	1.3	15
7	Effect of proximity to specialty care on outcomes for biliary cancers: a population-based retrospective cohort study. CMAJ Open, 2019, 7, E131-E139.	2.4	7
8	Comparison of therapeutic effects of acarbose and metformin under different \hat{l}^2 -cell function status in Chinese patients with type 2 diabetes. Endocrine Journal, 2019, 66, 443-450.	1.6	7
9	Role of irisin in Chinese patients with hypothyroidism: an interventional study. Journal of International Medical Research, 2019, 47, 1592-1601.	1.0	8
10	Diffuse Myocardial Injuries are Present in Subclinical Hypothyroidism: A Clinical Study Using Myocardial T1-mapping Quantification. Scientific Reports, 2018, 8, 4999.	3.3	11
11	Factors That Influence Pancreatic Beta Cell Function and Insulin Resistance in Newly Diagnosed Type 2 Diabetes Patients: A Sub-Analysis of the MARCH Trial. Diabetes Therapy, 2018, 9, 743-752.	2.5	8
12	Impact of hyperhomocysteinemia on insulin resistance in patients with H-type hypertension. Clinical and Experimental Hypertension, 2018, 40, 28-31.	1.3	8
13	Distinctive pattern of <scp>AHNAK</scp> methylation level in peripheral blood mononuclear cells and the association with <scp>HBV</scp> â€related liver diseases. Cancer Medicine, 2018, 7, 5178-5186.	2.8	9
14	Subthreshold depression among diabetes patients in Beijing: Cross-sectional associations among sociodemographic, clinical, and behavior factors. Journal of Affective Disorders, 2018, 237, 80-86.	4.1	9
15	The Effects of Exenatide and Metformin on Endothelial Function in Newly Diagnosed Type 2 Diabetes Mellitus Patients: A Case–Control Study. Diabetes Therapy, 2018, 9, 1295-1305.	2.5	14
16	Atorvastatin Decreased Circulating RANTES Levels in Impaired Glucose Tolerance Patients with Hypercholesterolemia: An Interventional Study. Diabetes Therapy, 2017, 8, 309-319.	2.5	3
17	PPAR-α Agonist Fenofibrate Reduces Insulin Resistance in Impaired Glucose Tolerance Patients with Hypertriglyceridemia: A Cross-Sectional Study. Diabetes Therapy, 2017, 8, 433-444.	2.5	8
18	Comparison of risk adjustment methods in patients with liver disease using electronic medical record data. BMC Gastroenterology, 2017, 17, 5.	2.0	13

#	Article	IF	Citations
19	Effects of shortâ€term levothyroxine therapy on myocardial injuries in patients with severe overt hypothyroidism: Evidence from a cardiac MRI Study. Journal of Magnetic Resonance Imaging, 2017, 46, 897-904.	3.4	5
20	Low apoA-I is associated with insulin resistance in patients with impaired glucose tolerance: a cross-sectional study. Lipids in Health and Disease, 2017, 16, 69.	3.0	20
21	Antithyroid Drug Therapy for Graves' Disease and Implications for Recurrence. International Journal of Endocrinology, 2017, 2017, 1-8.	1.5	25
22	Comparison of Exenatide and Metformin Monotherapy in Overweight/Obese Patients with Newly Diagnosed Type 2 Diabetes. International Journal of Endocrinology, 2017, 2017, 1-6.	1.5	8
23	Hyperhomocysteinemia as a Metabolic Risk Factor for Glucose Intolerance Among High-Risk Groups of Chinese Adults. Medical Science Monitor, 2017, 23, 2775-2781.	1.1	12
24	The effects of patient cost sharing on inpatient utilization, cost, and outcome. PLoS ONE, 2017, 12, e0187096.	2.5	5
25	Value of Kidney Disease Improving Global Outcomes Urine Output Criteria in Critically Ill Patients. Chinese Medical Journal, 2016, 129, 2050-2057.	2.3	15
26	Lung-protective Ventilation in Patients with Brain Injury. Chinese Medical Journal, 2016, 129, 1643-1651.	2.3	4
27	Levothyroxine treatment restored the decreased circulating fibroblast growth factor 21 levels in patients with hypothyroidism. European Journal of Internal Medicine, 2016, 31, 94-98.	2.2	11
28	Exenatide treatment increases serum irisin levels in patients with obesity and newly diagnosed type 2 diabetes. Journal of Diabetes and Its Complications, 2016, 30, 1555-1559.	2.3	34
29	Homocysteine diminishes apolipoprotein A-I function and expression in patients with hypothyroidism: a cross-sectional study. Lipids in Health and Disease, 2016, 15, 123.	3.0	9
30	Development and validation of method for defining conditions using Chinese electronic medical record. BMC Medical Informatics and Decision Making, 2016, 16, 110.	3.0	13
31	Effect of Metformin on Fibroblast Growth Factor-21 Levels in Patients with Newly Diagnosed Type 2 Diabetes. Diabetes Technology and Therapeutics, 2016, 18, 120-126.	4.4	15
32	High prevalence of vitamin D deficiency in urban health checkup population. Clinical Nutrition, 2016, 35, 859-863.	5.0	41
33	PPAR-α Agonist Fenofibrate Decreased RANTES Levels in Type 2 Diabetes Patients with Hypertriglyceridemia. Medical Science Monitor, 2016, 22, 743-751.	1.1	21
34	Clinical Study of Serum Homocysteine and Non-Alcoholic Fatty Liver Disease in Euglycemic Patients. Medical Science Monitor, 2016, 22, 4146-4151.	1.1	10
35	Native Magnetic Resonance T1-Mapping Identifies Diffuse Myocardial Injury in Hypothyroidism. PLoS ONE, 2016, 11, e0151266.	2.5	21
36	MEN1 c.825-1G>A mutation in a family with multiple endocrine neoplasia type 1: A case report. Molecular Medicine Reports, 2015, 12, 6152-6156.	2.4	3

#	Article	IF	CITATION
37	PPAR- $\langle i \rangle \hat{l} \pm \langle i \rangle$ Agonist Fenofibrate Decreased Serum Irisin Levels in Type 2 Diabetes Patients with Hypertriglyceridemia. PPAR Research, 2015, 2015, 1-8.	2.4	17
38	Resveratrol prevents hepatic steatosis and endoplasmic reticulum stress and regulates the expression of genes involved in lipid metabolism, insulin resistance, and inflammation in rats. Nutrition Research, 2015, 35, 576-584.	2.9	57
39	The role of fibroblast growth factor 21 in the pathogenesis of non-alcoholic fatty liver disease and implications for therapy. Metabolism: Clinical and Experimental, 2015, 64, 380-390.	3.4	96
40	Weaning critically ill patients from mechanical ventilation: A prospective cohort study. Journal of Critical Care, 2015, 30, 862.e7-862.e13.	2.2	31
41	Novel Clinical Evidence of an Association between Homocysteine and Insulin Resistance in Patients with Hypothyroidism or Subclinical Hypothyroidism. PLoS ONE, 2015, 10, e0125922.	2.5	44
42	Effect of obesity and hyperglycemia on benign prostatic hyperplasia in elderly patients with newly diagnosed type 2 diabetes. International Journal of Clinical and Experimental Medicine, 2015, 8, 11289-94.	1.3	15
43	Resveratrol Increases Nephrin and Podocin Expression and Alleviates Renal Damage in Rats Fed a High-Fat Diet. Nutrients, 2014, 6, 2619-2631.	4.1	39
44	Epidemiology and Outcome of Severe Sepsis and Septic Shock in Intensive Care Units in Mainland China. PLoS ONE, 2014, 9, e107181.	2.5	147