# Kenneth G Andersen

List of Publications by Year
in descending order



| | | | |
|---|---|---|---|
| 31 papers | 1,354 citations | 430874 18 h-index | 477307 29 g-index |
| 31 all docs | 31 docs citations | 31 times ranked | 1463 citing authors |

| # | Article | IF | Citations |
|---|---|---|---|
| 1 | Long-term results of a standardized enhanced recovery protocol in unilateral, secondary autologous breast reconstructions using an abdominal free flap. Journal of Plastic, Reconstructive and Aesthetic Surgery, 2022, 75, 1117-1122. | 1.0 | 7 |
| 2 | Use of Medical Supplies at the Roskilde Festival 2016: A Prospective Observational Study. Prehospital and Disaster Medicine, 2021, 36, 306-312. | 1.3 | 2 |
| 3 | Effect of progressive resistance training on persistent pain after axillary dissection in breast cancer: a randomized controlled trial. Breast Cancer Research and Treatment, 2020, 179, 173-183. | 2.5 | 20 |
| 4 | Use of Medication at the Roskilde (Denmark) Music Festival 2015 â€“ A Prospective Observational Study of 15,133 Treated Attendees. Prehospital and Disaster Medicine, 2019, 34, 407-414. | 1.3 | 4 |
| 5 | Development and Validation of a Screening Tool for Surgery-Specific Neuropathic Pain: Neuropathic Pain Scale for Postsurgical Patients. Pain Physician, 2019, 2, E81-E90. | 0.4 | 4 |
| 6 | Development and Validation of a Screening Tool for Surgery-Specific Neuropathic Pain: Neuropathic Pain Scale for Postsurgical Patients. Pain Physician, 2019, 22, E81-E90. | 0.4 | 6 |
| 7 | Analgesic and Sensory Effects of the Pecs Local Anesthetic Block in Patients with Persistent Pain after Breast Cancer Surgery: A Pilot Study. Pain Practice, 2017, 17, 185-191. | 1.9 | 17 |
| 8 | The Relationship Between Sensory Loss and Persistent Pain 1ÂYear After Breast Cancer Surgery. Journal of Pain, 2017, 18, 1129-1138. | 1.4 | 27 |
| 9 | Pain, sensory disturbances and psychological distress are common sequelae after treatment of ductal carcinoma in situ: a cross-sectional study. Acta OncolÃ³gica, 2017, 56, 724-729. | 1.8 | 19 |
| 10 | Pain, Sensory Disturbances, and Psychological Distress among Danish Women Treated for Ductal Carcinoma In Situ: An Exploratory Study. Pain Management Nursing, 2017, 18, 309-317. | 0.9 | 7 |
| 11 | The effects of individually tailored nurse navigation for patients with newly diagnosed breast cancer: a randomized pilot study. Acta OncolÃ³gica, 2017, 56, 1682-1689. | 1.8 | 40 |
| 12 | Clinical Prediction Model and Tool for Assessing Risk of Persistent Pain After Breast Cancer Surgery. Journal of Clinical Oncology, 2017, 35, 1660-1667. | 1.6 | 80 |
| 13 | Association between sensory dysfunction and pain 1 week after breast cancer surgery: a psychophysical study. Acta Anaesthesiologica Scandinavica, 2016, 60, 259-269. | 1.6 | 17 |
| 14 | Reply. Pain, 2016, 157, 1174. | 4.2 | 2 |
| 15 | Ultrasound Guided Intercostobrachial Nerve Blockade in Patients with Persistent Pain after Breast Cancer Surgery: A Pilot Study. Pain Physician, 2016, 19, E309-18. | 0.4 | 31 |
| 16 | Preoperative Distress Predicts Persistent Pain After Breast Cancer Treatment: A Prospective Cohort Study. Journal of the National Comprehensive Cancer Network: JNCCN, 2015, 13, 995-1003. | 4.9 | 37 |
| 17 | The Effect of Pain on Physical Functioning After Breast Cancer Treatment. Clinical Journal of Pain, 2015, 31, 794-802. | 1.9 | 25 |
| 18 | Predictive factors for the development of persistent pain after breast cancer surgery. Pain, 2015, 156, 2413-2422. | 4.2 | 158 |

| # | Article | IF | Citations |
|---|---|---|---|
| 19 | Test-retest Agreement and Reliability of Quantitative Sensory Testing 1 Year After Breast Cancer Surgery. Clinical Journal of Pain, 2015, 31, 393-403. | 1.9 | 17 |
| 20 | Intercostobrachial nerve handling and pain after axillary lymph node dissection for breast cancer. Acta Anaesthesiologica Scandinavica, 2014, 58, 1240-1248. | 1.6 | 46 |
| 21 | Neural Blockade for Persistent Pain After Breast Cancer Surgery. Regional Anesthesia and Pain Medicine, 2014, 39, 272-278. | 2.3 | 37 |
| 22 | Development in self-reported arm-lymphedema in Danish women treated for early-stage breast cancer in 2005 and 2006 â€“ A nationwide follow-up study. Breast, 2014, 23, 445-452. | 2.2 | 24 |
| 23 | Targeted Intraoperative Radiotherapy and Persistent Pain After Treatment. , 2014, , 85-91. | | 0 |
| 24 | Persistent pain, sensory disturbances and functional impairment after immediate or delayed axillary lymph node dissection. European Journal of Surgical Oncology, 2013, 39, 31-35. | 1.0 | 20 |
| 25 | Breast reconstruction with an expander prosthesis following mastectomy does not cause additional persistent pain: A nationwide cross-sectional study. Journal of Plastic, Reconstructive and Aesthetic Surgery, 2013, 66, 1652-1658. | 1.0 | 24 |
| 26 | Persistent pain and sensory disturbances after treatment for breast cancer: six year nationwide follow-up study. BMJ, The, 2013, 346, f1865-f1865. | 6.0 | 178 |
| 27 | Persistent pain, sensory disturbances and functional impairment after adjuvant chemotherapy for breast cancer. Scandinavian Journal of Pain, 2012, 3, 189-189. | 1.3 | 0 |
| 28 | Persistent pain, sensory disturbances and functional impairment after adjuvant chemotherapy for breast cancer: Cyclophosphamide, epirubicin and fluorouracil compared with docetaxel + epirubicin and cyclophosphamide. Acta OncolÃ³gica, 2012, 51, 1036-1044. | 1.8 | 39 |
| 29 | Persistent pain after targeted intraoperative radiotherapy (TARGIT) or external breast radiotherapy for breast cancer: A randomized trial. Breast, 2012, 21, 46-49. | 2.2 | 33 |
| 30 | Persistent Pain After Breast Cancer Treatment: A Critical Review of Risk Factors and Strategies for Prevention. Journal of Pain, 2011, 12, 725-746. | 1.4 | 408 |
| 31 | Pain following the repair of an abdominal hernia. Surgery Today, 2010, 40, 8-21. | 1.5 | 25 |