

Chaturong Pornrattanamaneewong

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1617368/publications.pdf>

Version: 2024-02-01

19
papers

206
citations

1163117

8
h-index

1058476

14
g-index

25
all docs

25
docs citations

25
times ranked

215
citing authors

#	ARTICLE	IF	CITATIONS
1	Temporary clamping of drain combined with tranexamic acid reduce blood loss after total knee arthroplasty: a prospective randomized controlled trial. BMC Musculoskeletal Disorders, 2012, 13, 124.	1.9	51
2	Three-hour interval drain clamping reduces postoperative bleeding in total knee arthroplasty: a prospective randomized controlled trial. Archives of Orthopaedic and Trauma Surgery, 2012, 132, 1059-1063.	2.4	34
3	A retrospective analysis of medial opening wedge high tibial osteotomy for varus osteoarthritic knee. Indian Journal of Orthopaedics, 2012, 46, 455-461.	1.1	20
4	Survivorship of medial opening wedge high tibial osteotomy in the elderly: two to ten years of follow up. International Orthopaedics, 2017, 41, 2045-2052.	1.9	20
5	Medial proximal tibial angle after medial opening wedge HTO: A retrospective diagnostic test study. Indian Journal of Orthopaedics, 2012, 46, 525-530.	1.1	17
6	Reasons for revision of failed hemiarthroplasty: Are there any differences between unipolar and bipolar?. European Journal of Orthopaedic Surgery and Traumatology, 2018, 28, 1117-1123.	1.4	10
7	Diaphragmatic height index: new diagnostic test for phrenic nerve dysfunction. Journal of Neurosurgery, 2012, 117, 890-896.	1.6	9
8	Risk factors of early periprosthetic femoral fracture after total knee arthroplasty. BMC Musculoskeletal Disorders, 2021, 22, 1009.	1.9	9
9	Modified Robert Jones bandage can not reduce invisible blood loss after total knee arthroplasty: a randomized-controlled trial. Archives of Orthopaedic and Trauma Surgery, 2018, 138, 1151-1157.	2.4	8
10	Blood loss and transfusion rate compared among different dosing regimens of tranexamic acid administration in patients undergoing hip hemiarthroplasty for femoral neck fracture: A randomized controlled trial. Injury, 2021, 52, 2986-2990.	1.7	7
11	Loss of correction after medial opening wedge high tibial osteotomy: a comparison of locking plates without bone grafts and non-locking compression plates with bone grafts. Journal of the Medical Association of Thailand = Chotmaihet Thangphaet, 2012, 95 Suppl 9, S21-8.	0.1	6
12	Revision primary total hip replacement: causes and risk factors. Journal of the Medical Association of Thailand = Chotmaihet Thangphaet, 2015, 98, 93-9.	0.1	4
13	Accuracy of empirical distal femoral valgus cut angle of 4° to 6° in total knee arthroplasty: a randomized controlled trial. European Journal of Orthopaedic Surgery and Traumatology, 2021, , 1.	1.4	2
14	Validity and Reliability of the Thai Version of the International Hip Outcome Tool in Young to Middle-Aged Physically Active Adults With Hip Disorders. Orthopaedic Journal of Sports Medicine, 2021, 9, 232596712110420.	1.7	2
15	Reoperation rate and implant survivorship compared between cementless and cemented bipolar hemiarthroplasty in femoral neck fracture patients with chronic kidney disease. Injury, 2021, , .	1.7	1
16	Early Intraprosthetic Dislocation After Closed Reduction in Modern Dual-mobility Total Hip Arthroplasty. Arthroplasty Today, 2022, 13, 89-92.	1.6	1
17	CUSTOMIZED GUIDE FOR FEMORAL COMPONENT POSITIONING IN HIP RESURFACING ARTHROPLASTY. Acta Ortopedica Brasileira, 2017, 25, 103-106.	0.5	0
18	Delayed Femoral Artery Occlusion After Total Hip Arthroplasty via the Posterior Approach: A Single-Case Report From Thailand. Arthroplasty Today, 2021, 11, 38-40.	1.6	0

#	ARTICLE	IF	CITATIONS
19	Author's reply. Indian Journal of Orthopaedics, 2013, 47, 215-6.	1.1	0