

List of Publications by Citations

Source: <https://exaly.com/author-pdf/162067/bi-qun-zou-publications-by-citations.pdf>
Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

35 papers	525 citations	16 h-index	21 g-index
35 ext. papers	684 ext. citations	4.2 avg, IF	3.98 L-index

#	Paper	IF	Citations
35	Strong in vitro and vivo cytotoxicity of novel organoplatinum(II) complexes with quinoline-coumarin derivatives. <i>European Journal of Medicinal Chemistry</i> , 2019 , 184, 111751	6.8	40
34	Synthesis of two platinum(II) complexes with 2-methyl-8-quinolinol derivatives as ligands and study of their antitumor activities. <i>European Journal of Medicinal Chemistry</i> , 2019 , 161, 334-342	6.8	39
33	In vitro and in vivo antitumor activities of three novel binuclear platinum(II) complexes with 4'-substituted-2,2':6',2''-terpyridine ligands. <i>European Journal of Medicinal Chemistry</i> , 2019 , 170, 195-202	6.8	37
32	Synthesis, crystal structure and biological evaluation of a new dasatinib copper(II) complex as telomerase inhibitor. <i>European Journal of Medicinal Chemistry</i> , 2018 , 143, 1597-1603	6.8	31
31	Novel tacrine platinum(II) complexes display high anticancer activity via inhibition of telomerase activity, dysfunction of mitochondria, and activation of the p53 signaling pathway. <i>European Journal of Medicinal Chemistry</i> , 2018 , 158, 106-122	6.8	29
30	Synthesis and antitumor mechanisms of two novel platinum(ii) complexes with 3-(2'-benzimidazolyl)-7-methoxycoumarin. <i>Metallomics</i> , 2018 , 10, 1160-1169	4.5	26
29	Two telomerase-targeting Pt(ii) complexes of jatrorrhizine and berberine derivatives induce apoptosis in human bladder tumor cells. <i>Dalton Transactions</i> , 2019 , 48, 15247-15254	4.3	25
28	Complexes of lanthanides(iii) with mixed 2,2'-bipyridyl and 5,7-dibromo-8-quinolinoline chelating ligands as a new class of promising anti-cancer agents. <i>Metallomics</i> , 2019 , 11, 1005-1015	4.5	24
27	Cytotoxicity, DNA binding and cell apoptosis induction of a zinc(II) complex of HBrQ. <i>MedChemComm</i> , 2015 , 6, 2224-2231	5	24
26	Synthesis and in vitro biological evaluation of three 4'-(4-methoxyphenyl)-2,2':6',2''-terpyridine iridium(III) complexes as new telomerase inhibitors. <i>European Journal of Medicinal Chemistry</i> , 2018 , 143, 1387-1395	6.8	21
25	Design, synthesis and pharmacological evaluation of new 3-(1H-benzimidazol-2-yl)quinolin-2(1H)-one derivatives as potential antitumor agents. <i>European Journal of Medicinal Chemistry</i> , 2018 , 157, 139-150	6.8	21
24	Cobalt(II) 8-hydroxyquinoline complexes: structure, cytotoxicity and action mechanism. <i>MedChemComm</i> , 2016 , 7, 806-812	5	21
23	Three novel transition metal complexes of 6-methyl-2-oxo-quinoline-3-carbaldehyde thiosemicarbazone: synthesis, crystal structure, cytotoxicity, and mechanism of action. <i>RSC Advances</i> , 2017 , 7, 17923-17933	3.7	20
22	High in vitro and in vivo antitumor activities of luminescent platinum(II) complexes with jatrorrhizine derivatives. <i>European Journal of Medicinal Chemistry</i> , 2019 , 183, 111727	6.8	20
21	Synthesis and antitumor mechanism of a new iron(iii) complex with 5,7-dichloro-2-methyl-8-quinolinol as ligands. <i>MedChemComm</i> , 2017 , 8, 633-639	5	17
20	Complexes of oxoplatin with rhein and ferulic acid ligands as platinum(iv) prodrugs with high anti-tumor activity. <i>Dalton Transactions</i> , 2020 , 49, 1613-1619	4.3	17
19	Studies on the structures, cytotoxicity and apoptosis mechanism of 8-hydroxylquinoline rhodium(III) complexes in T-24 cells. <i>New Journal of Chemistry</i> , 2016 , 40, 6005-6014	3.6	15

18	Synthesis, characterization and biological evaluation of six highly cytotoxic ruthenium(II) complexes with 4'-substituted-2,2':6',2''-terpyridine. <i>MedChemComm</i> , 2018 , 9, 525-533	5	13
17	High in vitro anticancer activity of a dinuclear palladium(II) complex with a 2-phenylpyridine ligand. <i>Inorganic Chemistry Communication</i> , 2018 , 96, 106-110	3.1	12
16	Tryptanthrin derivative copper(II) complexes with high antitumor activity by inhibiting telomerase activity, and inducing mitochondria-mediated apoptosis and S-phase arrest in BEL-7402. <i>New Journal of Chemistry</i> , 2018 , 42, 15479-15487	3.6	10
15	Novel Quinoline-based Ir(III) Complexes Exhibit High Antitumor Activity and. <i>ACS Medicinal Chemistry Letters</i> , 2019 , 10, 1614-1619	4.3	9
14	Highly cytotoxic, cyclometalated iridium(III)-5-fluoro-8-quinolinol complexes as cancer cell mitochondriotropic agents. <i>New Journal of Chemistry</i> , 2020 , 44, 7832-7837	3.6	7
13	3-(1H-benzimidazol-2-yl)-chromen-2-ylideneamine platinum(II) and ruthenium(II) complexes exert their high in vitro antitumor activity by inducing S-phase arrest and disrupting mitochondrial functions in SK-OV-3/DDP tumor cells. <i>Polyhedron</i> , 2019 , 157, 219-224	2.7	6
12	In vitro and in vivo activity of novel platinum(II) complexes with naphthalene imide derivatives inhibiting human non-small cell lung cancer cells. <i>New Journal of Chemistry</i> , 2019 , 43, 8146-8152	3.6	5
11	Two novel platinum(II) complexes with sorafenib and regorafenib: Synthesis, structural characterization, and evaluation of in vitro antitumor activity. <i>Inorganic Chemistry Communication</i> , 2019 , 104, 27-30	3.1	5
10	Platinum(II) complexes with rutaecarpine and tryptanthrin derivatives induce apoptosis by inhibiting telomerase activity and disrupting mitochondrial function. <i>MedChemComm</i> , 2018 , 9, 1639-1648	5	5
9	Complexes of platinum(II/IV) with 2-phenylpyridine derivatives as a new class of promising anti-cancer agents. <i>Inorganic Chemistry Communication</i> , 2019 , 108, 107510	3.1	4
8	Transition metal complexes with 6,7-dichloro-5,8-quinolinedione as mitochondria-targeted anticancer agents. <i>Polyhedron</i> , 2020 , 181, 114482	2.7	4
7	A 9-chloro-5,6,7,8-tetrahydroacridine Pt(II) complex induces apoptosis of Hep-G2 cells via inhibiting telomerase activity and disrupting mitochondrial pathway. <i>Inorganic Chemistry Communication</i> , 2019 , 99, 77-81	3.1	4
6	Inhibition of telomerase activity and SK-OV-3/DDP cell apoptosis by rhodium(III) and iron(III) complexes with 4?-(3-thiophenecarboxaldehyde)-2,2':6',2''-terpyridine. <i>Inorganic Chemistry Communication</i> , 2019 , 102, 180-184	3.1	3
5	Cyclometallated iridium(III)-5-bromo-8-quinolinol complexes as mitochondria-targeted anticancer agents. <i>Inorganic Chemistry Communication</i> , 2020 , 115, 107854	3.1	3
4	Synthesis, Characterization, and Cytotoxicity of the Cobalt (III) Complex with N,N-Diethyl-4-(2,2':6',2''-terpyridin-4'-yl)aniline. <i>Chemistry and Biodiversity</i> , 2018 , 15, e1800215	2.5	3
3	Preparation of platinum(II) complexes with naphthalene imide derivatives and exploration of their in vitro cytotoxic activities. <i>Inorganic Chemistry Communication</i> , 2019 , 104, 124-128	3.1	2
2	Synthesis, characterization and the anticancer activity of six lanthanides(III) complexes with 5,7-dihalogenated-8-quinolinol and 2,2'-bipyridine derivatives. <i>Transition Metal Chemistry</i> , 2020 , 45, 477-483	2.1	2
1	New 5-chloro-8-hydroxyquinoline derivatives organometallic Ru(II)-arene complexes as antitumor agents. <i>Inorganic Chemistry Communication</i> , 2019 , 108, 107537	3.1	1

