

L Lee Grismer

List of Publications by Citations

Source: <https://exaly.com/author-pdf/1639766/l-lee-grismer-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

159
papers

1,909
citations

22
h-index

32
g-index

175
ext. papers

2,383
ext. citations

1.3
avg, IF

5.21
L-index

#	Paper	IF	Citations
159	A Species-Level Phylogeny of Extant Snakes with Description of a New Colubrid Subfamily and Genus. <i>PLoS ONE</i> , 2016, 11, e0161070	3.7	131
158	A phylogeny and taxonomy of the Thai-Malay Peninsula Bent-toed Geckos of the <i>Cyrtodactylus pulchellus</i> complex (Squamata: Gekkonidae): combined morphological and molecular analyses with descriptions of seven new species. <i>Zootaxa</i> , 2012, 3520, 1	0.5	79
157	Systematics and natural history of Southeast Asian Rock Geckos (genus <i>Cnemaspis</i> Strauch, 1887) with descriptions of eight new species from Malaysia, Thailand, and Indonesia. <i>Zootaxa</i> , 2014, 3880, 1-147	0.5	63
156	Twelve new species of <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from isolated limestone habitats in east-central and southern Myanmar demonstrate high localized diversity and unprecedented microendemism. <i>Zoological Journal of the Linnean Society</i> , 2018, 182, 862-959	2.4	55
155	Phylogeography, geographic variation, and taxonomy of the Bent-toed Gecko <i>Cyrtodactylus quadrivirgatus</i> Taylor, 1962 from Peninsular Malaysia with the description of a new swamp dwelling species. <i>Zootaxa</i> , 2012, 3406, 39	0.5	37
154	Species delimitation with gene flow: A methodological comparison and population genomics approach to elucidate cryptic species boundaries in Malaysian Torrent Frogs. <i>Molecular Ecology</i> , 2017, 26, 5435-5450	5.7	35
153	A new species of karst forest Bent-toed Gecko (genus <i>Cyrtodactylus</i> Gray) not yet threatened by foreign cement companies and a summary of Peninsular Malaysia's endemic karst forest herpetofauna and the need for its conservation. <i>Zootaxa</i> , 2016, 4061, 1-17	0.5	35
152	The phylogenetic relationships of three new species of the <i>Cyrtodactylus pulchellus</i> complex (Squamata: Gekkonidae) from poorly explored regions in northeastern Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3786, 359-81	0.5	32
151	FOUR NEW SPECIES OF THE GEKKONID GENUS <i>CNEMASPIS</i> STRAUCH 1887 (REPTILIA: SQUAMATA) FROM SOUTHERN VIETNAM. <i>Herpetologica</i> , 2007, 63, 482-500	1.9	32
150	Evolutionary biogeography on Mexico's Baja California peninsula: A synthesis of molecules and historical geology. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 14017-8	11.5	32
149	Integrative taxonomy uncovers high levels of cryptic species diversity in <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) and the description of a new species from Peninsular Malaysia. <i>Zoological Journal of the Linnean Society</i> , 2013, 169, 849-880	2.4	29
148	Three more new species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from the Salween Basin of eastern Myanmar underscore the urgent need for the conservation of karst habitats. <i>Journal of Natural History</i> , 2018, 52, 1243-1294	0.5	27
147	The systematics and independent evolution of cave ecomorphology in distantly related clades of Bent-toed Geckos (Genus <i>Cyrtodactylus</i> Gray, 1827) from the Mekong Delta and islands in the Gulf of Thailand. <i>Zootaxa</i> , 2015, 3980, 106-26	0.5	26
146	<i>Cyrtodactylus sanook</i> (Squamata: Gekkonidae), a new cave-dwelling gecko from Chumphon Province, southern Thailand. <i>Zootaxa</i> , 2013, 3635, 275-85	0.5	26
145	Phylogeny and cryptic diversification in Southeast Asian flying geckos. <i>Molecular Phylogenetics and Evolution</i> , 2012, 65, 351-61	4.1	25
144	An integrative taxonomic review of the agamid genus <i>Bronchocela</i> (Kuhl, 1820) from Peninsular Malaysia with descriptions of new montane and insular endemics. <i>Zootaxa</i> , 2015, 3948, 1-23	0.5	24
143	A new <i>Cyrtodactylus</i> Gray, 1827 (Squamata, Gekkonidae) from the Shan Hills and the biogeography of Bent-toed Geckos from eastern Myanmar. <i>Zootaxa</i> , 2018, 4446, 477-500	0.5	24

142	Six new species of the <i>Cyrtodactylus intermedius</i> complex (Squamata: Gekkonidae) from the Cardamom Mountains and associated highlands of Southeast Asia. <i>Zootaxa</i> , 2019 , 4554, 1-62	0.5	23
141	Cyrt in the city: a new Bent-toed Gecko (Genus <i>Cyrtodactylus</i>) is the only endemic species of vertebrate from Batu Caves, Selangor, Peninsular Malaysia. <i>Zootaxa</i> , 2014 , 3774, 381-94	0.5	23
140	A new species of insular skink (Genus <i>Sphenomorphus</i> Fitzinger 1843) from the Langkawi Archipelago, Kedah, West Malaysia with the first report of the herpetofauna of Pulau Singa Besar and an updated checklist of the herpetofauna of Pulau Langkawi. <i>Zootaxa</i> , 2008 , 1691, 53	0.5	23
139	Multiple origins of parthenogenesis, and a revised species phylogeny for the Southeast Asian butterfly lizards, <i>Leiolepis</i> . <i>Biological Journal of the Linnean Society</i> , 2014 , 113, 1080-1093	1.9	22
138	Cryptic speciation within <i>Asthenodipsas vertebralis</i> (Boulenger, 1900) (Squamata: Pareatidae), the description of a new species from Peninsular Malaysia, and the resurrection of <i>A. tropidonotus</i> (Lidth de Jude, 1923) from Sumatra: an integrative taxonomic analysis. <i>Zootaxa</i> , 2013 , 3664, 505-24	0.5	22
137	A new species of karst dwelling gecko (genus <i>Cnemaspis</i> Strauch 1887) from the border region of Thailand and Peninsular Malaysia. <i>Zootaxa</i> , 2008 , 1875, 51	0.5	22
136	Two new Bent-toed Geckos of the <i>Cyrtodactylus pulchellus</i> complex from Peninsular Malaysia and multiple instances of convergent adaptation to limestone forest ecosystems. <i>Zootaxa</i> , 2016 , 4105, 401-295	0.5	22
135	A re-evaluation of the phylogenetic relationships of the <i>Cyrtodactylus condorensis</i> group (Squamata; Gekkonidae) and a suggested protocol for the characterization of rock-dwelling ecomorphology in <i>Cyrtodactylus</i> . <i>Zootaxa</i> , 2017 , 4300, 486	0.5	21
134	A new endemic species of <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from Tho Chu Island, southwestern Vietnam. <i>Zootaxa</i> , 2012 , 3228, 48	0.5	21
133	The Rediscovery of <i>Sphenomorphus malayanus</i> Doria, 1888 (Squamata: Scincidae) from the Titiwangsa Mountain Range of Peninsular Malaysia and its re-description as <i>S. senja</i> sp. nov. <i>Zootaxa</i> , 2015 , 3931, 63-70	0.5	20
132	A new species of <i>Cyrtodactylus</i> Gray, 1827 (Squamata: Gekkonidae) in Phu Quoc National Park, Kien Giang Biosphere Reserve, Southwestern Vietnam. <i>Zootaxa</i> , 2010 , 2604, 37	0.5	20
131	A new spiny, prehensile-tailed species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from Peninsular Malaysia with a preliminary hypothesis of relationships based on morphology. <i>Zootaxa</i> , 2010 , 2625, 40	0.5	20
130	Two New Species of Torrent-dwelling Toads <i>Ansonia</i> Stoliczka, 1870 (Anura: Bufonidae) from Peninsular Malaysia. <i>Herpetologica</i> , 2008 , 64, 321-340	1.9	20
129	Comprehensive multi-locus phylogeny of Old World tree frogs (Anura: Rhacophoridae) reveals taxonomic uncertainties and potential cases of over- and underestimation of species diversity. <i>Molecular Phylogenetics and Evolution</i> , 2018 , 127, 1010-1019	4.1	19
128	A new species of karst forest-adapted Bent-toed Gecko (genus <i>Cyrtodactylus</i> Gray, 1827) belonging to the <i>C. sworderi</i> complex from a threatened karst forest in Perak, Peninsular Malaysia. <i>Zootaxa</i> , 2014 , 3755, 434-46	0.5	19
127	The global diversity and distribution of lizard clutch sizes. <i>Global Ecology and Biogeography</i> , 2020 , 29, 1515-1530	6.1	17
126	A new species of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) from northwestern Thailand. <i>Zootaxa</i> , 2014 , 3760, 67-78	0.5	17
125	A new lowland forest Bent-toed Gecko (Squamata: Gekkonidae: <i>Cyrtodactylus</i>) from Ranong Province, peninsular Thailand. <i>Zootaxa</i> , 2015 , 3911, 106-18	0.5	16

124	A new species of <i>Ansonia</i> Stoliczka 1872 (Anura: Bufonidae) from Central Peninsular Malaysia and a revised taxonomy for <i>Ansonia</i> from the Malay Peninsula. <i>Zootaxa</i> , 2006 , 1327, 1	0.5	16
123	TWO NEW SPECIES OF <i>CNEMASPIS</i> STRAUCH, 1887 (SQUAMATA: GEKKONIDAE) FROM THE SERIBUAT ARCHIPELAGO, PAHANG AND JOHOR STATES, WEST MALAYSIA. <i>Herpetologica</i> , 2003 , 59, 544-552 ^{1.9}	1.9	16
122	Phylogenetic relationships of geckos of the <i>Hemiphyllodactylus harterti</i> group, a new species from Penang Island, Peninsular Malaysia, and a likely case of true cryptic speciation. <i>Zootaxa</i> , 2016 , 4107, 367-80 ^{0.5}	0.5	16
121	Repeated evolution of sympatric, palaeoendemic species in closely related, co-distributed lineages of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) across a sky-island archipelago in Peninsular Malaysia. <i>Zoological Journal of the Linnean Society</i> , 2015 , 174, 859-876	2.4	15
120	Evolution of habitat preference in 243 species of Bent-toed geckos (Genus Gray, 1827) with a discussion of karst habitat conservation. <i>Ecology and Evolution</i> , 2020 , 10, 13717-13730	2.8	15
119	Elevational size variation and two new species of torrent frogs from Peninsular Malaysia (Anura: Ranidae: Amolops Cope). <i>Zootaxa</i> , 2018 , 4434, 250-264	0.5	14
118	A diminutive new species of cave-dwelling Wolf Snake (Colubridae: <i>Lycodon</i> Boie, 1826) from Peninsular Malaysia. <i>Zootaxa</i> , 2014 , 51-67	0.5	14
117	First record of <i>Batrachochytrium dendrobatidis</i> infecting four frog families from Peninsular Malaysia. <i>EcoHealth</i> , 2011 , 8, 121-8	3.1	14
116	A new species of <i>Cnemaspis</i> Strauch 1887 (Squamata: Gekkonidae) from Selangor, Peninsular Malaysia. <i>Zootaxa</i> , 2008 , 1877, 49	0.5	14
115	Gene flow creates a mirage of cryptic species in a Southeast Asian spotted stream frog complex. <i>Molecular Ecology</i> , 2020 , 29, 3970-3987	5.7	14
114	A re-description of <i>Cyrtodactylus chrysopylos</i> Bauer (Squamata: Gekkonidae) with comments on the adaptive significance of orange coloration in hatchlings and descriptions of two new species from eastern Myanmar (Burma). <i>Zootaxa</i> , 2018 , 4527, 151-185	0.5	14
113	An updated and annotated checklist of the lizards of Peninsular Malaysia, Singapore, and their adjacent archipelagos. <i>Zootaxa</i> , 2019 , 4545, 230-248	0.5	13
112	Larger, unfiltered datasets are more effective at resolving phylogenetic conflict: Introns, exons, and UCEs resolve ambiguities in Golden-backed frogs (Anura: Ranidae; genus <i>Hylarana</i>). <i>Molecular Phylogenetics and Evolution</i> , 2020 , 151, 106899	4.1	13
111	Unrealized diversity in an urban rainforest: A new species of <i>Lygosoma</i> (Squamata: Scincidae) from western Sarawak, Malaysia (Borneo). <i>Zootaxa</i> , 2018 , 4370, 345-362	0.5	13
110	A new species of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) from Ba Na-Nui Chua Nature Reserve, Central Vietnam. <i>Zootaxa</i> , 2014 , 3760, 539-52	0.5	13
109	A new species of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) from Pulau Enggano, southwestern Sumatra, Indonesia. <i>Zootaxa</i> , 2014 , 3821, 485-95	0.5	13
108	A NEW SPECIES OF SMALL MONTANE FOREST FLOOR SKINK (GENUS SPHENOMORPHUS FITZINGER 1843) FROM SOUTHERN PENINSULAR MALAYSIA. <i>Herpetologica</i> , 2007 , 63, 544-551 ^{1.9}	1.9	13
107	Redescription <i>Cyrtodactylus lateralis</i> (Werner) (Squamata: Gekkonidae) and Phylogeny of the Prehensile-tailed <i>Cyrtodactylus</i> . <i>Zootaxa</i> , 2016 , 4107, 517-40	0.5	13

106	Renewed classification within Goniurosaurus (Squamata: Eublepharidae) uncovers the dual roles of a continental island (Hainan) in species evolution. <i>Molecular Phylogenetics and Evolution</i> , 2018 , 127, 646-654	4.1	12
105	The fourth Bent-toed Gecko of the genus Cyrtodactylus (Squamata: Gekkonidae) from Java, Indonesia. <i>Zootaxa</i> , 2015 , 4059, 351-63	0.5	12
104	A new species of insular Rock Gecko (Genus Cnemaspis Strauch, 1887) from the Bidong Archipelago, Terengganu, Peninsular Malaysia. <i>Zootaxa</i> , 2014 , 3755, 447-56	0.5	12
103	Peninsular Malaysia's first limbless lizard: a new species of skink of the genus Larutia (Borneo) from Pulau Pinang with a phylogeny of the genus. <i>Zootaxa</i> , 2011 , 2799, 29	0.5	12
102	The Eurasian hot nightlife: Environmental forces associated with nocturnality in lizards. <i>Global Ecology and Biogeography</i> , 2017 , 26, 1316-1325	6.1	11
101	Out of Borneo, again and again: biogeography of the Stream Toad genus Ansonia Stoliczka (Anura: Bufonidae) and the discovery of the first limestone cave-dwelling species. <i>Biological Journal of the Linnean Society</i> , 2016 ,	1.9	11
100	Description of a New Species of Goniurosaurus (Squamata: Eublepharidae) from the Guangdong Province, China, Based on Molecular and Morphological Data. <i>Herpetologica</i> , 2014 , 70, 309-322	1.9	11
99	A new insular species of Rock Gecko (Cnemaspis Boulenger) from Pulau Langkawi, Kedah, Peninsular Malaysia. <i>Zootaxa</i> , 2015 , 3985, 203-18	0.5	11
98	A phylogenetic taxonomy of the group (Reptilia: Squamata: Gekkonidae) with descriptions of two new species from Myanmar. <i>PeerJ</i> , 2018 , 6, e5575	3.1	11
97	Parachute geckos free fall into synonymy: Gekko phylogeny, and a new subgeneric classification, inferred from thousands of ultraconserved elements. <i>Molecular Phylogenetics and Evolution</i> , 2020 , 146, 106731	4.1	11
96	Two new species of Cyrtodactylus Gray, 1827 (Squamata: Gekkonidae) from a karstic archipelago in the Salween Basin of southern Myanmar (Burma). <i>Zootaxa</i> , 2020 , 4718, zootaxa.4718.2.1	0.5	11
95	Two more new species of Hemiphyllodactylus Bleeker (Squamata: Gekkonidae) from the Shan Hills of eastern Myanmar (Burma). <i>Zootaxa</i> , 2018 , 4483, 295-316	0.5	11
94	The discovery and description of a new species of Bent-toed Gecko of the Cyrtodactylus pulchellus complex (Squamata: Gekkonidae) from the Langkawi Archipelago, Kedah, Peninsular Malaysia. <i>Zootaxa</i> , 2019 , 4668, zootaxa.4668.1.3	0.5	10
93	Dragons in the mist: three new species of Pseudocalotes Fitzinger (Squamata: Agamidae) from the sky island archipelago of Peninsular Malaysia. <i>Zootaxa</i> , 2016 , 4136, 461-90	0.5	10
92	The phylogenetic position and taxonomic status of the Rainbow Tree Snake <i>Gonyophis margaritatus</i> (Peters, 1871) (Squamata: Colubridae). <i>Zootaxa</i> , 2014 , 3881, 532-48	0.5	10
91	A new species of karst-adapted Cnemaspis Strauch, 1887 (Squamata: Gekkonidae) from a threatened karst region in Pahang, Peninsular Malaysia. <i>Zootaxa</i> , 2013 , 3746, 463-72	0.5	10
90	Another new, diminutive Rock Gecko (Cnemaspis Strauch) from Peninsular Malaysia and a discussion of resource partitioning in sympatric species pairs. <i>Zootaxa</i> , 2010 , 2569, 55	0.5	10
89	A new species of Gastrophrynoidea (Anura: Microhylidae): an addition to a previously monotypic genus and a new genus for Peninsular Malaysia. <i>Zootaxa</i> , 2009 , 2124, 63-68	0.5	10

88	Three new karst-dwelling Strauch, 1887 (Squamata; Gekkonidae) from Peninsular Thailand and the phylogenetic placement of and. <i>PeerJ</i> , 2017 , 5, e2884	3.1	10
87	Karstic Landscapes Are Foci of Species Diversity in the World's Third-Largest Vertebrate Genus <i>Cyrtodactylus</i> Gray, 1827 (Reptilia: Squamata; Gekkonidae). <i>Diversity</i> , 2021 , 13, 183	2.5	10
86	Systematics, ecomorphology, cryptic speciation and biogeography of the lizard genus <i>Tytthoscincus</i> Linkem, Diesmos & Brown (Squamata: Scincidae) from the sky-island archipelago of Peninsular Malaysia. <i>Zoological Journal of the Linnean Society</i> , 2018 , 183, 635-671	2.4	9
85	A new species of Bent-toed Gecko (Squamata, Gekkonidae, Cyrtodactylus) from the Shan Plateau in eastern Myanmar (Burma). <i>Zootaxa</i> , 2019 , 4624, zootaxa.4624.3.1	0.5	9
84	A new species of lowland skink (genus <i>Lipinia</i> Gray, 1845) from northeastern Peninsular Malaysia. <i>Zootaxa</i> , 2014 , 3821, 457-64	0.5	9
83	Two new diminutive species of <i>Cnemaspis</i> Strauch 1887 (Squamata: Gekkonidae) from Peninsular Malaysia. <i>Zootaxa</i> , 2009 , 2019, 40-56	0.5	9
82	Phylogenetic partitioning of the third-largest vertebrate genus in the world, <i>Cyrtodactylus</i> Gray, 1827 (Reptilia; Squamata; Gekkonidae) and its relevance to taxonomy and conservation. <i>Vertebrate Zoology</i> , 71 , 101-154	2	9
81	The first riparian skink (Genus: <i>Sphenomorphus</i> Strauch, 1887) from Peninsular Malaysia and its relationship to other Indochinese and Sundaic species. <i>Zootaxa</i> , 2016 , 4173, 29-44	0.5	9
80	To split or not to split? Multilocus phylogeny and molecular species delimitation of southeast Asian toads (family: Bufonidae). <i>BMC Evolutionary Biology</i> , 2019 , 19, 95	3	8
79	Multilocus phylogeny and a new classification for African, Asian and Indian supple and writhing skinks (Scincidae: Lygosominae). <i>Zoological Journal of the Linnean Society</i> , 2019 , 186, 1067-1096	2.4	8
78	Two more new species of the <i>Cyrtodactylus peguensis</i> group (Squamata: Gekkonidae) from the fringes of the Ayeyarwady Basin, Myanmar. <i>Zootaxa</i> , 2019 , 4577, zootaxa.4577.2.3	0.5	8
77	Multilocus phylogeny of Bornean Bent-Toed geckos (Gekkonidae: Cyrtodactylus) reveals hidden diversity, taxonomic disarray, and novel biogeographic patterns. <i>Molecular Phylogenetics and Evolution</i> , 2020 , 147, 106785	4.1	8
76	Integrative taxonomy of the Asian skinks <i>Sphenomorphus stellatus</i> (Boulenger, 1900) and <i>S. praesignis</i> (Boulenger, 1900) with the resurrection of <i>S. annamiticus</i> (Boettger, 1901) and the description of a new species from Cambodia. <i>Zootaxa</i> , 2019 , 4683, zootaxa.4683.3.4	0.5	8
75	A new species of insular pitviper of the genus <i>Cryptelytrops</i> (Squamata: Viperidae) from southern Vietnam. <i>Zootaxa</i> , 2008 , 1715, 57	0.5	8
74	A revised and updated checklist of the lizards of Peninsular Malaysia. <i>Zootaxa</i> , 2008 , 1860, 28	0.5	8
73	A new species <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from western Thailand and the phylogenetic placement of <i>C. inthanon</i> and <i>C. doisuthep</i> . <i>Zootaxa</i> , 2020 , 4838, zootaxa.4838.2.2	0.5	8
72	Polyphyly of Asian Tree Toads, Genus <i>Pedostibes</i> Günther, 1876 (Anura: Bufonidae), and the Description of a New Genus from Southeast Asia. <i>PLOS ONE</i> , 2016 , 11, e0145903	3.7	8
71	The phylogenetic relationships of a new Stream Toad of the genus <i>Ansonia</i> Stoliczka, 1870 (Anura: Bufonidae) from a montane region in Peninsular Malaysia. <i>Zootaxa</i> , 2016 , 4103, 137-53	0.5	8

70	On the taxonomy and phylogeny of the skinks <i>Lipinia sekayuensis</i> Grismer, Ismail, Awang, Rizal, & Ahmad and <i>Lipinia surda</i> Boulenger from Peninsular Malaysia. <i>Zootaxa</i> , 2016 , 4147, 59-66	0.5	8
69	A new species of Crocodile Newt <i>Tylototriton</i> (Caudata: Salamandridae) from Shan State, Myanmar (Burma). <i>Zootaxa</i> , 2018 , 4500, 553-573	0.5	8
68	Phylogeny and biogeography of Bent-toed Geckos (<i>Cyrtodactylus</i> Gray) of the Sundaic swamp clade. <i>Zootaxa</i> , 2018 , 4472, 365-374	0.5	8
67	A new species of forest-dwelling <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from the Indawgyi Wildlife Sanctuary, Kachin State, Myanmar. <i>Zootaxa</i> , 2019 , 4623, zootaxa.4623.1.1	0.5	7
66	Four new Burmese species of <i>Hemiphyllodactylus</i> Bleeker (Squamata: Gekkonidae) from distantly related parapatric clades from the Shan Plateau and Salween Basin. <i>Zootaxa</i> , 2020 , 4758, zootaxa.4758.1.2	0.5	7
65	A new species of <i>Aplectana</i> (Nematoda, Cosmocercidae) in <i>Goniurosaurus bawanglingensis</i> (Squamata, Eublepharidae), from Hainan Province, China. <i>Acta Parasitologica</i> , 2018 , 63, 190-197	1.7	7
64	Phylogenetic taxonomy of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) with descriptions of three new species from Myanmar. <i>Journal of Natural History</i> , 2018 , 52, 881-915	0.5	7
63	A New Species of Frog (Amphibia: Anura: Ranidae) of the <i>Hylarana signata</i> Complex from Peninsular Malaysia. <i>Herpetologica</i> , 2014 , 70, 228-240	1.9	7
62	Phylogenetic relationships and description of a new upland species of Bent-toed Gecko (<i>Cyrtodactylus</i> Gray, 1827) of the <i>C. sworderi</i> complex from northeastern Peninsular Malaysia. <i>Zootaxa</i> , 2013 , 3616, 239-52	0.5	7
61	Target-capture phylogenomics provide insights on gene and species tree discordances in Old World treefrogs (Anura: Rhacophoridae). <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2020 , 287, 202024102	4.5	6
60	Nematodes of Five Species of Bufonids (Anura: Bufonidae) from Peninsular Malaysia. <i>Pacific Science</i> , 2017 , 71, 367-375	0.9	6
59	A new potentially endangered limestone-associated Bent-toed Gecko of the <i>Cyrtodactylus pulchellus</i> (Squamata: Gekkonidae) complex from northern Peninsular Malaysia. <i>Zootaxa</i> , 2020 , 4751, zootaxa.4751.3.2	0.5	6
58	Description of a cryptic new species of <i>Cnemaspis</i> Strauch, 1887 (Squamata: Gekkonidae) from the Western Ghats of Kerala State of India. <i>Zootaxa</i> , 2019 , 4656, zootaxa.4656.3.7	0.5	5
57	A new species of crocodile newt <i>Tylototriton</i> (Caudata: Salamandridae) from northern Myanmar (Burma). <i>Journal of Natural History</i> , 2019 , 53, 475-495	0.5	5
56	Helminths of Five Species of <i>Gonocephalus</i> Lizards (Squamata: Agamidae) from Peninsular Malaysia. <i>Pacific Science</i> , 2016 , 70, 373-380	0.9	5
55	Geographic structure of genetic variation in the Parachute Gecko <i>Ptychozoon lionotum</i> Annandale, 1905 across Indochina and Sundaland with descriptions of three new species. <i>Zootaxa</i> , 2019 , 4638, zootaxa.4638.2.1	0.5	5
54	Reappraisal of the Javanese Bullfrog complex, <i>Kaloula baleata</i> (Müller, 1836) (Amphibia: Anura: Microhylidae), reveals a new species from Peninsular Malaysia. <i>Zootaxa</i> , 2014 , 3900, 569-80	0.5	5
53	Gastrointestinal helminths of nine species of <i>Cnemaspis</i> (Squamata: Gekkonidae) from Peninsular Malaysia, one species from Cambodia and Thailand and two species from Vietnam. <i>Journal of Natural History</i> , 2015 , 49, 2683-2691	0.5	5

52	The transfer of two clades of Malaysian <i>Sphenomorphus</i> Fitzinger (Squamata: Scincidae) into the genus <i>Typhloscincus</i> Linkem, Diesmos, & Brown and the description of a new Malaysian swamp-dwelling species. <i>Zootaxa</i> , 2016 , 4092, 231-42	0.5	5
51	A new species of Mud Snake (Serpentes, Homalopsidae, <i>Gyiophis</i> Murphy & Voris, 2014) from Myanmar with a first molecular phylogenetic assessment of the genus. <i>Zootaxa</i> , 2017 , 4238, 571-582	0.5	4
50	New <i>Ichthyophis</i> species from Indochina (Gymnophiona, Ichthyophiidae): 1. The unstriped forms with descriptions of three new species and the redescriptions of <i>I. acuminatus</i> Taylor, 1960, <i>I. youngorum</i> Taylor, 1960 and <i>I. laosensis</i> Taylor, 1969. <i>Organisms Diversity and Evolution</i> , 2015 , 15, 143-174	1.7	4
49	Rivers of Indochina as potential drivers of lineage diversification in the spotted flying lizard (<i>Draco maculatus</i>) species complex. <i>Molecular Phylogenetics and Evolution</i> , 2020 , 150, 106861	4.1	4
48	A new skink of the genus <i>Subdoluseps</i> (Hardwicke amp; Gray, 1828) from Peninsular Malaysia. <i>Zootaxa</i> , 2019 , 4609, zootaxa.4609.2.10	0.5	4
47	<i>Cyrtodactylus rosichonariefi</i> sp. nov. (Squamata: Gekkonidae), a new swamp-dwelling bent-toed gecko from Bunguran Island (Great Natuna), Indonesia. <i>Zootaxa</i> , 2015 , 3964, 114-24	0.5	4
46	A new insular species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from the Langkawi Archipelago, Kedah, Peninsular Malaysia. <i>Zootaxa</i> , 2008 , 1924, 53-68	0.5	4
45	A new species of <i>Hemiphyllodactylus</i> Bleeker (Squamata; Gekkonidae) from Peninsular Thailand that converges in morphology and color pattern on <i>Pseudogekko smaragdinus</i> (Taylor) from the Philippines. <i>Zootaxa</i> , 2020 , 4816, zootaxa.4816.2.2	0.5	4
44	A new insular species of the <i>Cyrtodactylus intermedius</i> (Squamata: Gekkonidae) group from Cambodia with a discussion of habitat preference and ecomorphology. <i>Zootaxa</i> , 2020 , 4830, zootaxa.4830.1.3	0.5	4
43	Phylogenetic partitioning of the third-largest vertebrate genus in the world, <i>Cyrtodactylus</i> Gray, 1827 (Reptilia; Squamata; Gekkonidae) and its relevance to taxonomy and conservation. <i>Vertebrate Zoology</i> , 71 , 101-154	2	4
42	Sex Chromosome Turnover in Bent-Toed Geckos (). <i>Genes</i> , 2021 , 12,	4.2	4
41	Geographically structured genetic variation in <i>Ptychozoon lionotum</i> (Squamata: Gekkonidae) and a new species from an isolated volcano in Myanmar. <i>Zootaxa</i> , 2018 , 4514, 202-214	0.5	4
40	Gastrointestinal Helminths from Eight Species of <i>Cyrtodactylus</i> Lizards (Squamata: Gekkonidae) from Peninsular Malaysia and Two Species from Vietnam. <i>Pacific Science</i> , 2017 , 71, 91-98	0.9	3
39	Nematoda of Eleven Species of Ranid Frogs (Anura: Ranidae) from Southeast Asia. <i>Pacific Science</i> , 2017 , 71, 229-235	0.9	3
38	A taxonomic reappraisal of the Smooth Slug Snake <i>Asthenodipsas laevis</i> (Boie, 1827) (Squamata: Pareidae) in Borneo with the description of two new species. <i>Zootaxa</i> , 2019 , 4646, zootaxa.4646.3.4	0.5	3
37	A new insular, endemic frog of the genus <i>Kalophryndus</i> Tschudi, 1838 (Anura: Microhylidae) from Tioman Island, Pahang, Peninsular MalaysiaDepartment of Biology, La Sierra University, 4500 Riverwalk Parkway, Riverside, California, 92515-8247 USA.. <i>Zootaxa</i> , 2011 , 3123, 60	0.5	3
36	A definition of the group (Squamata, Eublepharidae) with the description of a new species. <i>ZooKeys</i> , 2020 , 986, 127-155	1.2	3
35	Origin, diversity, and conservation of karst-associated Bent-toed Geckos (Genus <i>Cyrtodactylus</i>) in Myanmar (Burma). <i>Israel Journal of Ecology and Evolution</i> , 2020 , 66, 202-208	0.8	3

34	A new species of <i>Cyrtodactylus</i> Gray (Squamata; Gekkonidae) from the Thai Highlands with a discussion on the evolution of habitat preference. <i>Zootaxa</i> , 2020 , 4852, zootaxa.4852.4.1	0.5	3
33	Taxonomy, phylogeny, and distribution of <i>Bronchocela rayaensis</i> (Squamata: Agamidae) on the Thai-Malay Peninsula. <i>Zootaxa</i> , 2016 , 4092, 414-20	0.5	3
32	Influence of niche breadth and position on the historical biogeography of seafaring scincid lizards. <i>Biological Journal of the Linnean Society</i> , 2021 , 132, 74-92	1.9	3
31	On the taxonomy of <i>Lygosoma bampfyldei</i> Bartlett, 1895 (Squamata: Scincidae) with descriptions of new species from Borneo and Peninsular Malaysia and the resurrection of <i>Lygosoma schneideri</i> Werner, 1900. <i>Zootaxa</i> , 2018 , 4438, 528-550	0.5	3
30	Nematodes of Five Species of Dicroidioid Frogs (Anura: Dicroidiidae) from Southeast Asia. <i>Pacific Science</i> , 2017 , 71, 535-540	0.9	2
29	New Supple Skink, Genus <i>Lygosoma</i> (Reptilia: Squamata: Scincidae), from Indochina and Redescription of <i>Lygosoma quadrupes</i> . <i>Journal of Herpetology</i> , 2018 , 52, 332-347	1.1	2
28	The Stoor Hobbit of Guangdong: sp. nov., a cave-dwelling Leopard Gecko (Squamata, Eublepharidae) from South China. <i>ZooKeys</i> , 2020 , 991, 137-153	1.2	2
27	Convergent evolution of karst habitat preference and its ecomorphological correlation in three species of Bent-toed Geckos (<i>Cyrtodactylus</i>) from Peninsular Malaysia. <i>Vertebrate Zoology</i> , 2016 , 71, 367-386	2	2
26	Archipelago-wide survey of Philippine forest dragons (Agamidae: <i>Gonocephalus</i>): multilocus phylogeny uncovers unprecedented levels of genetic diversity in a biodiversity hotspot. <i>Biological Journal of the Linnean Society</i> , 2016 ,	1.9	2
25	Two new <i>Typhloscincus</i> Linkem, Diesmos, Brown (Squamata: Scincidae) from Peninsular Malaysia and another case of microsyntopy between ecologically specialized, unrelated, leaf-litter species. <i>Zootaxa</i> , 2018 , 4425, 87-107	0.5	2
24	GroupStruct: An R Package for Allometric Size Correction.. <i>Zootaxa</i> , 2022 , 5124, 471-482	0.5	2
23	Discovery of the westernmost population of the genus <i>Ansonia</i> Stoliczka (Anura, Bufonidae) with the description of a new species from the Shan Plateau of eastern Myanmar. <i>Zootaxa</i> , 2019 , 4656, zootaxa.4656.3.11	0.5	1
22	Integrative taxonomic and geographic variation analyses in <i>Cyrtodactylus aequalis</i> (Squamata: Gekkonidae) from southern Myanmar (Burma): one species, two different stories. <i>Israel Journal of Ecology and Evolution</i> , 2020 , 66, 151-179	0.8	1
21	A taxonomic revision of <i>Asthenodipsas malaccana</i> Peters, 1864 (Squamata: Pareidae) with a description of a new species from Borneo. <i>Zootaxa</i> , 2020 , 4729, zootaxa.4729.1.1	0.5	1
20	Gastrointestinal Helminths of the Spotted Forest Skink, <i>Sphenomorphus scotophilus</i> (Squamata: Scincidae), from Peninsular Malaysia. <i>Comparative Parasitology</i> , 2018 , 85, 83-88	0.3	1
19	Three new species of the genus <i>Boulenophrys</i> (Anura, Megophryidae) from southern China.. <i>Zootaxa</i> , 2021 , 5072, 401-438	0.5	1
18	A new species of <i>Cnemaspis</i> Strauch (Squamata: Gekkonidae) of the <i>C. siamensis</i> group from Tenasserim Mountains, Thailand. <i>Zootaxa</i> , 2020 , 4852, zootaxa.4852.5.3	0.5	1
17	Taxonomic reassessment and phylogenetic placement of (Reptilia, Gekkonidae) based on morphological and molecular evidence. <i>ZooKeys</i> , 2021 , 1040, 91-121	1.2	1

16	First record of the <i>Cyrtodactylus brevipalmatus</i> group (Squamata: Gekkonidae) from Vietnam with description of a new species. <i>Zootaxa</i> , 2021 , 4969, 492510	0.5	1
15	On the taxonomy and phylogeny of the rare Selangor Mud Snake (<i>Raclitia indica</i>) Gray (Serpentes, Homalopsidae) from Peninsular Malaysia. <i>Zootaxa</i> , 2018 , 4514, 53-64	0.5	1
14	The rediscovery of Schaefer's Spine-jawed Snake (<i>Xenophidion schaeferi</i> GÜther Manthey, 1995) (Serpentes, Xenophidiidae) from Peninsular Malaysia with notes on its variation and the first record of the genus from Sumatra, Indonesia. <i>Zootaxa</i> , 2018 , 4441, 366-378	0.5	1
13	A standardized and statistically defensible framework for quantitative morphological analyses in taxonomic studies. <i>Zootaxa</i> , 2021 , 5023, 293-300	0.5	1
12	Another New Bent-toed Gecko, genus <i>Cyrtodactylus</i> Gray 1837 (Squamata: Gekkonidae), from Borneo. <i>Zootaxa</i> , 2021 , 5026, 286-300	0.5	1
11	The biogeography of bent-toed geckos, (Squamata: Gekkonidae).. <i>PeerJ</i> , 2022 , 10, e13153	3.1	1
10	Mountain endemism in Malesian geckos: can biotic interactions push lizards up hills?. <i>Israel Journal of Ecology and Evolution</i> , 2020 , 66, 190-201	0.8	0
9	Molecular phylogenetics, PCA, and MFA recover a new species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from an isolated sandstone massif in northwestern Cambodia. <i>Zootaxa</i> , 2021 , 4949, zootaxa.4949.2.3	0.5	0.3
8	Integrating spatial, phylogenetic, and threat assessment data from frogs and lizards to identify areas for conservation priorities in Peninsular Malaysia. <i>Global Ecology and Conservation</i> , 2021 , 28, e01650	2.8	0
7	The taxonomy and phylogeny of the <i>Cyrtodactylus brevipalmatus</i> group (Squamata: Gekkonidae) with emphasis on <i>C. interdigitalis</i> and <i>C. ngati</i> . <i>Vertebrate Zoology</i> , 72 , 245-269	2	0
6	A new species in the <i>Cyrtodactylus oldhami</i> group (Squamata, Gekkonidae) from Kanchanaburi Province, western Thailand. <i>ZooKeys</i> , 1103 , 139-169	1.2	0
5	A new species of <i>Sphenomorphus</i> (Squamata: Scincidae) from Phu Quoc Island, Vietnam with a discussion of biogeography and character state evolution in the <i>S. stellatus</i> group. <i>Zootaxa</i> , 2020 , 4801, zootaxa.4801.3.3	0.5	
4	The purportedly Indo-Australian gecko species <i>Cnemaspis timoriensis</i> (DumÉRÉ & Bibron, 1836) is actually the first named species of the neotropical genus <i>Gonatodes</i> Fitzinger, 1843 (Squamata: Gekkonidae). <i>Zootaxa</i> , 2019 , 4576, zootaxa.4576.3.4	0.5	
3	Herpetofauna of gunung panti forest reserve, johor, peninsular malaysia. <i>Tropical Life Sciences Research</i> , 2010 , 21, 71-82	1.1	
2	The phylogeny of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) with a description of a new species from the Mangin Range, Sagaing Region, northern Myanmar. <i>Journal of Natural History</i> , 2020 , 54, 1913-1931	0.5	
1	A new species of <i>Hemiphyllodactylus</i> Bleeker (Squamata: Gekkonidae) from Yunnan, China and its phylogenetic relationship to other congeners. <i>Zootaxa</i> , 2021 , 4980, 127	0.5	