

L Lee Grismer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1639766/publications.pdf>

Version: 2024-02-01

171
papers

3,036
citations

270111

25
h-index

355658

38
g-index

175
all docs

175
docs citations

175
times ranked

1575
citing authors

#	ARTICLE	IF	CITATIONS
1	The biogeography of bent-toed geckos, <i>Cyrtodactylus</i> (Squamata: Gekkonidae). PeerJ, 2022, 10, e13153.	0.9	11
2	GroupStruct: An R Package for Allometric Size Correction. Zootaxa, 2022, 5124, 471-482.	0.2	33
3	Cryptic extinction risk in a western Pacific lizard radiation. Biodiversity and Conservation, 2022, 31, 2045-2062.	1.2	8
4	Influence of niche breadth and position on the historical biogeography of seafaring scincid lizards. Biological Journal of the Linnean Society, 2021, 132, 74-92.	0.7	10
5	Sex Chromosome Turnover in Bent-Toed Geckos (<i>Cyrtodactylus</i>). Genes, 2021, 12, 116.	1.0	21
6	Molecular phylogenetics, PCA, and MFA recover a new species of <i>Cyrtodactylus</i> (Squamata: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 547 T zootaxa.4949.2.3.	0.2	5
7	Karstic Landscapes Are Foci of Species Diversity in the World's Third-Largest Vertebrate Genus <i>Cyrtodactylus</i> Gray, 1827 (Reptilia: Squamata; Gekkonidae). Diversity, 2021, 13, 183.	0.7	29
8	Taxonomic reassessment and phylogenetic placement of <i>Cyrtodactylus phuketensis</i> (Reptilia,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 462	0.5	3
9	A new species of <i>Hemiphyllodactylus</i> Bleeker (Squamata: Gekkonidae) from Yunnan, China and its phylogenetic relationship to other congeners. Zootaxa, 2021, 4980, 127.	0.2	4
10	First record of the <i>Cyrtodactylus brevipalmatus</i> group (Squamata: Gekkonidae) from Vietnam with description of a new species. Zootaxa, 2021, 4969, 492510.	0.2	7
11	A sheep in wolf's clothing: <i>Elaphe xiphodonta</i> sp. nov. (Squamata, Colubridae) and its possible mimicry to <i>Protobothrops jerdonii</i> . ZooKeys, 2021, 1048, 23-47.	0.5	4
12	A standardized and statistically defensible framework for quantitative morphological analyses in taxonomic studies. Zootaxa, 2021, 5023, 293-300.	0.2	19
13	Integrating spatial, phylogenetic, and threat assessment data from frogs and lizards to identify areas for conservation priorities in Peninsular Malaysia. Global Ecology and Conservation, 2021, 28, e01650.	1.0	3
14	Another New Bent-toed Gecko, genus <i>Cyrtodactylus</i> Gray 1837 (Squamata: Gekkonidae), from Borneo. Zootaxa, 2021, 5026, 286-300.	0.2	3
15	Three new species of the genus <i>Boulenophrys</i> (Anura, Megophryidae) from southern China. Zootaxa, 2021, 5072, 401-438.	0.2	5
16	Parachute geckos free fall into synonymy: <i>Gekko</i> phylogeny, and a new subgeneric classification, inferred from thousands of ultraconserved elements. Molecular Phylogenetics and Evolution, 2020, 146, 106731.	1.2	31
17	Two new species of <i>Cyrtodactylus</i> Gray, 1827 (Squamata: Gekkonidae) from a karstic archipelago in the Salween Basin of southern Myanmar (Burma). Zootaxa, 2020, 4718, zootaxa.4718.2.1.	0.2	16
18	A new species of <i>Hemiphyllodactylus</i> Bleeker (Squamata; Gekkonidae) from Peninsular Thailand that converges in morphology and color pattern on <i>Pseudogekko smaragdinus</i> (Taylor) from the Philippines. Zootaxa, 2020, 4816, zootaxa.4816.2.2.	0.2	9

#	ARTICLE	IF	CITATIONS
19	Evolution of habitat preference in 243 species of Bent-toed geckos (Genus <i>Cyrtodactylus</i> Gray,) Tj ETQq1 1.0784314 rgBT / Overlock 10 Tf 50	0.8	33
20	Origin, diversity, and conservation of karst-associated Bent-toed Geckos (Genus <i>Cyrtodactylus</i>) in Myanmar (Burma). <i>Israel Journal of Ecology and Evolution</i> , 2020, 66, 202-208.	0.2	3
21	A new species of <i>Cnemaspis</i> Strauch (Squamata: Gekkonidae) of the <i>C. siamensis</i> group from Tenasserim Mountains, Thailand. <i>Zootaxa</i> , 2020, 4852, zootaxa.4852.5.3.	0.2	5
22	A new species of <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from western Thailand and the phylogenetic placement of <i>C. inthanon</i> and <i>C. doisuthep</i> . <i>Zootaxa</i> , 2020, 4838, 179-209.	0.2	9
23	A new species of <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from the Thai Highlands with a discussion on the evolution of habitat preference. <i>Zootaxa</i> , 2020, 4852, 401-427.	0.2	6
24	Gene flow creates a mirage of cryptic species in a Southeast Asian spotted stream frog complex. <i>Molecular Ecology</i> , 2020, 29, 3970-3987.	2.0	49
25	A new insular species of the <i>Cyrtodactylus intermedius</i> (Squamata:) Tj ETQq1 1.0784314 rgBT / Overlock 10 Tf 50 ecomorphology. <i>Zootaxa</i> , 2020, 4830, 75-102.	0.2	11
26	Target-capture phylogenomics provide insights on gene and species tree discordances in Old World treefrogs (Anura: Rhacophoridae). <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2020, 287, 20202102.	1.2	17
27	Integrative taxonomic and geographic variation analyses in <i>Cyrtodactylus aequalis</i> (Squamata:) Tj ETQq1 1.0784314 rgBT / Overlock 10 Tf 50 Ecology and Evolution, 2020, 66, 151-179.	0.2	5
28	Rivers of Indochina as potential drivers of lineage diversification in the spotted flying lizard (<i>Draco</i>) Tj ETQq0 0.0 rgBT / Overlock 10 Tf 50	1.2	10
29	The global diversity and distribution of lizard clutch sizes. <i>Global Ecology and Biogeography</i> , 2020, 29, 1515-1530.	2.7	49
30	Larger, unfiltered datasets are more effective at resolving phylogenetic conflict: Introns, exons, and UCEs resolve ambiguities in Golden-backed frogs (Anura: Ranidae; genus <i>Hylarana</i>). <i>Molecular Phylogenetics and Evolution</i> , 2020, 151, 106899.	1.2	30
31	A new species of <i>Sphenomorphus</i> (Squamata: Scincidae) from Phu Quoc Island, Vietnam with a discussion of biogeography and character state evolution in the <i>S. stellatus</i> group. <i>Zootaxa</i> , 2020, 4801, zootaxa.4801.3.3.	0.2	3
32	A taxonomic revision of <i>Asthenodipsas malaccana</i> Peters, 1864 (Squamata: Pareidae) with a description of a new species from Borneo. <i>Zootaxa</i> , 2020, 4729, 1-24.	0.2	3
33	Four new Burmese species of <i>Hemiphyllodactylus</i> Bleeker (Squamata: Gekkonidae) from distantly related parapatric clades from the Shan Plateau and Salween Basin. <i>Zootaxa</i> , 2020, 4758, zootaxa.4758.1.2.	0.2	12
34	Multilocus phylogeny of Bornean Bent-Toed geckos (Gekkonidae: <i>Cyrtodactylus</i>) reveals hidden diversity, taxonomic disarray, and novel biogeographic patterns. <i>Molecular Phylogenetics and Evolution</i> , 2020, 147, 106785.	1.2	13
35	A new potentially endangered limestone-associated Bent-toed Gecko of the <i>Cyrtodactylus pulchellus</i> (Squamata: Gekkonidae) complex from northern Peninsular Malaysia. <i>Zootaxa</i> , 2020, 4751, zootaxa.4751.3.2.	0.2	8
36	A definition of the <i>Goniurosaurus yingdeensis</i> group (Squamata, Eublepharidae) with the description of a new species. <i>ZooKeys</i> , 2020, 986, 127-155.	0.5	8

#	ARTICLE	IF	CITATIONS
37	The Stoor Hobbit of Guangdong: <i>Goniurosaurus gollum</i> sp. nov., a cave-dwelling Leopard Gecko (Squamata, Eublepharidae) from South China. <i>ZooKeys</i> , 2020, 991, 137-153.	0.5	6
38	The phylogeny of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) with a description of a new species from the Mangin Range, Sagaing Region, northern Myanmar. <i>Journal of Natural History</i> , 2020, 54, 1913-1931.	0.2	2
39	Mountain endemism in Malesian geckos: can biotic interactions push lizards up hills?. <i>Israel Journal of Ecology and Evolution</i> , 2020, 66, 190-201.	0.2	3
40	Geographic structure of genetic variation in the Parachute Gecko <i>Ptychozoon lionotum</i> Annandale, 1905 across Indochina and Sundaland with descriptions of three new species. <i>Zootaxa</i> , 2019, 4638, 151-198.	0.2	9
41	A taxonomic reappraisal of the Smooth Slug Snake <i>Asthenodipsas laevis</i> (Boie, 1827) (Squamata: Tj ETQq1 1 0.784314 rgBT/Overlo	0.2	7
42	A new skink of the genus <i>Subdoluseps</i> (Hardwicke & Gray, 1828) from Peninsular Malaysia. <i>Zootaxa</i> , 2019, 4609, zootaxa.4609.2.10.	0.2	7
43	The purportedly Indo-Australian gecko species <i>Cnemaspis timoriensis</i> (DumÃ©nil & Bibron, 1836) is actually the first named species of the neotropical genus <i>Gonatodes</i> Fitzinger, 1843 (Squamata: Tj ETQq1 1 0.784314 rgBT/Overlo	0.2	14
44	A new species of Bent-toed Gecko (Squamata, Gekkonidae, <i>Cyrtodactylus</i>) from the Shan Plateau in eastern Myanmar (Burma). <i>Zootaxa</i> , 2019, 4624, zootaxa.4624.3.1.	0.2	11
45	Integrative taxonomy of the Asian skinks <i>Sphenomorphus stellatus</i> (Boulenger, 1900) and <i>S. praesignis</i> (Boulenger, 1900) with the resurrection of <i>S. annamiticus</i> (Boettger, 1901) and the description of a new species from Cambodia. <i>Zootaxa</i> , 2019, 4683, zootaxa.4683.3.4.	0.2	14
46	Discovery of the westernmost population of the genus <i>Ansonia</i> (Anura, Bufonidae) with the description of a new species from the Shan Plateau of eastern Myanmar. <i>Zootaxa</i> , 2019, 4656, 545-571.	0.2	4
47	A new species of forest-dwelling <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from the Indawgyi Wildlife Sanctuary, Kachin State, Myanmar. <i>Zootaxa</i> , 2019, 4623, 1-25.	0.2	10
48	Description of a cryptic new species of <i>Cnemaspis</i> Strauch, 1887 (Squamata: Gekkonidae) from the Western Ghats of Kerala State of India. <i>Zootaxa</i> , 2019, 4656, zootaxa.4656.3.7.	0.2	8
49	The discovery and description of a new species of Bent-toed Gecko of the <i>Cyrtodactylus pulchellus</i> complex (Squamata: Tj ETQq1 1 0.784314 rgBT/Overlo	0.2	12
50	An updated and annotated checklist of the lizards of Peninsular Malaysia, Singapore, and their adjacent archipelagos. <i>Zootaxa</i> , 2019, 4545, 230-248.	0.2	21
51	A new species of crocodile newt <i>Tylototriton</i> (Caudata: Salamandridae) from northern Myanmar (Burma). <i>Journal of Natural History</i> , 2019, 53, 475-495.	0.2	10
52	To split or not to split? Multilocus phylogeny and molecular species delimitation of southeast Asian toads (family: Bufonidae). <i>BMC Evolutionary Biology</i> , 2019, 19, 95.	3.2	16
53	Multilocus phylogeny and a new classification for African, Asian and Indian supple and writhing skinks (Scincidae: Lygosominae). <i>Zoological Journal of the Linnean Society</i> , 2019, 186, 1067-1096.	1.0	15
54	Two more new species of the <i>Cyrtodactylus peguensis</i> group (Squamata: Gekkonidae) from the fringes of the Ayeyarwady Basin, Myanmar. <i>Zootaxa</i> , 2019, 4577, 274.	0.2	11

#	ARTICLE	IF	CITATIONS
55	Six new species of the <i>Cyrtodactylus intermedius</i> complex (Squamata: Gekkonidae) from the Cardamom Mountains and associated highlands of Southeast Asia. <i>Zootaxa</i> , 2019, 4554, 1-62.	0.2	31
56	ENDOPARASITES IN THE THREE-BANDED LARUT SKINK, <i>LARUTIA TRIFASCIATA</i> (SQUAMATA: SCINCIDAE) FROM PENINSULAR MALAYSIA. <i>Texas Journal of Science</i> , 2019, 71, .	0.3	0
57	Three more new species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from the Salween Basin of eastern Myanmar underscore the urgent need for the conservation of karst habitats. <i>Journal of Natural History</i> , 2018, 52, 1243-1294.	0.2	35
58	Twelve new species of <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from isolated limestone habitats in east-central and southern Myanmar demonstrate high localized diversity and unprecedented microendemism. <i>Zoological Journal of the Linnean Society</i> , 2018, 182, 862-959.	1.0	71
59	A new species of <i>Aplectana</i> (Nematoda, Cosmocercidae) in <i>Goniurosaurus bawanglingensis</i> (Squamata), <i>Tj ETQq1 1,0,784314 rgBT /Ov</i>	0.4	10
60	Systematics, ecomorphology, cryptic speciation and biogeography of the lizard genus <i>Tytthoscincus</i> Linkem, Diesmos & Brown (Squamata: Scincidae) from the sky-island archipelago of Peninsular Malaysia. <i>Zoological Journal of the Linnean Society</i> , 2018, 183, 635-671.	1.0	18
61	Phylogenetic taxonomy of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) with descriptions of three new species from Myanmar. <i>Journal of Natural History</i> , 2018, 52, 881-915.	0.2	14
62	On the taxonomy and phylogeny of the rare Selangor Mud Snake (<i>Raclitia indica</i>) Gray (Serpentes), <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50</i>	0.2	4
63	Geographically structured genetic variation in <i>Ptychozoon lionotum</i> (Squamata: Gekkonidae) and a new species from an isolated volcano in Myanmar. <i>Zootaxa</i> , 2018, 4514, 202.	0.2	5
64	A re-description of <i>Cyrtodactylus chrysopylos</i> Bauer (Squamata: Gekkonidae) with comments on the adaptive significance of orange coloration in hatchlings and descriptions of two new species from eastern Myanmar (Burma). <i>Zootaxa</i> , 2018, 4527, 151.	0.2	16
65	A new species of Crocodile Newt <i>Tylostotriton</i> (Caudata: Salamandridae) from Shan State, Myanmar (Burma). <i>Zootaxa</i> , 2018, 4500, 553-573.	0.2	13
66	Two more new species of <i>Hemiphyllodactylus</i> Bleeker (Squamata: Gekkonidae) from the Shan Hills of eastern Myanmar (Burma). <i>Zootaxa</i> , 2018, 4483, 295-316.	0.2	16
67	Phylogeny and biogeography of Bent-toed Geckos (<i>Cyrtodactylus</i> Gray) of the Sundaic swamp clade. <i>Zootaxa</i> , 2018, 4472, 365.	0.2	12
68	On the taxonomy of <i>Lygosoma bampfyldei</i> Bartlett, 1895 (Squamata: Scincidae) with descriptions of new species from Borneo and Peninsular Malaysia and the resurrection of <i>Lygosoma schneideri</i> Werner, 1900. <i>Zootaxa</i> , 2018, 4438, 528.	0.2	7
69	Two new <i>Tytthoscincus</i> Linkem, Diesmos, & Brown (Squamata; Scincidae) from Peninsular Malaysia and another case of microsyntopy between ecologically specialized, unrelated, leaf-litter species. <i>Zootaxa</i> , 2018, 4425, 87.	0.2	5
70	The rediscovery of Schaefer's Spine-jawed Snake (<i>Xenophidion schaeferi</i> Günther & Manthey), <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50</i> record of the genus from Sumatra, Indonesia. <i>Zootaxa</i> , 2018, 4441, 366-378.	0.2	5
71	Unrealized diversity in an urban rainforest: A new species of <i>Lygosoma</i> (Squamata: Scincidae) from western Sarawak, Malaysia (Borneo). <i>Zootaxa</i> , 2018, 4370, 345-362.	0.2	15
72	Comprehensive multi-locus phylogeny of Old World tree frogs (Anura: Rhacophoridae) reveals taxonomic uncertainties and potential cases of over- and underestimation of species diversity. <i>Molecular Phylogenetics and Evolution</i> , 2018, 127, 1010-1019.	1.2	33

#	ARTICLE	IF	CITATIONS
73	A new <i>Cyrtodactylus</i> Gray, 1827 (Squamata, Gekkonidae) from the Shan Hills and the biogeography of Bent-toed Geckos from eastern Myanmar. <i>Zootaxa</i> , 2018, 4446, 477-500.	0.2	27
74	Elevational size variation and two new species of torrent frogs from Peninsular Malaysia (Anura: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 7	0.2	24
75	Gastrointestinal Helminths of the Spotted Forest Skink, <i>Sphenomorphus scotophilus</i> (Squamata: Tj ETQq1 1 0.784314 rgBT /Overlock 1	0.0	1
76	New Supple Skink, Genus <i>Lygosoma</i> (Reptilia: Squamata: Scincidae), from Indochina and Redescription of <i>Lygosoma quadrupes</i> (). <i>Journal of Herpetology</i> , 2018, 52, 332-347.	0.2	8
77	Renewed classification within <i>Goniurosaurus</i> (Squamata: Eublepharidae) uncovers the dual roles of a continental island (Hainan) in species evolution. <i>Molecular Phylogenetics and Evolution</i> , 2018, 127, 646-654.	1.2	19
78	A phylogenetic taxonomy of the <i>Cyrtodactylus peguensis</i> group (Reptilia: Squamata: Gekkonidae) with descriptions of two new species from Myanmar. <i>PeerJ</i> , 2018, 6, e5575.	0.9	16
79	Gastrointestinal Helminths from Eight Species of <i>Cyrtodactylus</i> Lizards (Squamata: Gekkonidae) from Peninsular Malaysia and Two Species from Vietnam. <i>Pacific Science</i> , 2017, 71, 91-98.	0.2	3
80	A new species of Mud Snake (Serpentes, Homalopsidae, <i>Gyiophis</i> Murphy & Voris, 2014) from Myanmar with a first molecular phylogenetic assessment of the genus. <i>Zootaxa</i> , 2017, 4238, 571.	0.2	6
81	Nematoda of Eleven Species of Ranid Frogs (Anura: Ranidae) from Southeast Asia. <i>Pacific Science</i> , 2017, 71, 229-235.	0.2	6
82	Nematodes of Five Species of Dicroglossid Frogs (Anura: Dicroglossidae) from Southeast Asia. <i>Pacific Science</i> , 2017, 71, 535-540.	0.2	3
83	The Eurasian hot nightlife: Environmental forces associated with nocturnality in lizards. <i>Global Ecology and Biogeography</i> , 2017, 26, 1316-1325.	2.7	22
84	Nematodes of Five Species of Bufonids (Anura: Bufonidae) from Peninsular Malaysia. <i>Pacific Science</i> , 2017, 71, 367-375.	0.2	9
85	Species delimitation with gene flow: A methodological comparison and population genomics approach to elucidate cryptic species boundaries in Malaysian Torrent Frogs. <i>Molecular Ecology</i> , 2017, 26, 5435-5450.	2.0	56
86	A re-evaluation of the phylogenetic relationships of the <i>Cyrtodactylus condorensis</i> group (Squamata: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 7 <i>Cyrtodactylus</i> . <i>Zootaxa</i> , 2017, 4300, .	0.2	33
87	Three new karst-dwelling <i>Cnemaspis</i> Strauch, 1887 (Squamata; Gekkonidae) from Peninsular Thailand and the phylogenetic placement of <i>C. Apunctatonuchalis</i> and <i>C. Avandeventeri</i> . <i>PeerJ</i> , 2017, 5, e2884.	0.9	15
88	Polyphyly of Asian Tree Toads, Genus <i>Pedostibes</i> Günther, 1876 (Anura: Bufonidae), and the Description of a New Genus from Southeast Asia. <i>PLoS ONE</i> , 2016, 11, e0145903.	1.1	11
89	Species composition, diversity and relative abundance of amphibians in forests and non-forest habitats on Langkawi Island, Peninsular Malaysia. , 2016, , .		1
90	Herpetofauna diversity at Gunung Raya, Pulau Langkawi, Kedah, Peninsular Malaysia. <i>AIP Conference Proceedings</i> , 2016, , .	0.3	2

#	ARTICLE	IF	CITATIONS
91	A new species of karst forest Bent-toed Gecko (genus <i>Tytthoscincus</i>) from Peninsular Malaysia's endemic karst forest herpetofauna and the need for its conservation. <i>Zootaxa</i> , 2016, 4061, 1.	0.2	42
92	The transfer of two clades of Malaysian <i>Sphenomorphus</i> Fitzinger (Squamata: Scincidae) into the genus <i>Tytthoscincus</i> Linkem, Diesmos, & Brown and the description of a new Malaysian swamp-dwelling species. <i>Zootaxa</i> , 2016, 4092, 231-42.	0.2	6
93	Taxonomy, phylogeny, and distribution of <i>Bronchocela rayaensis</i> (Squamata: Agamidae) on the Thai-Malay Peninsula. <i>Zootaxa</i> , 2016, 4092, 414-20.	0.2	4
94	The phylogenetic relationships of a new Stream Toad of the genus <i>Ansonia</i> Stoliczka, 1870 (Anura: Anura) from Peninsular Malaysia. <i>Zootaxa</i> , 2016, 4107, 517-40.	0.2	13
95	Phylogenetic relationships of geckos of the <i>Hemiphyllodactylus harterti</i> group, a new species from Penang Island, Peninsular Malaysia, and a likely case of true cryptic speciation. <i>Zootaxa</i> , 2016, 4107, 367-80.	0.2	19
96	Redescription <i>Cyrtodactylus lateralis</i> (Werner) (Squamata: Gekkonidae) and Phylogeny of the Prehensile-tailed <i>Cyrtodactylus</i> . <i>Zootaxa</i> , 2016, 4107, 517-40.	0.2	19
97	Two new Bent-toed Geckos of the <i>Cyrtodactylus pulchellus</i> complex from Peninsular Malaysia and multiple instances of convergent adaptation to limestone forest ecosystems. <i>Zootaxa</i> , 2016, 4105, 401-29.	0.2	26
98	The first riparian skink (Genus: <i>Sphenomorphus</i> Strauch, 1887) from Peninsular Malaysia and its relationship to other Indochinese and Sundaic species. <i>Zootaxa</i> , 2016, 4173, 29.	0.2	15
99	Archipelago-wide survey of Philippine forest dragons (Agamidae: <i>Gonocephalus</i>): multilocus phylogeny uncovers unprecedented levels of genetic diversity in a biodiversity hotspot. <i>Biological Journal of the Linnean Society</i> , 2016, , .	0.7	5
100	On the taxonomy and phylogeny of the skinks <i>Lipinia sekayuensis</i> Grismer, Ismail, Awang, Rizal, & Ahmad and <i>Lipinia surda</i> Boulenger from Peninsular Malaysia. <i>Zootaxa</i> , 2016, 4147, 59.	0.2	11
101	Out of Borneo, again and again: biogeography of the Stream Toad genus <i>Ansonia</i> Stoliczka (Anura: Bufonidae) and the discovery of the first limestone cave-dwelling species. <i>Biological Journal of the Linnean Society</i> , 2016, , .	0.7	19
102	Dragons in the mist: three new species of <i>Pseudocalotes</i> Fitzinger (Squamata: Agamidae) from the sky island archipelago of Peninsular Malaysia. <i>Zootaxa</i> , 2016, 4136, 461-90.	0.2	19
103	Helminths of Five Species of <i>Gonocephalus</i> Lizards (Squamata: Agamidae) from Peninsular Malaysia. <i>Pacific Science</i> , 2016, 70, 373-380.	0.2	7
104	A Species-Level Phylogeny of Extant Snakes with Description of a New Colubrid Subfamily and Genus. <i>PLoS ONE</i> , 2016, 11, e0161070.	1.1	206
105	A new lowland forest Bent-toed Gecko (Squamata: Gekkonidae: <i>Cyrtodactylus</i>) from Ranong Province, peninsular Thailand. <i>Zootaxa</i> , 2015, 3911, 106.	0.2	16
106	Repeated evolution of sympatric, palaeoendemic species in closely related, co-distributed lineages of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae) across a sky-island archipelago in Peninsular Malaysia. <i>Zoological Journal of the Linnean Society</i> , 2015, 174, 859-876.	1.0	22
107	The Rediscovery of <i>Sphenomorphus malayanus</i> Doria, 1888 (Squamata: Scincidae) from the Titiwangsa Mountain Range of Peninsular Malaysia and its re-description as <i>S. senja</i> sp. nov.. <i>Zootaxa</i> , 2015, 3931, 63-70.	0.2	25
108	A new insular species of Rock Gecko (<i>Cnemaspis</i> Boulenger) from Pulau Langkawi, Kedah, Peninsular Malaysia. <i>Zootaxa</i> , 2015, 3985, 203-18.	0.2	17

#	ARTICLE	IF	CITATIONS
109	The systematics and independent evolution of cave ecomorphology in distantly related clades of Bent-toed Geckos (Genus <i>Cyrtodactylus</i> ; Gray, 1827) from the Mekong Delta and islands in the Gulf of Thailand. <i>Zootaxa</i> , 2015, 3980, 106.	0.2	37
110	<i>Cyrtodactylus rosichonariefi</i> sp. nov. (Squamata: Gekkonidae), a new swamp-dwelling bent-toed gecko from Bunguran Island (Great Natuna), Indonesia. <i>Zootaxa</i> , 2015, 3964, 114-24.	0.2	11
111	The fourth Bent-toed Gecko of the genus <i>Cyrtodactylus</i> (Squamata: Gekkonidae). <i>Zootaxa</i> , 2015, 3980, 106.	0.2	17
112	Gastrointestinal helminths of nine species of <i>Cnemaspis</i> (Squamata: Gekkonidae) from Peninsular Malaysia, one species from Cambodia and Thailand and two species from Vietnam. <i>Journal of Natural History</i> , 2015, 49, 2683-2691.	0.2	5
113	An integrative taxonomic review of the agamid genus <i>Bronchocela</i> (Kuhl, 1820) from Peninsular Malaysia with descriptions of new montane and insular endemics. <i>Zootaxa</i> , 2015, 3948, 1.	0.2	28
114	New <i>Ichthyophis</i> species from Indochina (Gymnophiona, Ichthyophiidae): 1. The unstriped forms with descriptions of three new species and the redescription of <i>I. acuminatus</i> Taylor, 1960, <i>I. youngorum</i> Taylor, 1960 and <i>I. laosensis</i> Taylor, 1969. <i>Organisms Diversity and Evolution</i> , 2015, 15, 143-174.	0.7	9
115	A new species of karst forest-adapted Bent-toed Gecko (genus <i>Cyrtodactylus</i> Gray, 1827) belonging to the <i>C. sworderi</i> complex from a threatened karst forest in Perak, Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3755, 434-46.	0.2	28
116	A new species of insular Rock Gecko (Genus <i>Cnemaspis</i> Strauch, 1887) from the Bidong Archipelago, Terengganu, Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3755, 447-56.	0.2	15
117	The phylogenetic relationships of three new species of the <i>Cyrtodactylus pulchellus</i> complex (Squamata: Gekkonidae) from poorly explored regions in northeastern Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3786, 359.	0.2	37
118	A diminutive new species of cave-dwelling Wolf Snake (Colubridae: <i>Lycodon</i> Boie, 1826) from Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3815, 51-67.	0.2	16
119	Systematics and natural history of Southeast Asian Rock Geckos (genus <i>Cnemaspis</i> Strauch, 1887) with descriptions of eight new species from Malaysia, Thailand, and Indonesia. <i>Zootaxa</i> , 2014, 3880, 1-147.	0.2	80
120	The phylogenetic position and taxonomic status of the Rainbow Tree Snake <i>Gonyophis margaritatus</i> (Peters, 1871) (Squamata: Colubridae). <i>Zootaxa</i> , 2014, 3881, 532-48.	0.2	16
121	A new species of lowland skink (genus <i>Lipinia</i>) from Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3821, 457.	0.2	13
122	Multiple origins of parthenogenesis, and a revised species phylogeny for the Southeast Asian butterfly lizards, <i>Leiolepis</i> . <i>Biological Journal of the Linnean Society</i> , 2014, 113, 1080-1093.	0.7	30
123	A New Species of Frog (Amphibia: Anura: Ranidae) of the <i>Hylarana signata</i> Complex from Peninsular Malaysia. <i>Herpetologica</i> , 2014, 70, 228-240.	0.2	9
124	Description of a New Species of <i>Goniurosaurus</i> (Squamata: Eublepharidae) from the Guangdong Province, China, Based on Molecular and Morphological Data. <i>Herpetologica</i> , 2014, 70, 309-322.	0.2	13
125	A new species of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae). <i>Zootaxa</i> , 2014, 3755, 539.	0.2	16
126	A new species of <i>Hemiphyllodactylus</i> Bleeker, 1860 (Squamata: Gekkonidae). <i>Zootaxa</i> , 2014, 3755, 539.	0.2	21

#	ARTICLE	IF	CITATIONS
127	<i>Cyrtodactylus</i> in the city: A new Bent-toed Gecko (Genus <i>Cyrtodactylus</i>) is the only endemic species of vertebrate from Batu Caves, Selangor, Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3774, 381.	0.2	28
128	A new species of <i>Hemiphyllodactylus</i> ; Bleeker, 1860 (Squamata: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 3821, 485.	0.2	18
129	Reappraisal of the Javanese Bullfrog complex, <i>Kaloula</i> ; (MÄ;4ller, 1836) (Amphibia: Anura: Microhylidae), reveals a new species from Peninsular Malaysia. <i>Zootaxa</i> , 2014, 3900, 569.	0.2	8
130	Integrative taxonomy uncovers high levels of cryptic species diversity in <i>Hemiphyllodactylus</i> ...Bleeker, 1860 (Squamata: Gekkonidae) and the description of a new species from Peninsular Malaysia. <i>Zoological Journal of the Linnean Society</i> , 2013, 169, 849-880.	1.0	54
131	A new species of karst-adapted <i>Cnemaspis</i> ; Strauch, 1887 (Squamata: Gekkonidae) from a threatened karst region in Pahang, Peninsular Malaysia. <i>Zootaxa</i> , 2013, 3746, 463.	0.2	14
132	Cryptic speciation within <i>Asthenodipsas vertebralis</i> ; (Boulenger, 1900) (Squamata: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 3664, 505-24.	0.2	24
133	Phylogenetic relationships and description of a new upland species of Bent-toed Gecko (<i>Cyrtodactylus</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 239-52.	0.2	23
134	<i>Cyrtodactylus sanook</i> (Squamata: Gekkonidae), a new cave-dwelling gecko from Chumphon Province, southern Thailand. <i>Zootaxa</i> , 2013, 3635, 275-285.	0.2	26
135	Phylogeny and cryptic diversification in Southeast Asian flying geckos. <i>Molecular Phylogenetics and Evolution</i> , 2012, 65, 351-361.	1.2	33
136	Phylogeography, geographic variation, and taxonomy of the Bent-toed Gecko <i>Cyrtodactylus quadrivirgatus</i> Taylor, 1962 from Peninsular Malaysia with the description of a new swamp dwelling species. <i>Zootaxa</i> , 2012, 3406, 39.	0.2	45
137	A new endemic species of <i>Cyrtodactylus</i> Gray (Squamata: Gekkonidae) from Tho Chu Island, southwestern Vietnam. <i>Zootaxa</i> , 2012, 3228, .	0.2	24
138	A phylogeny and taxonomy of the Thai-Malay Peninsula Bent-toed Geckos of the <i>Cyrtodactylus pulchellus</i> complex (Squamata: Gekkonidae): combined morphological and molecular analyses with descriptions of seven new species. <i>Zootaxa</i> , 2012, 3520, 1.	0.2	85
139	A new insular, endemic frog of the genus <i>Kalophrynus</i> Tschudi, 1838 (Anura: Microhylidae) from Tioman Island, Pahang, Peninsular Malaysia Department of Biology, La Sierra University, 4500 Riverwalk Parkway, Riverside, California, 92515-8247 USA.. <i>Zootaxa</i> , 2011, 3123, .	0.2	6
140	Peninsular Malaysia's first limbless lizard: a new species of skink of the genus <i>Larutia</i> (BÄ;hme) from Pulau Pinang with a phylogeny of the genus. <i>Zootaxa</i> , 2011, 2799, .	0.2	16
141	First Record of <i>Batrachochytrium dendrobatidis</i> Infecting Four Frog Families from Peninsular Malaysia. <i>EcoHealth</i> , 2011, 8, 121-128.	0.9	20
142	Another new, diminutive Rock Gecko (<i>Cnemaspis</i> Strauch) from Peninsular Malaysia and a discussion of resource partitioning in sympatric species pairs. <i>Zootaxa</i> , 2010, 2569, .	0.2	12
143	A new species of <i>Cyrtodactylus</i> Gray, 1827 (Squamata: Gekkonidae) in Phu Quoc National Park, Kien Giang Biosphere Reserve, Southwestern Vietnam. <i>Zootaxa</i> , 2010, 2604, 37.	0.2	26
144	A new spiny, prehensile-tailed species of <i>Cyrtodactylus</i> (Squamata: Gekkonidae) from Peninsular Malaysia with a preliminary hypothesis of relationships based on morphology. <i>Zootaxa</i> , 2010, 2625, .	0.2	25

#	ARTICLE	IF	CITATIONS
145	Herpetofauna of gunung panti forest reserve, johor, peninsular malaysia. Tropical Life Sciences Research, 2010, 21, 71-82.	0.5	0
146	Two new diminutive species of Cnemaspis Strauch 1887 (Squamata: Gekkonidae) from Peninsular Malaysia. Zootaxa, 2009, 2019, 40-56.	0.2	27
147	A new species of Gastrophryonoides (Anura: Microhylidae): an addition to a previously monotypic genus and a new genus for Peninsular Malaysia. Zootaxa, 2009, 2124, 63-68.	0.2	10
148	Two New Species of Torrent-dwelling Toads Ansonia Stoliczka, 1870 (Anura: Bufonidae) from Peninsular Malaysia. Herpetologica, 2008, 64, 321-340.	0.2	24
149	A new species of insular pitviper of the genus Cryptelytrops (Squamata: Viperidae) from southern Vietnam. Zootaxa, 2008, 1715, 57.	0.2	12
150	A new species of karst dwelling gecko (genus Cnemaspis Strauch 1887) from the border region of Thailand and Peninsular Malaysia. Zootaxa, 2008, 1875, 51.	0.2	31
151	A new insular species of Cyrtodactylus (Squamata: Gekkonidae) from the Langkawi Archipelago, Kedah, Peninsular Malaysia. Zootaxa, 2008, 1924, 53-68.	0.2	24
152	A new species of insular skink (Genus Sphenomorphus Fitzinger 1843) from the Langkawi Archipelago, Kedah, West Malaysia with the first report of the herpetofauna of Pulau Singa Besar and an updated checklist of the herpetofauna of Pulau Langkawi. Zootaxa, 2008, 1691, 53.	0.2	29
153	A new species of Cnemaspis Strauch 1887 (Squamata: Gekkonidae) from Selangor, Peninsular Malaysia. Zootaxa, 2008, 1877, 49.	0.2	20
154	A revised and updated checklist of the lizards of Peninsular Malaysia. Zootaxa, 2008, 1860, 28.	0.2	13
155	A NEW SPECIES OF SMALL MONTANE FOREST FLOOR SKINK (GENUS SPHENOMORPHUS FITZINGER 1843) FROM SOUTHERN PENINSULAR MALAYSIA. Herpetologica, 2007, 63, 544-551.	0.2	14
156	FOUR NEW SPECIES OF THE GEKKONID GENUS CNEMASPIS STRAUCH 1887 (REPTILIA: SQUAMATA) FROM SOUTHERN VIETNAM. Herpetologica, 2007, 63, 482-500.	0.2	42
157	A new species of Ansonia Stoliczka 1872 (Anura: Bufonidae) from Central Peninsular Malaysia and a revised taxonomy for Ansonia from the Malay Peninsula. Zootaxa, 2006, 1327, 1.	0.2	16
158	TWO NEW SPECIES OF CNEMASPIS STRAUCH, 1887 (SQUAMATA: GEKKONIDAE) FROM THE SERIBUAT ARCHIPELAGO, PAHANG AND JOHOR STATES, WEST MALAYSIA. Herpetologica, 2003, 59, 544-552.	0.2	29
159	Evolutionary biogeography on Mexico's Baja California peninsula: A synthesis of molecules and historical geology. Proceedings of the National Academy of Sciences of the United States of America, 2000, 97, 14017-14018.	3.3	38
160	Phylogenetic partitioning of the third-largest vertebrate genus in the world, Cyrtodactylus Gray, 1827 (Reptilia; Squamata; Gekkonidae) and its relevance to taxonomy and conservation. Vertebrate Zoology, 0, 71, 101-154.	2.0	25
161	Phylogenetic partitioning of the third-largest vertebrate genus in the world, Cyrtodactylus Gray, 1827 (Reptilia; Squamata; Gekkonidae) and its relevance to taxonomy and conservation. Vertebrate Zoology, 0, 71, 101-154.	2.0	16
162	Phylogeny and evolution of habitat preference in Goniurosaurus (Squamata: Eublepharidae) and their correlation with karst and granite-stream-adapted ecomorphologies in species groups from Vietnam. Vertebrate Zoology, 0, 71, 335-352.	2.0	6

