

Haruhiko Koseki

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1640843/publications.pdf>

Version: 2024-02-01

350
papers

41,273
citations

2963

93
h-index

2940

189
g-index

367
all docs

367
docs citations

367
times ranked

46094
citing authors

#	ARTICLE	IF	CITATIONS
1	Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. <i>Nature</i> , 2013, 504, 446-450.	13.7	3,901
2	CD1d-Restricted and TCR-Mediated Activation of $V\alpha 14$ NKT Cells by Glycosylceramides. <i>Science</i> , 1997, 278, 1626-1629.	6.0	2,274
3	Control of Developmental Regulators by Polycomb in Human Embryonic Stem Cells. <i>Cell</i> , 2006, 125, 301-313.	13.5	2,059
4	A promoter-level mammalian expression atlas. <i>Nature</i> , 2014, 507, 462-470.	13.7	1,838
5	Requirement for $V\alpha 14$ NKT Cells in IL-12-Mediated Rejection of Tumors. <i>Science</i> , 1997, 278, 1623-1626.	6.0	1,190
6	The SRA protein Np95 mediates epigenetic inheritance by recruiting Dnmt1 to methylated DNA. <i>Nature</i> , 2007, 450, 908-912.	13.7	1,096
7	Genomewide Analysis of PRC1 and PRC2 Occupancy Identifies Two Classes of Bivalent Domains. <i>PLoS Genetics</i> , 2008, 4, e1000242.	1.5	878
8	Polycomb Group Proteins Ring1A/B Link Ubiquitylation of Histone H2A to Heritable Gene Silencing and X Inactivation. <i>Developmental Cell</i> , 2004, 7, 663-676.	3.1	829
9	Variant PRC1 Complex-Dependent H2A Ubiquitylation Drives PRC2 Recruitment and Polycomb Domain Formation. <i>Cell</i> , 2014, 157, 1445-1459.	13.5	613
10	Ring1-mediated ubiquitination of H2A restrains poised RNA polymerase II at bivalent genes in mouse ES cells. <i>Nature Cell Biology</i> , 2007, 9, 1428-1435.	4.6	584
11	Transcribed enhancers lead waves of coordinated transcription in transitioning mammalian cells. <i>Science</i> , 2015, 347, 1010-1014.	6.0	517
12	Enhanced Self-Renewal of Hematopoietic Stem Cells Mediated by the Polycomb Gene Product Bmi-1. <i>Immunity</i> , 2004, 21, 843-851.	6.6	486
13	Natural killer-like nonspecific tumor cell lysis mediated by specific ligand-activated $V\alpha 14$ NKT cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 5690-5693.	3.3	443
14	Polycomb Limits the Neurogenic Competence of Neural Precursor Cells to Promote Astrogenic Fate Transition. <i>Neuron</i> , 2009, 63, 600-613.	3.8	420
15	KDM2B links the Polycomb Repressive Complex 1 (PRC1) to recognition of CpG islands. <i>ELife</i> , 2012, 1, e00205.	2.8	414
16	Pax genes and organogenesis. <i>BioEssays</i> , 1997, 19, 755-765.	1.2	360
17	Recruitment of PRC1 function at the initiation of X inactivation independent of PRC2 and silencing. <i>EMBO Journal</i> , 2006, 25, 3110-3122.	3.5	353
18	Disruption of the Bcl6 Gene Results in an Impaired Germinal Center Formation. <i>Journal of Experimental Medicine</i> , 1997, 186, 439-448.	4.2	336

#	ARTICLE	IF	CITATIONS
19	Polycomb repressive complex PRC1 spatially constrains the mouse embryonic stem cell genome. <i>Nature Genetics</i> , 2015, 47, 1179-1186.	9.4	330
20	Generation of Rejuvenated Antigen-Specific T Cells by Reprogramming to Pluripotency and Redifferentiation. <i>Cell Stem Cell</i> , 2013, 12, 114-126.	5.2	327
21	Mouse model of Prinzmetal angina by disruption of the inward rectifier Kir6.1. <i>Nature Medicine</i> , 2002, 8, 466-472.	15.2	306
22	Uhrf1-dependent H3K23 ubiquitylation couples maintenance DNA methylation and replication. <i>Nature</i> , 2013, 502, 249-253.	13.7	305
23	Mesp2: a novel mouse gene expressed in the presegmented mesoderm and essential for segmentation initiation.. <i>Genes and Development</i> , 1997, 11, 1827-1839.	2.7	300
24	The adaptor protein CARD9 is essential for the activation of myeloid cells through ITAM-associated and Toll-like receptors. <i>Nature Immunology</i> , 2007, 8, 619-629.	7.0	300
25	PRC1 and Suv39h specify parental asymmetry at constitutive heterochromatin in early mouse embryos. <i>Nature Genetics</i> , 2008, 40, 411-420.	9.4	291
26	Targeting Polycomb to Pericentric Heterochromatin in Embryonic Stem Cells Reveals a Role for H2AK119u1 in PRC2 Recruitment. <i>Cell Reports</i> , 2014, 7, 1456-1470.	2.9	283
27	UHRF1 targets DNMT1 for DNA methylation through cooperative binding of hemi-methylated DNA and methylated H3K9. <i>Nature Communications</i> , 2013, 4, 1563.	5.8	275
28	Selective ablation of basophils in mice reveals their nonredundant role in acquired immunity against ticks. <i>Journal of Clinical Investigation</i> , 2010, 120, 2867-2875.	3.9	272
29	SAM Domain Polymerization Links Subnuclear Clustering of PRC1 to Gene Silencing. <i>Developmental Cell</i> , 2013, 26, 565-577.	3.1	271
30	Deep transcriptome profiling of mammalian stem cells supports a regulatory role for retrotransposons in pluripotency maintenance. <i>Nature Genetics</i> , 2014, 46, 558-566.	9.4	271
31	Regeneration of Human Tumor Antigen-Specific T Cells from iPSCs Derived from Mature CD8 + T Cells. <i>Cell Stem Cell</i> , 2013, 12, 31-36.	5.2	270
32	Characterization and Developmental Expression of Pax9, a Paired-Box-Containing Gene Related to Pax1. <i>Developmental Biology</i> , 1995, 170, 701-716.	0.9	266
33	Polycomb group proteins Ring1A/B are functionally linked to the core transcriptional regulatory circuitry to maintain ES cell identity. <i>Development (Cambridge)</i> , 2008, 135, 1513-1524.	1.2	265
34	The Polycomb Protein Ezh2 Regulates Differentiation and Plasticity of CD4+ T Helper Type 1 and Type 2 Cells. <i>Immunity</i> , 2013, 39, 819-832.	6.6	260
35	The Zinc Transporter SLC39A13/ZIP13 Is Required for Connective Tissue Development; Its Involvement in BMP/TGF- β Signaling Pathways. <i>PLoS ONE</i> , 2008, 3, e3642.	1.1	240
36	Histone H2A Mono-Ubiquitination Is a Crucial Step to Mediate PRC1-Dependent Repression of Developmental Genes to Maintain ES Cell Identity. <i>PLoS Genetics</i> , 2012, 8, e1002774.	1.5	233

#	ARTICLE	IF	CITATIONS
37	PCGF3/5â€“PRC1 initiates Polycomb recruitment in X chromosome inactivation. <i>Science</i> , 2017, 356, 1081-1084.	6.0	220
38	Dynamic Reprogramming of DNA Methylation at an Epigenetically Sensitive Allele in Mice. <i>PLoS Genetics</i> , 2006, 2, e49.	1.5	218
39	Prolonged Mek1/2 suppression impairs the developmental potential of embryonic stem cells. <i>Nature</i> , 2017, 548, 219-223.	13.7	211
40	Synergy between Variant PRC1 Complexes Defines Polycomb-Mediated Gene Repression. <i>Molecular Cell</i> , 2019, 74, 1020-1036.e8.	4.5	200
41	Dependency on the polycomb gene Ezh2 distinguishes fetal from adult hematopoietic stem cells. <i>Blood</i> , 2011, 118, 6553-6561.	0.6	198
42	Mesp2 initiates somite segmentation through the Notch signalling pathway. <i>Nature Genetics</i> , 2000, 25, 390-396.	9.4	197
43	FANTOM5 CAGE profiles of human and mouse samples. <i>Scientific Data</i> , 2017, 4, 170112.	2.4	195
44	Angiotensin-related growth factor antagonizes obesity and insulin resistance. <i>Nature Medicine</i> , 2005, 11, 400-408.	15.2	194
45	Repression of the Transcription Factor Bach2 Contributes to Predisposition of IgG1 Memory B Cells toward Plasma Cell Differentiation. <i>Immunity</i> , 2013, 39, 136-147.	6.6	187
46	Targeted disruption of Traf5 gene causes defects in CD40- and CD27-mediated lymphocyte activation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 9803-9808.	3.3	183
47	Development and Function of Invariant Natural Killer T Cells Producing TH2- and TH17-Cytokines. <i>PLoS Biology</i> , 2012, 10, e1001255.	2.6	180
48	Bypass of senescence by the polycomb group protein CBX8 through direct binding to the INK4A-ARF locus. <i>EMBO Journal</i> , 2007, 26, 1637-1648.	3.5	175
49	SF3B1 haploinsufficiency leads to formation of ring sideroblasts in myelodysplastic syndromes. <i>Blood</i> , 2012, 120, 3173-3186.	0.6	173
50	The Hbo1-Brd1/Brpf2 complex is responsible for global acetylation of H3K14 and required for fetal liver erythropoiesis. <i>Blood</i> , 2011, 118, 2443-2453.	0.6	168
51	Involvement of decidual Valpha 14 NKT cells in abortion. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 740-744.	3.3	167
52	H2A.Z landscapes and dual modifications in pluripotent and multipotent stem cells underlie complex genome regulatory functions. <i>Genome Biology</i> , 2012, 13, R85.	13.9	166
53	ESCs Require PRC2 to Direct the Successful Reprogramming of Differentiated Cells toward Pluripotency. <i>Cell Stem Cell</i> , 2010, 6, 547-556.	5.2	162
54	Concurrent loss of <i>Ezh2</i> and <i>Tet2</i> cooperates in the pathogenesis of myelodysplastic disorders. <i>Journal of Experimental Medicine</i> , 2013, 210, 2627-2639.	4.2	162

#	ARTICLE	IF	CITATIONS
55	Ezh2 augments leukemogenicity by reinforcing differentiation blockage in acute myeloid leukemia. <i>Blood</i> , 2012, 120, 1107-1117.	0.6	161
56	Novel regulation of MHC class II function in B cells. <i>EMBO Journal</i> , 2007, 26, 846-854.	3.5	158
57	Senescence Marker Protein-30 Knockout Mouse Liver Is Highly Susceptible to Tumor Necrosis Factor- α - and Fas-Mediated Apoptosis. <i>American Journal of Pathology</i> , 2002, 161, 1273-1281.	1.9	156
58	CBX8, a Polycomb Group Protein, Is Essential for MLL-AF9-Induced Leukemogenesis. <i>Cancer Cell</i> , 2011, 20, 563-575.	7.7	156
59	Inhibition of T Helper Cell Type 2 Cell Differentiation and Immunoglobulin E Response by Ligand-Activated V α 14 Natural Killer T Cells. <i>Journal of Experimental Medicine</i> , 1999, 190, 783-792.	4.2	153
60	Dilated Cardiomyopathy Caused by Aberrant Endoplasmic Reticulum Quality Control in Mutant KDEL Receptor Transgenic Mice. <i>Molecular and Cellular Biology</i> , 2004, 24, 8007-8017.	1.1	152
61	The epigenetic regulator Uhrf1 facilitates the proliferation and maturation of colonic regulatory T cells. <i>Nature Immunology</i> , 2014, 15, 571-579.	7.0	147
62	The Zinc Transporter SLC39A14/ZIP14 Controls G-Protein Coupled Receptor-Mediated Signaling Required for Systemic Growth. <i>PLoS ONE</i> , 2011, 6, e18059.	1.1	147
63	Cooperative B7-1/2 (CD80/CD86) and B7-DC Costimulation of CD4+ T Cells Independent of the PD-1 Receptor. <i>Journal of Experimental Medicine</i> , 2003, 198, 31-38.	4.2	144
64	Ezh2 loss promotes development of myelodysplastic syndrome but attenuates its predisposition to leukaemic transformation. <i>Nature Communications</i> , 2014, 5, 4177.	5.8	143
65	Generation of Cloned Mice by Direct Nuclear Transfer from Natural Killer T Cells. <i>Current Biology</i> , 2005, 15, 1114-1118.	1.8	142
66	In vivo costimulatory role of B7-DC in tuning T helper cell 1 and cytotoxic T lymphocyte responses. <i>Journal of Experimental Medicine</i> , 2005, 201, 1531-1541.	4.2	140
67	PRC2.1 and PRC2.2 Synergize to Coordinate H3K27 Trimethylation. <i>Molecular Cell</i> , 2019, 76, 437-452.e6.	4.5	137
68	The KDEL receptor mediates a retrieval mechanism that contributes to quality control at the endoplasmic reticulum. <i>EMBO Journal</i> , 2001, 20, 3082-3091.	3.5	135
69	Mammalian Polyhomeotic Homologues Phc2 and Phc1 Act in Synergy To Mediate Polycomb Repression of Hox Genes. <i>Molecular and Cellular Biology</i> , 2005, 25, 6694-6706.	1.1	133
70	Extrathymic development of V alpha 14-positive T cells.. <i>Journal of Experimental Medicine</i> , 1993, 177, 1399-1408.	4.2	132
71	Roles of HIPK1 and HIPK2 in AML1- and p300-dependent transcription, hematopoiesis and blood vessel formation. <i>EMBO Journal</i> , 2006, 25, 3955-3965.	3.5	124
72	Type II membrane protein CD69 regulates the formation of resting T-helper memory. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 7409-7414.	3.3	121

#	ARTICLE	IF	CITATIONS
73	A Family of Vertebrate-Specific Polycombs Encoded by the LCOR/LCORL Genes Balance PRC2 Subtype Activities. <i>Molecular Cell</i> , 2018, 70, 408-421.e8.	4.5	121
74	A Phosphorylated Form of Mel-18 Targets the Ring1B Histone H2A Ubiquitin Ligase to Chromatin. <i>Molecular Cell</i> , 2007, 28, 107-120.	4.5	118
75	Opposing roles of polycomb repressive complexes in hematopoietic stem and progenitor cells. <i>Blood</i> , 2010, 116, 731-739.	0.6	117
76	The Role of mel-18, a Mammalian Polycomb Group Gene, during IL-7-Dependent Proliferation of Lymphocyte Precursors. <i>Immunity</i> , 1997, 7, 135-146.	6.6	112
77	A novel pathogenesis of megacolon in Ncx/Hox11L.1 deficient mice.. <i>Journal of Clinical Investigation</i> , 1997, 100, 795-801.	3.9	109
78	Homogenous junctional sequence of the V14+ T-cell antigen receptor alpha chain expanded in unprimed mice.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 5248-5252.	3.3	108
79	Regulation of Th2 Cell Differentiation by mel-18, a Mammalian Polycomb Group Gene. <i>Immunity</i> , 2001, 15, 275-287.	6.6	107
80	S phase-dependent interaction with DNMT1 dictates the role of UHRF1 but not UHRF2 in DNA methylation maintenance. <i>Cell Research</i> , 2011, 21, 1723-1739.	5.7	107
81	Zinc transporter SLC39A10/ZIP10 facilitates antiapoptotic signaling during early B-cell development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 11780-11785.	3.3	107
82	DNA polymerase β contributes to the generation of C/G mutations during somatic hypermutation of Ig genes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 13986-13991.	3.3	106
83	PRC1 Fine-tunes Gene Repression and Activation to Safeguard Skin Development and Stem Cell Specification. <i>Cell Stem Cell</i> , 2018, 22, 726-739.e7.	5.2	106
84	Pax-1, a regulator of sclerotome development is induced by notochord and floor plate signals in avian embryos. <i>Anatomy and Embryology</i> , 1995, 191, 297-310.	1.5	105
85	FGF9 monomer-dimer equilibrium regulates extracellular matrix affinity and tissue diffusion. <i>Nature Genetics</i> , 2009, 41, 289-298.	9.4	104
86	Nucleotide-sugar transporter SLC35D1 is critical to chondroitin sulfate synthesis in cartilage and skeletal development in mouse and human. <i>Nature Medicine</i> , 2007, 13, 1363-1367.	15.2	103
87	Zinc transporter SLC39A10/ZIP10 controls humoral immunity by modulating B-cell receptor signal strength. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 11786-11791.	3.3	103
88	Expression of Avian Pax1 and Pax9 Is Intrinsically Regulated in the Pharyngeal Endoderm, but Depends on Environmental Influences in the Paraxial Mesoderm. <i>Developmental Biology</i> , 1996, 178, 403-417.	0.9	102
89	Mammalian Polycomb-mediated repression of Hox genes requires the essential spliceosomal protein Sf3b1. <i>Genes and Development</i> , 2005, 19, 536-541.	2.7	102
90	Overlapping Roles for Homeodomain-Interacting Protein Kinases Hipk1 and Hipk2 in the Mediation of Cell Growth in Response to Morphogenetic and Genotoxic Signals. <i>Molecular and Cellular Biology</i> , 2006, 26, 2758-2771.	1.1	102

#	ARTICLE	IF	CITATIONS
91	The Generation of Mature, Single-Positive Thymocytes In Vivo Is Dysregulated by CD69 Blockade or Overexpression. <i>Journal of Immunology</i> , 2002, 168, 87-94.	0.4	101
92	Growth retardation and skin abnormalities of the Recql4-deficient mouse. <i>Human Molecular Genetics</i> , 2003, 12, 2293-2299.	1.4	101
93	Ezh2 is required for neural crest-derived cartilage and bone formation. <i>Development (Cambridge)</i> , 2014, 141, 867-877.	1.2	101
94	Dominant expression of a distinctive V14+ T-cell antigen receptor alpha chain in mice.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 7518-7522.	3.3	99
95	Model mice for tissue-specific deletion of the manganese superoxide dismutase (MnSOD) gene. <i>Biochemical and Biophysical Research Communications</i> , 2002, 296, 729-736.	1.0	99
96	PCGF6-PRC1 suppresses premature differentiation of mouse embryonic stem cells by regulating germ cell-related genes. <i>ELife</i> , 2017, 6, .	2.8	99
97	Polycomb Complex PRC1 Preserves Intestinal Stem Cell Identity by Sustaining Wnt/ β -Catenin Transcriptional Activity. <i>Cell Stem Cell</i> , 2016, 18, 91-103.	5.2	97
98	Membrane-bound human SCF/KL promotes in vivo human hematopoietic engraftment and myeloid differentiation. <i>Blood</i> , 2012, 119, 2768-2777.	0.6	96
99	Polycomb Potentiates Meis2 Activation in Midbrain by Mediating Interaction of the Promoter with a Tissue-Specific Enhancer. <i>Developmental Cell</i> , 2014, 28, 94-101.	3.1	96
100	Essential requirement of an invariant V alpha 14 T cell antigen receptor expression in the development of natural killer T cells.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 11025-11028.	3.3	95
101	Notochord-Dependent Expression of MFH1 and PAX1 Cooperates to Maintain the Proliferation of Sclerotome Cells during the Vertebral Column Development. <i>Developmental Biology</i> , 1999, 210, 15-29.	0.9	95
102	Role of UHRF1 in de novo DNA methylation in oocytes and maintenance methylation in preimplantation embryos. <i>PLoS Genetics</i> , 2017, 13, e1007042.	1.5	95
103	Enhancers are activated by p300/CBP activity-dependent PIC assembly, RNAPII recruitment, and pause release. <i>Molecular Cell</i> , 2021, 81, 2166-2182.e6.	4.5	94
104	Regeneration of CD8 α β T Cells from T-cell-Derived iPSC Imparts Potent Tumor Antigen-Specific Cytotoxicity. <i>Cancer Research</i> , 2016, 76, 6839-6850.	0.4	93
105	The Importance of Disinfection Therapy Using Povidone-Iodine Solution in Atopic Dermatitis. <i>Dermatology</i> , 2002, 204, 63-69.	0.9	91
106	Targeted Disruption of Dermopontin Causes Abnormal Collagen Fibrillogenesis. <i>Journal of Investigative Dermatology</i> , 2002, 119, 678-683.	0.3	91
107	The KDEL Receptor Modulates the Endoplasmic Reticulum Stress Response through Mitogen-activated Protein Kinase Signaling Cascades. <i>Journal of Biological Chemistry</i> , 2003, 278, 34525-34532.	1.6	91
108	A polycomb group protein, PHF1, is involved in the response to DNA double-strand breaks in human cell. <i>Nucleic Acids Research</i> , 2008, 36, 2939-2947.	6.5	89

#	ARTICLE	IF	CITATIONS
109	<i>Sall4</i> Is Essential for Stabilization, But Not for Pluripotency, of Embryonic Stem Cells by Repressing Aberrant Trophectoderm Gene Expression. <i>Stem Cells</i> , 2009, 27, 796-805.	1.4	89
110	Role of SOX17 in hematopoietic development from human embryonic stem cells. <i>Blood</i> , 2013, 121, 447-458.	0.6	87
111	Inactivation of the Polycomb Group Protein Ring1B Unveils an Antiproliferative Role in Hematopoietic Cell Expansion and Cooperation with Tumorigenesis Associated with <i>Ink4a</i> Deletion. <i>Molecular and Cellular Biology</i> , 2008, 28, 1018-1028.	1.1	86
112	The SET1 Complex Selects Actively Transcribed Target Genes via Multivalent Interaction with CpG Island Chromatin. <i>Cell Reports</i> , 2017, 20, 2313-2327.	2.9	86
113	Polycomblike 2 facilitates the recruitment of PRC2 Polycomb group complexes to the inactive X chromosome and to target loci in embryonic stem cells. <i>Development (Cambridge)</i> , 2011, 138, 1471-1482.	1.2	85
114	Involvement of the Polycomb-group gene <i>Ring1B</i> in the specification of the anterior-posterior axis in mice. <i>Development (Cambridge)</i> , 2002, 129, 4171-4183.	1.2	85
115	RYBP Represses Endogenous Retroviruses and Preimplantation- and Germ Line-Specific Genes in Mouse Embryonic Stem Cells. <i>Molecular and Cellular Biology</i> , 2012, 32, 1139-1149.	1.1	84
116	Functional analysis of AEBP2, a PRC2 Polycomb protein, reveals a Trithorax phenotype in embryonic development and in ES cells. <i>Development (Cambridge)</i> , 2016, 143, 2716-23.	1.2	84
117	Mouse Homologue of <i>coq7/clk-1</i> , Longevity Gene in <i>Caenorhabditis elegans</i> , Is Essential for Coenzyme Q Synthesis, Maintenance of Mitochondrial Integrity, and Neurogenesis. <i>Biochemical and Biophysical Research Communications</i> , 2001, 289, 463-471.	1.0	82
118	Cell-autonomous involvement of <i>Mab21l1</i> is essential for lens placode development. <i>Development (Cambridge)</i> , 2003, 130, 1759-1770.	1.2	82
119	Zinc Transporter SLC39A7/ZIP7 Promotes Intestinal Epithelial Self-Renewal by Resolving ER Stress. <i>PLoS Genetics</i> , 2016, 12, e1006349.	1.5	80
120	Polycomb proteins control proliferation and transformation independently of cell cycle checkpoints by regulating DNA replication. <i>Nature Communications</i> , 2014, 5, 3649.	5.8	79
121	<i>Bmi1</i> is a MYCN target gene that regulates tumorigenesis through repression of <i>KIF1B</i> and <i>TSLC1</i> in neuroblastoma. <i>Oncogene</i> , 2010, 29, 2681-2690.	2.6	77
122	Activation of Endogenous Retroviruses in <i>Dnmt1</i> ESCs Involves Disruption of SETDB1-Mediated Repression by NP95 Binding to Hemimethylated DNA. <i>Cell Stem Cell</i> , 2016, 19, 81-94.	5.2	77
123	H2AK119ub1 guides maternal inheritance and zygotic deposition of H3K27me3 in mouse embryos. <i>Nature Genetics</i> , 2021, 53, 539-550.	9.4	77
124	<i>Zic1</i> regulates the patterning of vertebral arches in cooperation with <i>Gli3</i> . <i>Mechanisms of Development</i> , 1999, 89, 141-150.	1.7	76
125	Forkhead transcription factor <i>Foxf2</i> (LUN)-deficient mice exhibit abnormal development of secondary palate. <i>Developmental Biology</i> , 2003, 259, 83-94.	0.9	75
126	<i>Noc2</i> is essential in normal regulation of exocytosis in endocrine and exocrine cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004, 101, 8313-8318.	3.3	75

#	ARTICLE	IF	CITATIONS
127	Positive selection of invariant V alpha 14+ T cells by non-major histocompatibility complex-encoded class I-like molecules expressed on bone marrow-derived cells.. Proceedings of the National Academy of Sciences of the United States of America, 1995, 92, 1200-1204.	3.3	74
128	Construction of an open-access database that integrates cross-reference information from the transcriptome and proteome of immune cells. Bioinformatics, 2007, 23, 2934-2941.	1.8	74
129	CCN3 Inhibits Neointimal Hyperplasia Through Modulation of Smooth Muscle Cell Growth and Migration. Arteriosclerosis, Thrombosis, and Vascular Biology, 2010, 30, 675-682.	1.1	74
130	Cell Cycle-Dependent Turnover of 5-Hydroxymethyl Cytosine in Mouse Embryonic Stem Cells. PLoS ONE, 2013, 8, e82961.	1.1	73
131	HP1 links histone methylation marks to meiotic synapsis in mice. Development (Cambridge), 2011, 138, 4207-4217.	1.2	71
132	CRTAM Confers Late-Stage Activation of CD8+ T Cells to Regulate Retention within Lymph Node. Journal of Immunology, 2009, 183, 4220-4228.	0.4	70
133	An epigenetic switch is crucial for spermatogonia to exit the undifferentiated state toward a Kit-positive identity. Development (Cambridge), 2013, 140, 3565-3576.	1.2	70
134	Ash1l Methylates Lys36 of Histone H3 Independently of Transcriptional Elongation to Counteract Polycomb Silencing. PLoS Genetics, 2013, 9, e1003897.	1.5	69
135	Mammalian Polycomb-Like Pcl2/Mtf2 Is a Novel Regulatory Component of PRC2 That Can Differentially Modulate Polycomb Activity both at the Hox Gene Cluster and at Cdkn2a Genes. Molecular and Cellular Biology, 2011, 31, 351-364.	1.1	68
136	Deficiency of the macrophage migration inhibitory factor gene has no significant effect on endotoxaemia. Immunology, 2000, 100, 84-90.	2.0	67
137	Requirement for Mab21l2 during development of murine retina and ventral body wall. Developmental Biology, 2004, 274, 295-307.	0.9	67
138	Ubiquitination-Independent Repression of PRC1 Targets during Neuronal Fate Restriction in the Developing Mouse Neocortex. Developmental Cell, 2018, 47, 758-772.e5.	3.1	67
139	A noncoding RNA regulates the neurogenin1 gene locus during mouse neocortical development. Proceedings of the National Academy of Sciences of the United States of America, 2012, 109, 16939-16944.	3.3	66
140	The polycomb component Ring1B regulates the timed termination of subcerebral projection neuron production during mouse neocortical development. Development (Cambridge), 2014, 141, 4343-4353.	1.2	66
141	Efficient Regeneration of Human V α 24+ Invariant Natural Killer T Cells and Their Anti-Tumor Activity In Vivo. Stem Cells, 2016, 34, 2852-2860.	1.4	65
142	Estrogen, Insulin, and Dietary Signals Cooperatively Regulate Longevity Signals to Enhance Resistance to Oxidative Stress in Mice. Journal of Biological Chemistry, 2005, 280, 16417-16426.	1.6	64
143	Polycomb Repressive Complexes Restrain the Expression of Lineage-Specific Regulators in Embryonic Stem Cells. Cell Cycle, 2006, 5, 1411-1414.	1.3	64
144	A Lysosomal Protein Negatively Regulates Surface T Cell Antigen Receptor Expression by Promoting CD3 ϵ -Chain Degradation. Immunity, 2008, 29, 33-43.	6.6	64

#	ARTICLE	IF	CITATIONS
145	Physiological Srsf2 P95H expression causes impaired hematopoietic stem cell functions and aberrant RNA splicing in mice. <i>Blood</i> , 2018, 131, 621-635.	0.6	64
146	Abnormal PcG protein expression in Hodgkin's lymphoma. Relation with E2F6 and NF κ B transcription factors. <i>Journal of Pathology</i> , 2004, 204, 528-537.	2.1	63
147	Identification of Epha4 enhancer required for segmental expression and the regulation by Mesp2. <i>Development (Cambridge)</i> , 2006, 133, 2517-2525.	1.2	63
148	Aberrant quality control in the endoplasmic reticulum impairs the biosynthesis of pulmonary surfactant in mice expressing mutant BiP. <i>Cell Death and Differentiation</i> , 2007, 14, 1475-1485.	5.0	63
149	The role of Bcl6 in mature cardiac myocytes. <i>Cardiovascular Research</i> , 1999, 42, 670-679.	1.8	62
150	Protective Role of Macrophages in Noninflammatory Lung Injury Caused by Selective Ablation of Alveolar Epithelial Type II Cells. <i>Journal of Immunology</i> , 2007, 178, 5001-5009.	0.4	60
151	Bmi1 cooperates with Dnmt1-associated protein 1 in gene silencing. <i>Biochemical and Biophysical Research Communications</i> , 2007, 353, 992-998.	1.0	60
152	CD69 α null mice protected from arthritis induced with anti α type II collagen antibodies. <i>International Immunology</i> , 2003, 15, 987-992.	1.8	59
153	Altered Quality Control in the Endoplasmic Reticulum Causes Cortical Dysplasia in Knock-In Mice Expressing a Mutant BiP. <i>Molecular and Cellular Biology</i> , 2008, 28, 293-301.	1.1	59
154	Oxygen Affinity of Hemoglobin Regulates O ₂ Consumption, Metabolism, and Physical Activity. <i>Journal of Biological Chemistry</i> , 2003, 278, 5035-5043.	1.6	58
155	WIP1, a Homeostatic Regulator of the DNA Damage Response, Is Targeted by HIPK2 for Phosphorylation and Degradation. <i>Molecular Cell</i> , 2013, 51, 374-385.	4.5	58
156	Mammalian Polycomb Scmh1 mediates exclusion of Polycomb complexes from the XY body in the pachytene spermatocytes. <i>Development (Cambridge)</i> , 2007, 134, 579-590.	1.2	57
157	A Novel Gene Essential for the Development of Single Positive Thymocytes. <i>Molecular and Cellular Biology</i> , 2009, 29, 5128-5135.	1.1	57
158	Maintenance of Undifferentiated State and Self-Renewal of Embryonic Neural Stem Cells by Polycomb Protein Ring1B. <i>Stem Cells</i> , 2009, 27, 1559-1570.	1.4	57
159	Production of Monoclonal Antibodies Against Mammalian Ring1B Proteins. <i>Hybridoma</i> , 2001, 20, 43-46.	0.9	56
160	Repressor of GATA regulates TH2-driven allergic airway inflammation and airway hyperresponsiveness. <i>Journal of Allergy and Clinical Immunology</i> , 2008, 122, 512-520.e11.	1.5	56
161	Smchd1 Targeting to the Inactive X Is Dependent on the Xist-HnrnpK-PRC1 Pathway. <i>Cell Reports</i> , 2018, 25, 1912-1923.e9.	2.9	56
162	Predominant use of a particular κ -chain in suppressor T cell hybridomas specific for keyhole limpet hemocyanin. <i>International Immunology</i> , 1989, 1, 557-564.	1.8	55

#	ARTICLE	IF	CITATIONS
163	Development of Valpha4+ NK T cells in the early stages of embryogenesis.. Proceedings of the National Academy of Sciences of the United States of America, 1996, 93, 6516-6520.	3.3	55
164	Abnormal blood vessel development in mice lacking presenilin-1. Mechanisms of Development, 2003, 120, 657-667.	1.7	55
165	Requirement of Zinc Transporter SLC39A7/ZIP7 for Dermal Development to Fine-Tune Endoplasmic Reticulum Function by Regulating Protein Disulfide Isomerase. Journal of Investigative Dermatology, 2017, 137, 1682-1691.	0.3	55
166	Tet2 and Tet3 in B cells are required to repress CD86 and prevent autoimmunity. Nature Immunology, 2020, 21, 950-961.	7.0	55
167	Murine induced pluripotent stem cells can be derived from and differentiate into natural killer T cells. Journal of Clinical Investigation, 2010, 120, 2610-2618.	3.9	55
168	Three developmental compartments involved in rib formation. International Journal of Developmental Biology, 2005, 49, 325-333.	0.3	52
169	Regulation of Th2 Cell Development by Polycomb Group Gene bmi-1 through the Stabilization of GATA3. Journal of Immunology, 2006, 177, 7656-7664.	0.4	52
170	Polycomb directs timely activation of germline genes in spermatogenesis. Genes and Development, 2017, 31, 1693-1703.	2.7	52
171	Histone H2A T120 Phosphorylation Promotes Oncogenic Transformation via Upregulation of Cyclin D1. Molecular Cell, 2016, 64, 176-188.	4.5	51
172	Combined Cohesin-RUNX1 Deficiency Synergistically Perturbs Chromatin Looping and Causes Myelodysplastic Syndromes. Cancer Discovery, 2020, 10, 836-853.	7.7	51
173	Microglia integration into human midbrain organoids leads to increased neuronal maturation and functionality. Glia, 2022, 70, 1267-1288.	2.5	51
174	A new Pax gene, Pax-9, maps to mouse Chromosome 12. Mammalian Genome, 1993, 4, 354-358.	1.0	50
175	Cloning and chromosome mapping of the human Mel-18 gene which encodes a DNA-binding protein with a new RING-finger motif. Gene, 1993, 129, 249-255.	1.0	50
176	Inductive Signals from the Somatopleure Mediated by Bone Morphogenetic Proteins Are Essential for the Formation of the Sternal Component of Avian Ribs. Developmental Biology, 2001, 232, 284-300.	0.9	50
177	CCN3 Protein Participates in Bone Regeneration as an Inhibitory Factor. Journal of Biological Chemistry, 2013, 288, 19973-19985.	1.6	50
178	Lethal myelofibrosis induced by Bmi1-deficient hematopoietic cells unveils a tumor suppressor function of the polycomb group genes. Journal of Experimental Medicine, 2012, 209, 445-454.	4.2	49
179	NOV/CCN3: A New Adipocytokine Involved in Obesity-Associated Insulin Resistance. Diabetes, 2016, 65, 2502-2515.	0.3	48
180	topors, a p53 and topoisomerase I-binding RING finger protein, is a coactivator of p53 in growth suppression induced by DNA damage. Oncogene, 2005, 24, 3385-3396.	2.6	46

#	ARTICLE	IF	CITATIONS
181	The Polycomb Group Protein SUZ12 regulates histone H3 lysine 9 methylation and HP1 \pm distribution. <i>Chromosome Research</i> , 2007, 15, 299-314.	1.0	46
182	Nuclear Inhibitor of Protein Phosphatase-1 (NIPP1) Directs Protein Phosphatase-1 (PP1) to Dephosphorylate the U2 Small Nuclear Ribonucleoprotein Particle (snRNP) Component, Spliceosome-associated Protein 155 (Sap155). <i>Journal of Biological Chemistry</i> , 2008, 283, 35805-35814.	1.6	46
183	Ring1a/b polycomb proteins regulate the mesenchymal stem cell niche in continuously growing incisors. <i>Developmental Biology</i> , 2012, 367, 140-153.	0.9	46
184	Histone demethylase Jmjd3 is required for the development of subsets of retinal bipolar cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 3751-3756.	3.3	45
185	Roles of histone H3K27 trimethylase Ezh2 in retinal proliferation and differentiation. <i>Developmental Neurobiology</i> , 2015, 75, 947-960.	1.5	45
186	Requirement of zinc transporter ZIP10 for epidermal development: Implication of the ZIP10-p63 axis in epithelial homeostasis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, 12243-12248.	3.3	45
187	MBLR, a new RING finger protein resembling mammalian Polycomb gene products, is regulated by cell cycle-dependent phosphorylation. <i>Genes To Cells</i> , 2002, 7, 835-850.	0.5	44
188	Presenilin 1 is essential for cardiac morphogenesis. <i>Developmental Dynamics</i> , 2004, 230, 795-799.	0.8	44
189	Accumulation of p27KIP1 is associated with BMP2-induced growth arrest and neuronal differentiation of human neuroblastoma-derived cell lines. <i>Biochemical and Biophysical Research Communications</i> , 2003, 307, 206-213.	1.0	43
190	Reproductive Isolation in Hybrid Mice Due to Spermatogenesis Defects at Three Meiotic Stages. <i>Genetics</i> , 2010, 186, 339-351.	1.2	43
191	CNOT3 contributes to early B cell development by controlling <i>Igh</i> rearrangement and <i>p53</i> mRNA stability. <i>Journal of Experimental Medicine</i> , 2015, 212, 1465-1479.	4.2	43
192	Internal deletion of BCOR reveals a tumor suppressor function for BCOR in T lymphocyte malignancies. <i>Journal of Experimental Medicine</i> , 2017, 214, 2901-2913.	4.2	43
193	Dependence receptor UNC5D mediates nerve growth factor depletion-induced neuroblastoma regression. <i>Journal of Clinical Investigation</i> , 2013, 123, 2935-2947.	3.9	43
194	Fc receptor β 2 subunit is required for full activation of mast cells through Fc receptor engagement. <i>International Immunology</i> , 1999, 11, 199-207.	1.8	42
195	Genome-wide analysis of target genes regulated by HoxB4 in hematopoietic stem and progenitor cells developing from embryonic stem cells. <i>Blood</i> , 2011, 117, e142-e150.	0.6	42
196	Loss of Ezh2 promotes a midbrain-to-forebrain identity switch by direct gene derepression and Wnt-dependent regulation. <i>BMC Biology</i> , 2015, 13, 103.	1.7	42
197	Extrathymic Differentiation of a T Cell Bearing Invariant V β 14J β 281 TCR. <i>International Reviews of Immunology</i> , 1994, 11, 31-46.	1.5	41
198	Overlapping Roles of the Methylated DNA-binding Protein MBD1 and Polycomb Group Proteins in Transcriptional Repression of HOXA Genes and Heterochromatin Foci Formation. <i>Journal of Biological Chemistry</i> , 2007, 282, 16391-16400.	1.6	41

#	ARTICLE	IF	CITATIONS
199	Identification of loss-of-function mutations of SLC35D1 in patients with Schneckenbecken dysplasia, but not with other severe spondylodysplastic dysplasias group diseases. <i>Journal of Medical Genetics</i> , 2009, 46, 562-568.	1.5	41
200	Divergence of CpG island promoters: A consequence or cause of evolution?. <i>Development Growth and Differentiation</i> , 2010, 52, 545-554.	0.6	41
201	Human NK cell development in hIL-7 and hIL-15 knockin NOD/SCID/IL2rgKO mice. <i>Life Science Alliance</i> , 2019, 2, e201800195.	1.3	41
202	Gab2, via PI-3K, Regulates ARF1 in Fc μ RI-Mediated Granule Translocation and Mast Cell Degranulation. <i>Journal of Immunology</i> , 2011, 187, 932-941.	0.4	40
203	Bmi1 Confers Resistance to Oxidative Stress on Hematopoietic Stem Cells. <i>PLoS ONE</i> , 2012, 7, e36209.	1.1	40
204	Pax-1 in the development of the cervico-occipital transitional zone. <i>Anatomy and Embryology</i> , 1995, 192, 221-7.	1.5	38
205	Gene Resistance to Transcriptional Reprogramming following Nuclear Transfer Is Directly Mediated by Multiple Chromatin-Repressive Pathways. <i>Molecular Cell</i> , 2017, 65, 873-884.e8.	4.5	38
206	Impaired cell cycle control of neuronal precursor cells in the neocortical primordium of presenilin-1-deficient mice. <i>Journal of Neuroscience Research</i> , 2002, 70, 501-513.	1.3	36
207	Generation of functional NKT cells in vitro from embryonic stem cells bearing rearranged invariant V β 14-J β 18 TCR β gene. <i>Blood</i> , 2010, 115, 230-237.	0.6	36
208	Depletion of Sf3b1 impairs proliferative capacity of hematopoietic stem cells but is not sufficient to induce myelodysplasia. <i>Blood</i> , 2014, 123, 3336-3343.	0.6	36
209	Bcor insufficiency promotes initiation and progression of myelodysplastic syndrome. <i>Blood</i> , 2018, 132, 2470-2483.	0.6	36
210	PRC1 preserves epidermal tissue integrity independently of PRC2. <i>Genes and Development</i> , 2019, 33, 55-60.	2.7	36
211	An essential role for REV3 in mammalian cell survival: absence of REV3 induces p53-independent embryonic death. <i>Biochemical and Biophysical Research Communications</i> , 2002, 293, 1132-1137.	1.0	35
212	Distinct roles of Polycomb group gene products in transcriptionally repressed and active domains of Hoxb8. <i>Development (Cambridge)</i> , 2006, 133, 2371-2381.	1.2	35
213	KDEL receptor 1 regulates T-cell homeostasis via PP1 that is a key phosphatase for ISR. <i>Nature Communications</i> , 2015, 6, 7474.	5.8	35
214	Generation of Tumor Antigen-Specific iPSC-Derived Thymic Emigrants Using a 3D Thymic Culture System. <i>Cell Reports</i> , 2018, 22, 3175-3190.	2.9	35
215	Expression of the Dan gene during chicken embryonic development. <i>Mechanisms of Development</i> , 2001, 109, 363-365.	1.7	34
216	Late-Onset of Spinal Neurodegeneration in Knock-In Mice Expressing a Mutant BiP. <i>PLoS ONE</i> , 2014, 9, e112837.	1.1	34

#	ARTICLE	IF	CITATIONS
217	A Novel Zinc Finger Protein Zfp277 Mediates Transcriptional Repression of the Ink4a/Arf Locus through Polycomb Repressive Complex 1. PLoS ONE, 2010, 5, e12373.	1.1	33
218	Histone acetylation mediated by Brd1 is crucial for Cd8 gene activation during early thymocyte development. Nature Communications, 2014, 5, 5872.	5.8	33
219	Pramel7 mediates ground-state pluripotency through proteasomal epigenetic combined pathways. Nature Cell Biology, 2017, 19, 763-773.	4.6	33
220	Involvement of the Polycomb-group gene Ring1B in the specification of the anterior-posterior axis in mice. Development (Cambridge), 2002, 129, 4171-83.	1.2	32
221	Dissociation of mammalian Polycomb-group proteins, Ring1B and Rae28/Ph1, from the chromatin correlates with configuration changes of the chromatin in mitotic and meiotic prophase. Histochemistry and Cell Biology, 2003, 120, 111-119.	0.8	31
222	Is there a role for endogenous retroviruses to mediate long-term adaptive phenotypic response upon environmental inputs?. Philosophical Transactions of the Royal Society B: Biological Sciences, 2013, 368, 20110340.	1.8	30
223	Bromodomain-PHD finger protein 1 is critical for leukemogenesis associated with MOZ-TIF2 fusion. International Journal of Hematology, 2014, 99, 21-31.	0.7	30
224	Recruitment of Dnmt1. Progress in Molecular Biology and Translational Science, 2011, 101, 289-310.	0.9	29
225	Polycomb Group Protein Ezh2 Regulates Hepatic Progenitor Cell Proliferation and Differentiation in Murine Embryonic Liver. PLoS ONE, 2014, 9, e104776.	1.1	29
226	Conversion of T cells to B cells by inactivation of polycomb-mediated epigenetic suppression of the B-lineage program. Genes and Development, 2016, 30, 2475-2485.	2.7	29
227	Analysis of LacZ reporter genes in transgenic embryos suggests the presence of several cis-acting regulatory elements in the murine Hoxb-6 gene. Developmental Dynamics, 1993, 196, 205-216.	0.8	28
228	A gene targeting approach for functional characterization of KIAA genes encoding extremely large proteins. FASEB Journal, 2006, 20, 1718-1720.	0.2	28
229	MAPKAP kinase MK2 maintains self-renewal capacity of haematopoietic stem cells. EMBO Journal, 2009, 28, 1392-1406.	3.5	28
230	Samd7 is a cell type-specific PRC1 component essential for establishing retinal rod photoreceptor identity. Proceedings of the National Academy of Sciences of the United States of America, 2017, 114, E8264-E8273.	3.3	28
231	Mast cells play role in wound healing through the ZnT2/GPR39/IL-6 axis. Scientific Reports, 2019, 9, 10842.	1.6	28
232	<i>Irx1</i> mutant mice show reduced tendon differentiation and no patterning defects in musculoskeletal system development. Genesis, 2011, 49, 2-9.	0.8	27
233	Guanine- 5-carboxylcytosine base pairs mimic mismatches during DNA replication. Scientific Reports, 2014, 4, 5220.	1.6	27
234	Prediction of the Coding Sequences of Mouse Homologues of KIAA Gene: III. The Complete Nucleotide Sequences of 500 Mouse KIAA-homologous cDNAs Identified by Screening of Terminal Sequences of cDNA Clones Randomly Sampled from Size-fractionated Libraries. DNA Research, 2003, 10, 167-180.	1.5	26

#	ARTICLE	IF	CITATIONS
235	Bmi1 regulates cell fate via tumor suppressor <i>Wwox</i> repression in small cell lung cancer cells. <i>Cancer Science</i> , 2011, 102, 983-990.	1.7	26
236	KDM2 proteins constrain transcription from CpG island gene promoters independently of their histone demethylase activity. <i>Nucleic Acids Research</i> , 2019, 47, 9005-9023.	6.5	26
237	<i>Staphylococcus cohnii</i> is a potentially biotherapeutic skin commensal alleviating skin inflammation. <i>Cell Reports</i> , 2021, 35, 109052.	2.9	26
238	Generation of Novel Traj18-Deficient Mice Lacking ± 14 Natural Killer T Cells with an Undisturbed T Cell Receptor γ -Chain Repertoire. <i>PLoS ONE</i> , 2016, 11, e0153347.	1.1	26
239	Repression of germline genes by PRC1.6 and SETDB1 in the early embryo precedes DNA methylation-mediated silencing. <i>Nature Communications</i> , 2021, 12, 7020.	5.8	26
240	Identification, Tissue Expression, and Functional Characterization of Otx3, a Novel Member of the Otx Family. <i>Journal of Biological Chemistry</i> , 2002, 277, 28065-28069.	1.6	25
241	Induction of Wnt5a-Expressing Mesenchymal Cells Adjacent to the Cloacal Plate Is an Essential Process for Its Proximodistal Elongation and Subsequent Anorectal Development. <i>Pediatric Research</i> , 2009, 66, 149-154.	1.1	25
242	Reduced NOV/CCN3 Expression Limits Inflammation and Interstitial Renal Fibrosis after Obstructive Nephropathy in Mice. <i>PLoS ONE</i> , 2015, 10, e0137876.	1.1	25
243	Polycomb complexes redundantly maintain epidermal stem cell identity during development. <i>Genes and Development</i> , 2021, 35, 354-366.	2.7	25
244	Variant PCGF1-PRC1 links PRC2 recruitment with differentiation-associated transcriptional inactivation at target genes. <i>Nature Communications</i> , 2021, 12, 5341.	5.8	25
245	Characterization of Size-Fractionated cDNA Libraries Generated by the in vitro Recombination-Assisted Method. <i>DNA Research</i> , 2002, 9, 47-57.	1.5	24
246	Mesenchymal expression of Foxl1, a winged helix transcriptional factor, regulates generation and maintenance of gut-associated lymphoid organs. <i>Developmental Biology</i> , 2003, 255, 278-289.	0.9	24
247	Foxl1-deficient mice exhibit aberrant epithelial cell positioning resulting from dysregulated EphB/EphrinB expression in the small intestine. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 291, G163-G170.	1.6	24
248	Transcription factors Mesp2 and Paraxis have critical roles in axial musculoskeletal formation. <i>Developmental Dynamics</i> , 2007, 236, 1484-1494.	0.8	24
249	Dual Functions of the RFTS Domain of Dnmt1 in Replication-Coupled DNA Methylation and in Protection of the Genome from Aberrant Methylation. <i>PLoS ONE</i> , 2015, 10, e0137509.	1.1	24
250	Elbow knee synostosis (Eks): a new mutation on mouse Chromosome 14. <i>Mammalian Genome</i> , 2002, 13, 341-344.	1.0	23
251	Polycomb in Transcriptional Phase Transition of Developmental Genes. <i>Trends in Biochemical Sciences</i> , 2016, 41, 9-19.	3.7	23
252	<i>Polycomb</i> Group Gene Product Ring1B Regulates Th2-Driven Airway Inflammation through the Inhibition of Bim-Mediated Apoptosis of Effector Th2 Cells in the Lung. <i>Journal of Immunology</i> , 2010, 184, 4510-4520.	0.4	22

#	ARTICLE	IF	CITATIONS
253	Splicing factor 3b subunit 1 (Sf3b1) haploinsufficient mice display features of low risk Myelodysplastic syndromes with ring sideroblasts. <i>Journal of Hematology and Oncology</i> , 2014, 7, 89.	6.9	22
254	KDM2B in polycomb repressive complex 1.1 functions as a tumor suppressor in the initiation of T-cell leukemogenesis. <i>Blood Advances</i> , 2019, 3, 2537-2549.	2.5	22
255	FBXL19 recruits CDK-Mediator to CpG islands of developmental genes priming them for activation during lineage commitment. <i>ELife</i> , 2018, 7, .	2.8	22
256	Clustering of CARMA1 through SH3â€“GUK domain interactions is required for its activation of NF-Î²B signalling. <i>Nature Communications</i> , 2015, 6, 5555.	5.8	21
257	Polycomb group molecule PHC3 regulates polycomb complex composition and prognosis of osteosarcoma. <i>Cancer Science</i> , 2010, 101, 1646-1652.	1.7	20
258	Presenilin-1 controls the growth and differentiation of endothelial progenitor cells through its Î²-catenin-binding region. <i>Cell Biology International</i> , 2006, 30, 239-243.	1.4	19
259	VÎ±14+ NK T cells: A novel lymphoid cell lineage with regulatory function. <i>Journal of Allergy and Clinical Immunology</i> , 1996, 98, S263-S269.	1.5	18
260	Characterization of Np95 expression in mouse brain from embryo to adult: A novel marker for proliferating neural stem/precursor cells. <i>Neurogenesis (Austin, Tex)</i> , 2014, 1, e976026.	1.5	18
261	The Polycomb group protein Ring1 regulates dorsoventral patterning of the mouse telencephalon. <i>Nature Communications</i> , 2020, 11, 5709.	5.8	18
262	Randomly inserted and targeted Hox/reporter fusions transcriptionally silenced in Polycomb mutants. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 13362-13367.	3.3	17
263	Ring1B promotes hepatic stem/progenitor cell expansion through simultaneous suppression of Cdkn1a and Cdkn2a in mice. <i>Hepatology</i> , 2014, 60, 323-333.	3.6	16
264	Generating a transgenic mouse line stably expressing human MHC surface antigen from a HAC carrying multiple genomic BACs. <i>Chromosoma</i> , 2015, 124, 107-118.	1.0	16
265	DNMTs and SETDB1 function as co-repressors in MAX-mediated repression of germ cellâ€“related genes in mouse embryonic stem cells. <i>PLoS ONE</i> , 2018, 13, e0205969.	1.1	16
266	The USP21 Short Variant (USP21SV) Lacking NES, Located Mostly in the Nucleus In Vivo, Activates Transcription by Deubiquitylating ubH2A In Vitro. <i>PLoS ONE</i> , 2013, 8, e79813.	1.1	16
267	Loss of Pcgf5 Affects Global H2A Monoubiquitination but Not the Function of Hematopoietic Stem and Progenitor Cells. <i>PLoS ONE</i> , 2016, 11, e0154561.	1.1	16
268	Establishment of mouse stem cells that can recapitulate the developmental potential of primitive endoderm. <i>Science</i> , 2022, 375, 574-578.	6.0	16
269	Monoclonal antibody against murine T cell receptor VÎ±14 cross-reacts with human CD3Î± and detects disulfide-linked dimeric form. <i>International Immunology</i> , 1991, 3, 991-995.	1.8	15
270	Mesenchymal Transcription Factor Fkh6 Is Essential for the Development and Differentiation of Parietal Cells. <i>Biochemical and Biophysical Research Communications</i> , 2001, 280, 1069-1076.	1.0	15

#	ARTICLE	IF	CITATIONS
271	Bone Marrow Allograft Rejection Mediated by a Novel Murine NK Receptor, NKG2I. <i>Journal of Experimental Medicine</i> , 2004, 199, 137-144.	4.2	15
272	Insulin resistance and increased pancreatic β -cell proliferation in mice expressing a mutant insulin receptor (P1195L). <i>Journal of Endocrinology</i> , 2006, 190, 739-747.	1.2	15
273	Induced pluripotency as a potential path towards iNKT cell-mediated cancer immunotherapy. <i>International Journal of Hematology</i> , 2012, 95, 624-631.	0.7	15
274	RING1 contributes to early proximal-distal specification of the forelimb bud by restricting Meis2 expression. <i>Development (Cambridge)</i> , 2015, 143, 276-85.	1.2	15
275	Ring1A and Ring1B inhibit expression of Glis2 to maintain murine MOZ-TIF2 AML stem cells. <i>Blood</i> , 2018, 131, 1833-1845.	0.6	15
276	Variant PRC1 competes with retinoic acid-related signals to repress <i>Meis2</i> in distal forelimb bud. <i>Development (Cambridge)</i> , 2018, 145, .	1.2	15
277	Maintenance DNA methylation in pre-meiotic germ cells regulates meiotic prophase by facilitating homologous chromosome pairing. <i>Development (Cambridge)</i> , 2021, 148, .	1.2	15
278	Human cell clones, R _{Sa} and U _{Vr} -1, differing in their capability for UV-induced virus reactivation and phenotypic mutation. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1994, 306, 211-222.	0.4	14
279	Heat shock-induced homeotic transformations of the axial skeleton and associated shifts of Hox gene expression domains in mouse embryos. <i>Reproductive Toxicology</i> , 1997, 11, 761-770.	1.3	14
280	Molecular cloning, genetic mapping, and expression of the mouse Sf3b1 (SAP155) gene for the U2 snRNP component of spliceosome. <i>Mammalian Genome</i> , 2001, 12, 192-198.	1.0	14
281	Model mice for Presbyterian hemoglobinopathy (Asn ¹⁰⁸ →Lys) confer hemolytic anemia with altered oxygen affinity and instability of Hb. <i>Biochemical and Biophysical Research Communications</i> , 2002, 295, 869-876.	1.0	14
282	Prediction of the Coding Sequences of Mouse Homologues of KIAA Gene: IV. The Complete Nucleotide Sequences of 500 Mouse KIAA-homologous cDNAs Identified by Screening of Terminal Sequences of cDNA Clones Randomly Sampled from Size-Fractionated Libraries. <i>DNA Research</i> , 2004, 11, 205-218.	1.5	14
283	Nuclear transcriptome profiling of induced pluripotent stem cells and embryonic stem cells identify non-coding loci resistant to reprogramming. <i>Cell Cycle</i> , 2015, 14, 1148-1155.	1.3	14
284	Polycomb complexes regulate cellular senescence by repression of <i>ARF</i> in cooperation with E2F3. <i>Genes To Cells</i> , 2007, 12, 1371-1382.	0.5	13
285	UV-induced reduction in Polycomb repression promotes epidermal pigmentation. <i>Developmental Cell</i> , 2021, 56, 2547-2561.e8.	3.1	13
286	Osteoclasts adapt to physioxia perturbation through DNA demethylation. <i>EMBO Reports</i> , 2021, 22, e53035.	2.0	13
287	Role of Polycomb RYBP in Maintaining the B-1-to-B-2 B-Cell Lineage Switch in Adult Hematopoiesis. <i>Molecular and Cellular Biology</i> , 2016, 36, 900-912.	1.1	12
288	Double-step and inverse polymerase chain reaction for sensitive detection and cloning of T cell receptor variable region sequences. <i>International Immunology</i> , 1991, 3, 1053-1057.	1.8	11

#	ARTICLE	IF	CITATIONS
289	Fine genetic mapping of the proximal part of mouse Chromosome 2 excludes Pax-8 as a candidate gene for Danforth's short tail (Sd). <i>Mammalian Genome</i> , 1993, 4, 324-327.	1.0	11
290	Reduction of SNAP25 in acid secretion defect of Foxl1 ^{-/-} gastric parietal cells. <i>Biochemical and Biophysical Research Communications</i> , 2004, 320, 766-772.	1.0	11
291	Hemimethylation: DNA's lasting odd couple. <i>Science</i> , 2018, 359, 1102-1103.	6.0	11
292	Polycomb Repressive Complex 1 Controls Maintenance of Fungiform Papillae by Repressing Sonic Hedgehog Expression. <i>Cell Reports</i> , 2019, 28, 257-266.e5.	2.9	11
293	Discovery of widespread transcription initiation at microsatellites predictable by sequence-based deep neural network. <i>Nature Communications</i> , 2021, 12, 3297.	5.8	11
294	Bivalent-histone-marked immediate-early gene regulation is vital for VEGF-responsive angiogenesis. <i>Cell Reports</i> , 2022, 38, 110332.	2.9	11
295	The mouse <i>Edr2</i> (<i>Mph2</i>) gene has two forms of mRNA encoding 90- and 36-kDa polypeptides. <i>Gene</i> , 2002, 288, 103-110.	1.0	10
296	Expression of recombination-activating gene in mature peripheral T cells in Peyer's patch. <i>International Immunology</i> , 2003, 15, 393-402.	1.8	10
297	<i>Phc2</i> controls hematopoietic stem and progenitor cell mobilization from bone marrow by repressing <i>Vcam1</i> expression. <i>Nature Communications</i> , 2019, 10, 3496.	5.8	10
298	The <i>Cxxc1</i> subunit of the Trithorax complex directs epigenetic licensing of CD4+ T cell differentiation. <i>Journal of Experimental Medicine</i> , 2021, 218, .	4.2	10
299	The mouse <i>YAF2</i> gene generates two distinct transcripts and is expressed in pre-and postimplantation embryos. <i>Gene</i> , 2003, 315, 183-192.	1.0	9
300	Enhanced Th2 Cell Differentiation and Allergen-Induced Airway Inflammation in <i>Zfp35</i> -Deficient Mice. <i>Journal of Immunology</i> , 2009, 183, 5388-5396.	0.4	9
301	<i>DJ-1</i> depletion prevents immunoaging in T cell compartments. <i>EMBO Reports</i> , 2022, 23, e53302.	2.0	9
302	Autoreactive T-cell clones which suppress cytotoxic T cell responses. <i>International Immunology</i> , 1991, 3, 377-384.	1.8	8
303	Embracing change to remain the same: conservation of polycomb functions despite divergence of binding motifs among species. <i>Current Opinion in Cell Biology</i> , 2013, 25, 305-313.	2.6	8
304	Role of a NK receptor, KLRE-1, in bone marrow allograft rejection: analysis with KLRE-1-deficient mice. <i>Blood</i> , 2004, 104, 781-783.	0.6	7
305	Mammalian Polycomb complexes are required for Peyer's patch development by regulating lymphoid cell proliferation. <i>Gene</i> , 2006, 379, 166-174.	1.0	7
306	Unique Composition of Polycomb Repressive Complex 1 in Hematopoietic Stem Cells. <i>International Journal of Hematology</i> , 2007, 85, 179-181.	0.7	7

#	ARTICLE	IF	CITATIONS
307	Epigenetic Memory Meets G2/M: To Remember or To Forget?. <i>Developmental Cell</i> , 2011, 20, 5-6.	3.1	7
308	Foxc2 in pharyngeal arch mesenchyme is important for aortic arch artery remodelling and ventricular septum formation. <i>Biomedical Research</i> , 2015, 36, 235-245.	0.3	7
309	Identification of the Coiled-Coil Domain as an Essential Methyl-CpG-Binding Domain Protein 3 Element for Preserving Lineage Commitment Potential of Embryonic Stem Cells. <i>Stem Cells</i> , 2018, 36, 1355-1367.	1.4	7
310	Lack of whey acidic protein four disulphide core (WFDC) 2 protease inhibitor causes neonatal death from respiratory failure in mice. <i>DMM Disease Models and Mechanisms</i> , 2019, 12, .	1.2	7
311	Generation of induced pluripotent stem cell-derived mice by reprogramming of a mature NKT cell. <i>International Immunology</i> , 2014, 26, 551-561.	1.8	6
312	PCL2 regulates p53 stability and functions as a tumor suppressor in breast cancer. <i>Science Bulletin</i> , 2018, 63, 629-639.	4.3	6
313	Prediction of the Coding Sequences of Mouse Homologues of FLJ Genes: The Complete Nucleotide Sequences of 110 Mouse FLJ-Homologous cDNAs Identified by Screening of Terminal Sequences of cDNA Clones Randomly Sampled from Size-Fractionated Libraries. <i>DNA Research</i> , 2004, 11, 127-135.	1.5	5
314	Mammal-specific H2A Variant, H2ABbd, Is Involved in Apoptotic Induction via Activation of NF- κ B Signaling Pathway. <i>Journal of Biological Chemistry</i> , 2014, 289, 11656-11666.	1.6	5
315	Physical interaction between MPP8 and PRC1 complex and its implication for regulation of spermatogenesis. <i>Biochemical and Biophysical Research Communications</i> , 2015, 458, 470-475.	1.0	5
316	Live Imaging of Xist RNA. <i>Methods in Molecular Biology</i> , 2018, 1861, 67-72.	0.4	5
317	Isolation of Genomic DNA Controlling Mouse Melanoma Antigen Defined by Monoclonal Antibody. <i>Japanese Journal of Cancer Research</i> , 1988, 79, 718-725.	1.7	4
318	Reduced inflammatory pain in mice deficient in the differential screening-selected gene aberrative in neuroblastoma. <i>Molecular and Cellular Neurosciences</i> , 2004, 25, 504-514.	1.0	4
319	Insufficiency of non-canonical PRC1 synergizes with JAK2V617F in the development of myelofibrosis. <i>Leukemia</i> , 2021, .	3.3	4
320	Polycomb group ring finger protein 6 suppresses Myc-induced lymphomagenesis. <i>Life Science Alliance</i> , 2022, 5, e202101344.	1.3	4
321	TCR repertoire in early fetal mouse thymus. <i>International Immunology</i> , 1995, 7, 493-499.	1.8	3
322	A unique expression pattern of <i>Tbx10</i> in the hindbrain as revealed by <i>Tbx10^{sup>LacZ</sup>}</i> allele. <i>Genesis</i> , 2010, 48, 295-302.	0.8	3
323	Regulation of Fetal Genes by Transitions among RNA-Binding Proteins during Liver Development. <i>International Journal of Molecular Sciences</i> , 2020, 21, 9319.	1.8	3
324	Efficient Production of Functional Human NKT Cells from Induced Pluripotent Stem Cells $\hat{\wedge}$ Reprogramming of Human $\hat{\wedge}$ 24+iNKT Cells. <i>Bio-protocol</i> , 2017, 7, e2277.	0.2	3

#	ARTICLE	IF	CITATIONS
325	Microbiota-Independent Spontaneous Dermatitis Associated with Increased Sebaceous Lipid Production in Tmem79-Deficient Mice. <i>Journal of Investigative Dermatology</i> , 2022, 142, 2864-2872.e6.	0.3	3
326	Expression of the Proliferation-Related Ki-67 mRNA in the Early Development of Murine Embryo. <i>Biochemical and Biophysical Research Communications</i> , 1997, 235, 191-196.	1.0	2
327	Generation of a Monoclonal Antibody Against the Mouse Sf3b1 (SAP155) Gene Product for U2 snRNP Component of Spliceosome. <i>Hybridoma</i> , 2003, 22, 117-119.	0.6	2
328	Mammalian Polycomb-Like Pcl2/Mtf2 Is a Novel Regulatory Component of PRC2 That Can Differentially Modulate Polycomb Activity both at the <i>Hox</i> Gene Cluster and at <i>Cdkn2a</i> Genes. <i>Molecular and Cellular Biology</i> , 2014, 34, 2773-2773.	1.1	2
329	Sublethal endoplasmic reticulum stress caused by the mutation of immunoglobulin heavy chain-binding protein induces the synthesis of a mitochondrial protein, pyrroline-5-carboxylate reductase 1. <i>Cell Stress and Chaperones</i> , 2017, 22, 77-85.	1.2	2
330	Rewriting the past: de novo activity of PRC2 restores global H3K27 methylation patterns. <i>Nature Structural and Molecular Biology</i> , 2018, 25, 197-199.	3.6	2
331	Np95/Uhrf1 regulates tumor suppressor gene expression of neural stem/precursor cells, contributing to neurogenesis in the adult mouse brain. <i>Neuroscience Research</i> , 2019, 143, 31-43.	1.0	2
332	Association of SF3B1 with Ring Sideroblasts in patients, In Vivo, and In Vitro models of Spliceosomal Dysfunction. <i>Blood</i> , 2011, 118, 457-457.	0.6	2
333	Splicing Factor 3b Subunit 1 (SF3B1) Heterozygous Mice Manifest a Hematologic Phenotype Similar To Low Risk Myelodysplastic Syndromes With Ring Sideroblasts. <i>Blood</i> , 2013, 122, 259-259.	0.6	2
334	Molecular Analysis of Suppressor T Cell Receptors. <i>International Reviews of Immunology</i> , 1988, 3, 229-239.	1.5	1
335	Method of Genomic DNA Cloning by the Combination of Cosmid Shuttle Vector and Monoclonal Antibody. <i>Microbiology and Immunology</i> , 1988, 32, 1073-1078.	0.7	1
336	Accelerated acquisition of permeability barrier function in the skin of presenilin-1-deficient embryos. <i>Archives of Dermatological Research</i> , 2006, 298, 339-345.	1.1	1
337	Epigenetics in development: Decorating the genome to show or hide?. <i>Development Growth and Differentiation</i> , 2010, 52, 481-481.	0.6	1
338	O12. Polycomb-dependent regulation for differentiation programs of stem cells and progenitors. <i>Differentiation</i> , 2010, 80, S9-S10.	1.0	1
339	Stem Cell Epigenetics: Looking Forward. <i>Cell Stem Cell</i> , 2014, 14, 706-709.	5.2	1
340	No Winter Lasts Forever: Polycomb Complexes Convert Epigenetic Memory of Cold into Flowering. <i>Developmental Cell</i> , 2017, 42, 563-564.	3.1	1
341	Role Of Sf3b1 On Hematopoiesis. <i>Blood</i> , 2013, 122, 600-600.	0.6	1
342	Genomic DNA with Transformation-Related Activity and Melanoma Antigen Expression. <i>Journal of Investigative Dermatology</i> , 1989, 92, S284-S288.	0.3	0

#	ARTICLE	IF	CITATIONS
343	Genomic DNA with Transformation-Related Activity and Melanoma Antigen Expression.. Journal of Investigative Dermatology, 1989, 92, 284S-288S.	0.3	0
344	V-Alpha-14 T-Cell Antigen Receptor Repertoire and Its Function. International Archives of Allergy and Immunology, 1992, 99, 416-418.	0.9	0
345	Production of Monoclonal Antibodies against Quail Pax-1. Hybridoma, 1996, 15, 83-86.	0.9	0
346	An interview with Haruhiko Koseki. Development (Cambridge), 2012, 139, 3469-3470.	1.2	0
347	Distinct roles of Polycomb group gene products in transcriptionally repressed and active domains of <i>Hoxb8</i> . Development (Cambridge), 2014, 141, 3431-3436.	1.2	0
348	Role of the polycomb gene BCOR in hematopoiesis. Experimental Hematology, 2015, 43, S83.	0.2	0
349	Biological and genetic characterization of the role of SRSF2 mutations in the pathogenesis of myelodysplastic syndromes. Experimental Hematology, 2015, 43, S73.	0.2	0
350	Lethal myelofibrosis induced by Bmi1-deficient hematopoietic cells unveils a tumor suppressor function of the polycomb group genes. Journal of Cell Biology, 2012, 196, i5-i5.	2.3	0