

Eduardo Reyes

List of Publications by Citations

Source: <https://exaly.com/author-pdf/1643597/eduardo-reyes-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

20
papers

106
citations

7
h-index

9
g-index

21
ext. papers

123
ext. citations

2.2
avg, IF

2.37
L-index

#	Paper	IF	Citations
20	Circulating Prostate Cells Found in Men with Benign Prostate Disease Are P504S Negative: Clinical Implications. <i>Journal of Oncology</i> , 2013 , 2013, 165014	4.5	18
19	Minimum Residual Disease in Patients Post Radical Prostatectomy for Prostate Cancer: Theoretical Considerations, Clinical Implications and Treatment Outcome. <i>Asian Pacific Journal of Cancer Prevention</i> , 2018 , 19, 229-236	1.7	11
18	Prediction model for early biochemical recurrence after radical prostatectomy based on the Cancer of the Prostate Risk Assessment score and the presence of secondary circulating prostate cells. <i>BJU International</i> , 2016 , 118, 556-62	5.6	11
17	A comparative performance analysis of total PSA, percentage free PSA, PSA velocity, and PSA density versus the detection of primary circulating prostate cells in predicting initial prostate biopsy findings in Chilean men. <i>BioMed Research International</i> , 2014 , 2014, 676572	3	10
16	Primary circulating prostate cells are not detected in men with low grade small volume prostate cancer. <i>Journal of Oncology</i> , 2014 , 2014, 612674	4.5	8
15	Predictive Value of Neutrophil to Lymphocyte Ratio in the Diagnosis of Significant Prostate Cancer at Initial Biopsy: A Comparison with Free Percent Prostate Specific Antigen, Prostate Specific Antigen Density and Primary Circulating Prostate Cells. <i>Asian Pacific Journal of Cancer Prevention</i> , 2019 , 20, 3385-3389	1.7	7
14	Platelet-to-lymphocyte ratio and systemic immune-inflammation index versus circulating prostate cells to predict significant prostate cancer at first biopsy. <i>Turkish Journal of Urology</i> , 2020 , 46, 115-122	1.3	7
13	The presence of primary circulating prostate cells is associated with upgrading and upstaging in patients eligible for active surveillance. <i>Ecancermedicalscience</i> , 2017 , 11, 711	2.7	6
12	Limited improvement of incorporating primary circulating prostate cells with the CAPRA score to predict biochemical failure-free outcome of radical prostatectomy for prostate cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2016 , 34, 430.e17-25	2.8	6
11	Subtypes of minimal residual disease, association with Gleason score, risk and time to biochemical failure in pT2 prostate cancer treated with radical prostatectomy. <i>Ecancermedicalscience</i> , 2019 , 13, 934	2.7	5
10	Efficacy of Using Sequential Primary Circulating Prostate Cell Detection for Initial Prostate Biopsy in Men Suspected of Prostate Cancer. <i>Asian Pacific Journal of Cancer Prevention</i> , 2016 , 17, 3385-90	1.7	3
9	Head-to-head comparison of the Montreal nomogram with the detection of primary malignant circulating prostate cells to predict prostate cancer at initial biopsy in Chilean men with suspicion of prostate cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015 , 33, 203.e19-25	2.8	2
8	How localized is pathologically localized prostate cancer? The use of secondary circulating prostate cells as a marker of minimal residual disease and their association with patient outcome. <i>Turkish Journal of Urology</i> , 2017 , 43, 456-461	1.3	2
7	Renal infarction associated with asymptomatic Covid-19 infection. <i>Hematology, Transfusion and Cell Therapy</i> , 2021 , 43, 353-356	1.6	2
6	Minimal Residual Disease Defines the Risk and Time to Biochemical Failure in Patients with Pt2 and Pt3a Prostate Cancer Treated With Radical Prostatectomy: An Observational Prospective Study. <i>Urology Journal</i> , 2020 , 17, 262-270	0.9	2
5	The Epstein criteria predict for organ-confined prostate cancer but not for minimal residual disease and outcome after radical prostatectomy. <i>Turkish Journal of Urology</i> , 2020 , 46, 360-366	1.3	1
4	The CAPRA score versus sub-types of minimal residual disease to predict biochemical failure after external beam radiotherapy. <i>Ecancermedicalscience</i> , 2020 , 14, 1042	2.7	1

- 3 Outcome of radical prostatectomy in primary circulating prostate cell negative prostate cancer. *Ecancermedicalscience*, **2016**, 10, 671 2.7 1
- 2 The Epstein criteria predict for organ-confined prostate cancer but not for minimal residual disease and outcome after radical prostatectomy. *Turkish Journal of Urology*, **2020**, 46, 360-366 1.3
- 1 The expression of matrix-metalloproteinase-2 in bone marrow micro-metastasis is associated with the presence of circulating prostate cells and a worse prognosis in men treated with radical prostatectomy for prostate cancer. *Turkish Journal of Urology*, **2020**, 46, 186-195 1.3