
Mario Santinami

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1646579/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Genetic Layout of Melanoma Lesions Is Associated with BRAF/MEK-Targeted Therapy Resistance and Transcriptional Profiles. Journal of Investigative Dermatology, 2022, 142, 3030-3040.e5.	0.7	6
2	The role of sentinel node tumor burden in modeling the prognosis of melanoma patients with positive sentinel node biopsy: an Italian melanoma intergroup study (N = 2,086). BMC Cancer, 2022, 22, .	2.6	5
3	Melanoma recurrence patterns and management after adjuvant targeted therapy: a multicentre analysis. British Journal of Cancer, 2021, 124, 574-580.	6.4	27
4	Surgical treatment of melanoma metastases to the small bowel: A single cancer referral center real-life experience. European Journal of Surgical Oncology, 2021, 47, 409-415.	1.0	3
5	The role of sentinel lymph node status performed in melanoma patients with local recurrence or in transit metastasis. European Journal of Surgical Oncology, 2021, 47, 1152-1156.	1.0	0
6	Survival in Patients With Sentinel Node–Positive Melanoma With Extranodal Extension. Journal of the National Comprehensive Cancer Network: JNCCN, 2021, 19, 1165-1173.	4.9	3
7	miR-146a-5p impairs melanoma resistance to kinase inhibitors by targeting COX2 and regulating NFkB-mediated inflammatory mediators. Cell Communication and Signaling, 2020, 18, 156.	6.5	18
8	Clinical applications of receptor-binding radiopharmaceutical 99mTc-Tilmanocept: sentinel node biopsy and beyond. Clinical and Translational Imaging, 2020, 8, 413-418.	2.1	5
9	Reply to E. Hindié. Journal of Clinical Oncology, 2020, 38, 3238-3240.	1.6	3
10	Timing of sentinel node biopsy independently predicts disease-free and overall survival in clinical stage I-II melanoma patients: A multicentre study of the Italian Melanoma Intergroup (IMI). European Journal of Cancer, 2020, 137, 30-39.	2.8	4
11	Five-Year Analysis of Adjuvant Dabrafenib plus Trametinib in Stage III Melanoma. New England Journal of Medicine, 2020, 383, 1139-1148.	27.0	256
12	1100P Restricted mean survival time (RMST) and cure-rate modeling in estimating survival benefit with adjuvant dabrafenib (D) plus trametinib (T) treatment in melanoma. Annals of Oncology, 2020, 31, S743-S744.	1.2	0
13	Factors Affecting Sentinel Node Metastasis in Thin (T1) Cutaneous Melanomas: Development and External Validation of a Predictive Nomogram. Journal of Clinical Oncology, 2020, 38, 1591-1601.	1.6	50
14	Systemic Therapy for Melanoma: ASCO Guideline. Journal of Clinical Oncology, 2020, 38, 3947-3970.	1.6	190
15	Adjuvant dabrafenib plus trametinib versus placebo in patients with resected, BRAFV600-mutant, stage III melanoma (COMBI-AD): exploratory biomarker analyses from a randomised, phase 3 trial. Lancet Oncology, The, 2020, 21, 358-372.	10.7	94
16	Analysis of Sentinel Node Biopsy and Clinicopathologic Features as Prognostic Factors in Patients With Atypical Melanocytic Tumors. Journal of the National Comprehensive Cancer Network: JNCCN, 2020, 18, 1327-1336.	4.9	3
17	An actionable axis linking NFATc2 to EZH2 controls the EMT-like program of melanoma cells. Oncogene, 2019, 38, 4384-4396.	5.9	36
18	Patient-reported outcomes in patients with resected, high-risk melanoma with BRAFV600E or BRAFV600K mutations treated with adjuvant dabrafenib plus trametinib (COMBI-AD): a randomised, placebo-controlled, phase 3 trial. Lancet Oncology, The, 2019, 20, 701-710.	10.7	50

#	Article	IF	CITATIONS
19	Reply to E. Hindi $ ilde{A}$ $ ilde{C}$ and K.R. Hess. Journal of Clinical Oncology, 2019, 37, 1356-1358.	1.6	1
20	Association between baseline disease characteristics and relapse-free survival (RFS) in patients (pts) with BRAF V600-mutant resected stage III melanoma treated with adjuvant dabrafenib (D) + trametinib (T) or placebo (PBO) Journal of Clinical Oncology, 2019, 37, 9582-9582.	1.6	1
21	Longer Follow-Up Confirms Relapse-Free Survival Benefit With Adjuvant Dabrafenib Plus Trametinib in Patients With Resected <i>BRAF</i> V600–Mutant Stage III Melanoma. Journal of Clinical Oncology, 2018, 36, 3441-3449.	1.6	226
22	Estimate of long-term relapse-free survival (RFS) and analysis of baseline factors associated with RFS in the COMBI-AD trial. Annals of Oncology, 2018, 29, viii445.	1.2	2
23	MACE-A3 immunotherapeutic as adjuvant therapy for patients with resected, MACE-A3-positive, stage III melanoma (DERMA): a double-blind, randomised, placebo-controlled, phase 3 trial. Lancet Oncology, The, 2018, 19, 916-929.	10.7	131
24	Mutational and immune gene expression profiling at relapse in patients (pts) treated with adjuvant dabrafenib plus trametinib (D + T) or placebo (pbo) in the COMBI-AD trial Journal of Clinical Oncology, 2018, 36, 9574-9574.	1.6	1
25	Effect on health-related quality of life (HRQOL) of adjuvant treatment (tx) with dabrafenib plus trametinib (D + T) in patients (pts) with resected stage III <i>BRAF</i> -mutant melanoma Journal of Clinical Oncology, 2018, 36, 9590-9590.	1.6	9
26	Dabrafenib plus trametinib (D + T) as adjuvant treatment of resected <i>BRAF</i> -mutant stage III melanoma: Findings from the COMBI-AD trial analyzed based on AJCC 8 classification Journal of Clinical Oncology, 2018, 36, 9591-9591.	1.6	8
27	The inflammation markers in serum of tumor-bearing rats after plasmonic photothermal therapy. , 2018, , .		0
28	Long-Term Survival after Complete Surgical Resection and Adjuvant Immunotherapy for Distant Melanoma Metastases. Annals of Surgical Oncology, 2017, 24, 3991-4000.	1.5	102
29	Melanoma staging: Evidenceâ€based changes in the American Joint Committee on Cancer eighth edition cancer staging manual. Ca-A Cancer Journal for Clinicians, 2017, 67, 472-492.	329.8	1,662
30	Adjuvant Dabrafenib plus Trametinib in Stage III <i>BRAF</i> -Mutated Melanoma. New England Journal of Medicine, 2017, 377, 1813-1823.	27.0	1,192
31	Sex-specific effect of RNASEL rs486907 and miR-146a rs2910164 polymorphisms' interaction as a susceptibility factor for melanoma skin cancer. Melanoma Research, 2017, 27, 309-314.	1.2	13
32	Prognostic factors in Merkel cell carcinoma patients undergoing sentinel node biopsy. European Journal of Surgical Oncology, 2017, 43, 1536-1541.	1.0	13
33	microRNA Expression in Sentinel Nodes from Progressing Melanoma Patients Identifies Networks Associated with Dysfunctional Immune Response. Genes, 2016, 7, 124.	2.4	8
34	Immunomodulatory Factors Control the Fate of Melanoma Tumor Initiating Cells. Stem Cells, 2016, 34, 2449-2460.	3.2	21
35	NFATc2 is an intrinsic regulator of melanoma dedifferentiation. Oncogene, 2016, 35, 2862-2872.	5.9	43
36	Overcoming melanoma resistance to vemurafenib by targeting CCL2-induced miR-34a, miR-100 and miR-125b. Oncotarget, 2016, 7, 4428-4441.	1.8	84

#	Article	IF	CITATIONS
37	3321 The role of sentinel lymph node biopsy in Merkel Cell Carcinoma: Analysis of 64 patients from a single institution. European Journal of Cancer, 2015, 51, S671.	2.8	0
38	Intralesional administration of L19-IL2/L19-TNF in stage III or stage IVM1a melanoma patients: results of a phase II study. Cancer Immunology, Immunotherapy, 2015, 64, 999-1009.	4.2	138
39	Lymph-Node Ratio in Patients with Cutaneous Melanoma: A Multi-Institution Prognostic Study. Annals of Surgical Oncology, 2015, 22, 2127-2134.	1.5	18
40	Factors predictive of pelvic lymph node involvement and outcomes in melanoma patients with metastatic sentinel lymph node of the groin: A multicentre study. European Journal of Surgical Oncology, 2015, 41, 823-829.	1.0	11
41	Differences in Clinicopathological Features and Distribution of Risk Factors in Italian Melanoma Patients. Dermatology, 2015, 230, 256-262.	2.1	6
42	Three-year follow-up of advanced melanoma patients who received ipilimumab plus fotemustine in the Italian Network for Tumor Biotherapy (NIBIT)-M1 phase II study. Annals of Oncology, 2015, 26, 798-803.	1.2	118
43	"Cancer Bio-Immunotherapy in Sienaâ€; Eleventh Meeting of the Network Italiano per la Bioterapia dei Tumori (NIBIT), Siena, Italy, October 17–19, 2013. Cancer Immunology, Immunotherapy, 2015, 64, 131-135.	4.2	0
44	ITOC2 – 038. Role of exosomes in immune suppression. European Journal of Cancer, 2015, 51, S13.	2.8	3
45	Electrochemotherapy: a good idea in recurrent basal cell carcinoma treatment. Melanoma Management, 2015, 2, 27-31.	0.5	3
46	Armed antibodies for cancer treatment: a promising tool in a changing era. Cancer Immunology, Immunotherapy, 2015, 64, 113-121.	4.2	28
47	The number of excised lymph nodes is associated with survival of melanoma patients with lymph node metastasis. Annals of Oncology, 2014, 25, 240-246.	1.2	34
48	Enrichment of CD56dimKIR+CD57+ highly cytotoxic NK cells in tumour-infiltrated lymph nodes of melanoma patients. Nature Communications, 2014, 5, 5639.	12.8	109
49	Association of promoter polymorphism â^'765 <scp>G</scp> > <scp>C</scp> in the <scp>PTGS</scp> 2 gene with malignant melanoma in <scp>I</scp> talian patients and its correlation to gene expression in dermal fibroblasts. Experimental Dermatology, 2014, 23, 766-768.	2.9	4
50	Number of Excised Lymph Nodes as a Quality Assurance Measure for Lymphadenectomy in Melanoma. JAMA Surgery, 2014, 149, 700.	4.3	42
51	Transcriptional Profiling of Melanoma Sentinel Nodes Identify Patients with Poor Outcome and Reveal an Association of CD30+ T Lymphocytes with Progression. Cancer Research, 2014, 74, 130-140.	0.9	27
52	Accuracy and prognostic value of sentinel lymph node biopsy in head and neck melanomas. Journal of Surgical Research, 2014, 187, 518-524.	1.6	28
53	Alternative Activation of Human Plasmacytoid DCs In Vitro and in Melanoma Lesions: Involvement of LAG-3. Journal of Investigative Dermatology, 2014, 134, 1893-1902.	0.7	74
54	Prediction of Survival in Patients With Thin Melanoma: Results From a Multi-Institution Study. Journal of Clinical Oncology, 2014, 32, 2479-2485.	1.6	103

#	Article	IF	CITATIONS
55	Nonsentinel Lymph Node Status in Patients With Cutaneous Melanoma: Results From a Multi-Institution Prognostic Study. Journal of Clinical Oncology, 2014, 32, 935-941.	1.6	49
56	Enrichment of KIR+CD57+ highly cytotoxic NK cells in sentinel lymph nodes of melanoma patients. Journal of Translational Medicine, 2014, 12, P10.	4.4	0
57	Cutaneous Melanoma in Children and Adolescents: The Italian Rare Tumors in Pediatric Age Project Experience. Journal of Pediatrics, 2014, 164, 376-382.e2.	1.8	47
58	A phase II study of intratumoral application of L19IL2/L19TNF in melanoma patients in clinical stage III or stage IV M1a with presence of injectable cutaneous and/or subcutaneous lesions Journal of Clinical Oncology, 2014, 32, TPS9103-TPS9103.	1.6	4
59	Isolated limb perfusion with the tumorâ€ŧargeting human monoclonal antibody–cytokine fusion protein L19â€₹NF plus melphalan and mild hyperthermia in patients with locally advanced extremity melanoma. Journal of Surgical Oncology, 2013, 107, 173-179.	1.7	72
60	Analysis of surrogate gene expression markers in peripheral blood of melanoma patients to predict treatment outcome of adjuvant pegylated interferon alpha 2b (EORTC 18991 side study). Cancer Immunology, Immunotherapy, 2013, 62, 1223-1233.	4.2	5
61	Effects of cyclophosphamide and IL-2 on regulatory CD4+ T cell frequency and function in melanoma patients vaccinated with HLA-class I peptides: impact on the antigen-specific T cell response. Cancer Immunology, Immunotherapy, 2013, 62, 897-908.	4.2	31
62	The use of polytetrafluoroethylene to facilitate the vascular access in recurrent melanoma to limbs. International Journal of Surgery Case Reports, 2013, 4, 40-43.	0.6	1
63	A variant in FTO shows association with melanoma risk not due to BMI. Nature Genetics, 2013, 45, 428-432.	21.4	111
64	Biological insights into BRAF ^{V600} mutations in melanoma patient. Oncolmmunology, 2013, 2, e25594.	4.6	6
65	Clinical and immunologic responses in melanoma patients vaccinated with MAGEâ€A3â€genetically modified lymphocytes. International Journal of Cancer, 2013, 132, 2557-2566.	5.1	20
66	<scp>CDKN</scp> 2A and <scp>MC</scp> 1R variants influence dermoscopic and confocal features of benign melanocytic lesions in multiple melanoma patients. Experimental Dermatology, 2013, 22, 411-416.	2.9	26
67	Adjuvant Ganglioside GM2-KLH/QS-21 Vaccination Versus Observation After Resection of Primary Tumor > 1.5 mm in Patients With Stage II Melanoma: Results of the EORTC 18961 Randomized Phase III Trial. Journal of Clinical Oncology, 2013, 31, 3831-3837.	1.6	88
68	Long-Term Results of the Randomized Phase III Trial EORTC 18991 of Adjuvant Therapy With Pegylated Interferon Alfa-2b Versus Observation in Resected Stage III Melanoma. Journal of Clinical Oncology, 2012, 30, 3810-3818.	1.6	254
69	Modulation of Microenvironment Acidity Reverses Anergy in Human and Murine Tumor-Infiltrating T Lymphocytes. Cancer Research, 2012, 72, 2746-2756.	0.9	470
70	Limited Induction of Tumor Cross-Reactive T Cells without a Measurable Clinical Benefit in Early Melanoma Patients Vaccinated with Human Leukocyte Antigen Class I–Modified Peptides. Clinical Cancer Research, 2012, 18, 6485-6496.	7.0	61
71	Ulceration and stage are predictive of interferon efficacy in melanoma: Results of the phase III adjuvant trials EORTC 18952 and EORTC 18991. European Journal of Cancer, 2012, 48, 218-225.	2.8	182
72	lpilimumab and fotemustine in patients with advanced melanoma (NIBIT-M1): an open-label, single-arm phase 2 trial. Lancet Oncology, The, 2012, 13, 879-886.	10.7	273

#	Article	IF	CITATIONS
73	Small Nodular Melanoma: The Beginning of a Life-Threatening Lesion. A Clinical Study on 11 Cases. Tumori, 2011, 97, 35-38.	1.1	9
74	EORTC 18991 phase III trial: Long-term adjuvant pegylated interferon-α2b (PEG-IFN) versus observation in resected stage III melanoma: Long-term results at 7.6-years follow-up Journal of Clinical Oncology, 2011, 29, 8506b-8506b.	1.6	13
75	A phase II study combining ipilimumab and fotemustine in patients with metastatic melanoma: The NIBIT-M1 trial Journal of Clinical Oncology, 2011, 29, TPS230-TPS230.	1.6	5
76	Pure Desmoplastic Melanoma. Annals of Surgery, 2010, 252, 1052-1057.	4.2	49
77	pHâ€dependent antitumor activity of proton pump inhibitors against human melanoma is mediated by inhibition of tumor acidity. International Journal of Cancer, 2010, 127, 207-219.	5.1	237
78	Phase III Trial Comparing Adjuvant Treatment With Pegylated Interferon Alfa-2b Versus Observation: Prognostic Significance of Autoantibodies—EORTC 18991. Journal of Clinical Oncology, 2010, 28, 2460-2466.	1.6	69
79	Tumor-Reactive CD8+ Early Effector T Cells Identified at Tumor Site in Primary and Metastatic Melanoma. Cancer Research, 2010, 70, 8378-8387.	0.9	52
80	Heterogeneous Phenotype of Human Melanoma Cells with In Vitro and In Vivo Features of Tumor-Initiating Cells. Journal of Investigative Dermatology, 2010, 130, 1877-1886.	0.7	77
81	Response to Griewank and Bastian. Journal of Investigative Dermatology, 2010, 130, 2331-2332.	0.7	0
82	High Levels of Exosomes Expressing CD63 and Caveolin-1 in Plasma of Melanoma Patients. PLoS ONE, 2009, 4, e5219.	2.5	806
83	Adjuvant Therapy With Pegylated Interferon Alfa-2b Versus Observation in Resected Stage III Melanoma: A Phase III Randomized Controlled Trial of Health-Related Quality of Life and Symptoms by the European Organisation for Research and Treatment of Cancer Melanoma Group. Journal of Clinical Oncology, 2009, 27, 2916-2923.	1.6	119
84	Impaired STAT Phosphorylation in T Cells from Melanoma Patients in Response to IL-2: Association with Clinical Stage. Clinical Cancer Research, 2009, 15, 4085-4094.	7.0	29
85	Modified peptides in anti-cancer vaccines: are we eventually improving anti-tumour immunity?. Cancer Immunology, Immunotherapy, 2009, 58, 1159-1167.	4.2	21
86	Cutaneous Melanoma in Childhood and Adolescence Shows Frequent Loss of INK4A and Gain of KIT. Journal of Investigative Dermatology, 2009, 129, 1759-1768.	0.7	54
87	New common variants affecting susceptibility to basal cell carcinoma. Nature Genetics, 2009, 41, 909-914.	21.4	303
88	Single-Institution Series of Early-Stage Merkel Cell Carcinoma: Long-Term Outcomes in 95 Patients Managed with Surgery Alone. Annals of Surgical Oncology, 2009, 16, 2985-2993.	1.5	50
89	Circulating melanoma cells and distant metastasis-free survival in stage III melanoma patients with or without adjuvant interferon treatment (EORTC 18991 side study). European Journal of Cancer, 2009, 45, 3189-3197.	2.8	48
90	Radical dissection after positive groin sentinel biopsy in melanoma patients: rate of further positive nodes. Melanoma Research, 2009, 19, 112-118.	1.2	24

#	Article	IF	CITATIONS
91	Banked venous homograft replacement of the inferior vena cava for primary leiomyosarcoma. European Journal of Surgical Oncology, 2008, 34, 720-724.	1.0	18
92	Adjuvant therapy with pegylated interferon alfa-2b versus observation alone in resected stage III melanoma: final results of EORTC 18991, a randomised phase III trial. Lancet, The, 2008, 372, 117-126.	13.7	620
93	Metamorphosis of melanoma. Trends in size and thickness of cutaneous melanoma over one decade at the Istituto Nazionale Tumori, Milan. Tumori, 2008, 94, 11-13.	1.1	22
94	Adjuvant multipeptide vaccination in high-risk early melanoma patients. Journal of Clinical Oncology, 2008, 26, 3014-3014.	1.6	3
95	Opposite immune functions of GM-CSF administered as vaccine adjuvant in cancer patients. Annals of Oncology, 2007, 18, 226-232.	1.2	252
96	Melanoma immunology: past, present and future. Current Opinion in Oncology, 2007, 19, 121-127.	2.4	57
97	INV 4 Impaired response to ??c cytokines in T cells from melanoma patients. Melanoma Research, 2007, 17, A2.	1.2	0
98	Melanoma contains CD133 and ABCG2 positive cells with enhanced tumourigenic potential. European Journal of Cancer, 2007, 43, 935-946.	2.8	523
99	Multispectral imaging and artificial neural network: mimicking the management decision of the clinician facing pigmented skin lesions. Physics in Medicine and Biology, 2007, 52, 2599-2613.	3.0	58
100	Detection of mutated BRAFV600E variant in circulating DNA of stage Ill–IV melanoma patients. International Journal of Cancer, 2007, 120, 2439-2444.	5.1	76
101	Advanced Extremity Soft Tissue Sarcoma: Prognostic Effect of Isolated Limb Perfusion in a Series of 88 Patients Treated at a Single Institution. Annals of Surgical Oncology, 2007, 14, 553-559.	1.5	61
102	EORTC 18991: Long-term adjuvant pegylated interferon-alpha2b (PEG-IFN) compared to observation in resected stage III melanoma, final results of a randomized phase III trial. Journal of Clinical Oncology, 2007, 25, 8504-8504.	1.6	18
103	CD133 POSTIVE CELLULAR POPULATION IN HUMAN MELANOMA. FASEB Journal, 2007, 21, A32.	0.5	0
104	Identification of a new subset of myeloid suppressor cells in peripheral blood of melanoma patients and modulation by GM-CSF-based anti-tumor vaccine. Journal of Clinical Oncology, 2007, 25, 21082-21082.	1.6	1
105	Micro-melanoma detection: a clinical study on 206 consecutive cases of pigmented skin lesions with a diameter â‰≇€ƒ3 mm. British Journal of Dermatology, 2006, 155, 570-573.	1.5	66
106	A phase II trial of vaccination with autologous, tumor-derived heat-shock protein peptide complexes Gp96, in combination with GM-CSF and interferon-α in metastatic melanoma patients. Cancer Immunology, Immunotherapy, 2006, 55, 958-968.	4.2	134
107	Sentinel and Nonsentinel Node Status in Stage IB and II Melanoma Patients: Two-Step Prognostic Indicators of Survival. Journal of Clinical Oncology, 2006, 24, 4464-4471.	1.6	132
108	Vaccination: role in metastatic melanoma. Expert Review of Anticancer Therapy, 2006, 6, 1305-1318.	2.4	22

#	Article	IF	CITATIONS
109	Immunization of Stage IV Melanoma Patients with Melan-A/MART-1 and gp100 Peptides plus IFN-α Results in the Activation of Specific CD8+ T Cells and Monocyte/Dendritic Cell Precursors. Cancer Research, 2006, 66, 4943-4951.	0.9	108
110	Hyperthermic isolation limb perfusion with TNFalpha in the treatment of in-transit melanoma metastasis. In Vivo, 2006, 20, 739-42.	1.3	6
111	Evaluation of Myeloid Suppressive Cells in Peripheral Blood of Melanoma Patients and Their Modulation by A Heat-shock Protein (HSP)-96 and GM-CSF-based Vaccine. Journal of Immunotherapy, 2005, 28, 659.	2.4	0
112	Does Melanoma Behave Differently in Younger Children Than in Adults? A Retrospective Study of 33 Cases of Childhood Melanoma From a Single Institution. Pediatrics, 2005, 115, 649-654.	2.1	215
113	Automated segmentation of pigmented skin lesions in multispectral imaging. Physics in Medicine and Biology, 2005, 50, N345-N357.	3.0	17
114	Constitutive Expression and Costimulatory Function of LIGHT/TNFSF14 on Human Melanoma Cells and Melanoma-Derived Microvesicles. Cancer Research, 2005, 65, 3428-3436.	0.9	53
115	Narrower Surgical Margins Might be Sufficient in Invasive Horizontal Growth Phase Melanoma. Tumori, 2004, 90, 464-466.	1.1	0
116	Heat Shock Proteins and Their Use as Anticancer Vaccines. Clinical Cancer Research, 2004, 10, 8142-8146.	7.0	62
117	Atypical pleomorphic epithelioid angiomyolipoma localized to the pelvis: a case report and review of the literature. Histopathology, 2004, 44, 292-295.	2.9	17
118	Retroperitoneal soft tissue sarcomas. Cancer, 2004, 100, 2448-2455.	4.1	167
119	Immunotherapy of melanoma. Seminars in Cancer Biology, 2003, 13, 391-400.	9.6	48
120	World Health Organization experience in the treatment of melanoma. Surgical Clinics of North America, 2003, 83, 405-416.	1.5	17
121	Chordoma: Natural History and Results in 28 Patients Treated at a Single Institution. Annals of Surgical Oncology, 2003, 10, 291-296.	1.5	204
122	Quality of Surgery and Outcome in Extra-Abdominal Aggressive Fibromatosis: A Series of Patients Surgically Treated at a Single Institution. Journal of Clinical Oncology, 2003, 21, 1390-1397.	1.6	326
123	Hypoxic pelvic and limb perfusion with melphalan and mitomycin C for recurrent limb melanoma. Melanoma Research, 2003, 13, 51-58.	1.2	21
124	E-cadherin Expression on Fine Needle Aspiration Biopsies of Breast Invasive Ductal Carcinomas and Its Relationship to Clinicopathologic Factors. Acta Cytologica, 2003, 47, 363-367.	1.3	8
125	Surgical management of primary melanoma. , 2003, , 247-254.		0
126	Lack of terminally differentiated tumor-specific CD8+ T cells at tumor site in spite of antitumor immunity to self-antigens in human metastatic melanoma. Cancer Research, 2003, 63, 2535-45.	0.9	142

#	Article	IF	CITATIONS
127	Deliberate hypoxic pelvic and limb chemoperfusion in the treatment of recurrent melanoma. American Journal of Surgery, 2002, 183, 28-36.	1.8	25
128	DNA fragmentation and cell proliferation correlated with tumor grade in patients with hepatocellular carcinoma. Cancer, 2002, 96, 301-305.	4.1	9
129	Clinical and dermatoscopic diagnosis of small pigmented skin lesions. European Journal of Dermatology, 2002, 12, 573-6.	0.6	12
130	Effect of long-term adjuvant therapy with interferon alpha-2a in patients with regional node metastases from cutaneous melanoma: a randomised trial. Lancet, The, 2001, 358, 866-869.	13.7	248
131	Impact of Clinical Trials on the Treatment of Melanoma. Surgical Oncology Clinics of North America, 2001, 10, 935-947.	1.5	13
132	Isolated Limb Perfusion. Surgical Oncology Clinics of North America, 2001, 10, 821-832.	1.5	38
133	Sentinel Lymph Node Biopsy in Cutaneous Melanoma: The WHO Melanoma Program Experience. Annals of Surgical Oncology, 2000, 7, 469-474.	1.5	318
134	Sentinel Node Biopsy and Selective Lymph Node Dissection in Cutaneous Melanoma Patients. , 2000, , 235-242.		9
135	Hyperthermic Antiblastic Perfusion with Alpha Tumor Necrosis Factor and Doxorubicin for the Treatment of Soft Tissue Limb Sarcoma in Candidates for Amputation. Journal of Immunotherapy, 1999, 22, 407-414.	2.4	24
136	Dissection of regional lymph nodes in cutaneous melanoma. Lancet, The, 1998, 351, 1885.	13.7	0
137	Immediate or delayed dissection of regional nodes in patients with melanoma of the trunk: a randomised trial. Lancet, The, 1998, 351, 793-796.	13.7	625
138	Treatment of Recurrent Sarcoma of the Extremities by Isolated Limb Perfusion Using Tumor Necrosis Factor Alpha and Melphalan. Tumori, 1996, 82, 579-584.	1.1	25
139	Isolated Pelvic Perfusion for the Treatment of Unresectable Primary or Recurrent Rectal Cancer. Tumori, 1996, 82, 459-462.	1.1	8
140	Intraperitoneal hyperthermic perfusion (IPHP). Oncology Reports, 1996, 3, 1103-6.	2.6	5
141	Current results of pelvic perfusion for non-resectable relapsing of pelvic cancer. Oncology Reports, 1996, 3, 1097-1102.	2.6	0
142	Isolation perfusion in extracorporeal circulation with interleukin-2 and lymphokine-activated killer cells in the treatment of in-transit metastases from limb cutaneous melanoma. Annals of Surgical Oncology, 1995, 2, 61-70.	1.5	14
143	Cardiac and Pulmonary Effects of Alpha Tumor Necrosis Factor Administered by Isolation Perfusion. Tumori, 1995, 81, 197-200.	1.1	2
144	Treatment of primary or relapsing limb cancer by isolation perfusion with high-dose alpha-tumor necrosis factor, gamma-interferon, and melphalan. Cancer, 1994, 73, 483-492.	4.1	75

#	Article	IF	CITATIONS
145	Hyperthermic antiblastic perfusion in the treatment of stage IIIA-IIIAB melanoma patients. Comparison of two experiences. Melanoma Research, 1994, 4, 269.	1.2	3
146	Excision of primary melanoma. Melanoma Research, 1993, 3, 13.	1.2	4
147	The role of tamoxifen (T) in association with chemotherapy in metastatic melanoma (MM). Melanoma Research, 1993, 3, 53.	1.2	Ο
148	Adoptive Immunotherapy of Melanoma with Interleukin-2 and Lymphocytes. , 1993, , 243-252.		0
149	Treatment of recurrent in transit metastases from cutaneous melanoma by isolation perfusion in extracorporeal circulation with interleukin-2 and lymphokine activated killer cells. A pilot study. Melanoma Research, 1992, 2, 263-272.	1.2	10
150	Elective lymph node dissection for melanoma: Two perspectives. World Journal of Surgery, 1992, 16, 203-213.	1.6	40
151	Antitumor activity of hyperthermia alone or in combination with cisplatin and melphalan in primary cultures of human malignant melanoma. International Journal of Cell Cloning, 1989, 7, 385-394.	1.6	17
152	Management of nodal metastases from head and neck melanoma. Journal of Surgical Oncology, 1989, 42, 47-53.	1.7	7
153	Seven years experience with hyperthermic perfusions in extracorporeal circulation for melanoma of the extremities. Journal of Surgical Oncology, 1989, 42, 201-208.	1.7	47
154	Clinical status of diagnosis and therapy of malignant melanoma. International Journal of Radiation Applications and Instrumentation Part B, Nuclear Medicine and Biology, 1989, 16, 621-624.	0.3	1
155	Isolation perfusion of the lower limb with platinum. World Journal of Surgery, 1988, 12, 307-309.	1.6	10
156	HLA Antigens in Familial and Sporadic Cutaneous Melanoma. Tumori, 1988, 74, 657-664.	1.1	2
157	Intra-arterial infusion and perfusion chemotherapy for soft tissue sarcomas of the extremities. Cancer Treatment and Research, 1986, , 103-129.	0.5	5
158	Limb-Salvage Procedure in Two Patients with Spindle Cell Carcinoma of the Lower Extremities using Isolation Perfusion. Tumori, 1985, 71, 271-275.	1.1	4
159	Hyperthermic Antiblastic Perfusion in the Treatment of Cancer of the Extremities. Tumori, 1985, 71, 355-359.	1.1	25
160	Prognosis of skin melanoma with regional node metastases (stage II). Journal of Surgical Oncology, 1984, 25, 240-247.	1.7	83
161	Regional Perfusion at High Temperature in Treatment of Stage IIIA-IIIAB Melanoma Patients. Tumori, 1983, 69, 585-588.	1.1	20
162	The advent of minimally invasive lymphadenectomy in melanoma. Annals of Laparoscopic and Endoscopic Surgery, 0, 2, 122-122.	0.5	1

#	Article	IF	CITATIONS
163	Cutaneous Melanoma â^' Surgical Treatment. , 0, , .		0