
Stefanie Kreutmair

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1660963/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Distinct immunological signatures discriminate severe COVID-19 from non-SARS-CoV-2-driven critical pneumonia. Immunity, 2021, 54, 1578-1593.e5.	14.3	75
2	Phosphorylation of BECLIN-1 by BCR-ABL suppresses autophagy in chronic myeloid leukemia. Haematologica, 2020, 105, 1285-1293.	3.5	22
3	Identification of the Lymphoma-Initiating Cell Population in a T Cell Lymphoma Mouse Model Resembling Human Anaplastic Large Cell Lymphoma. Blood, 2014, 124, 2213-2213.	1.4	16
4	NPM1c alters FLT3-D835Y localization and signaling in acute myeloid leukemia. Blood, 2019, 134, 383-388.	1.4	14
5	The IL-3, IL-5, and GM-CSF common receptor beta chain mediates oncogenic activity of FLT3-ITD-positive AML. Leukemia, 2021, , .	7.2	12
6	Extracellular Signal-regulated Kinase 2 (ERK2) Mediates Phosphorylation and Inactivation of Nuclear Interaction Partner of Anaplastic Lymphoma Kinase (NIPA) at G2/M. Journal of Biological Chemistry, 2012, 287, 37997-38005.	3.4	10
7	Inhibition of Aurora Kinase B Is Important for Biologic Activity of the Dual Inhibitors of BCR-ABL and Aurora Kinases R763/AS703569 and PHA-739358 in BCR-ABL Transformed Cells. PLoS ONE, 2014, 9, e112318.	2.5	9
8	Loss of the Fanconi anemia–associated protein NIPA causes bone marrow failure. Journal of Clinical Investigation, 2020, 130, 2827-2844.	8.2	8
9	First-in-human study of WT1 recombinant protein vaccination in elderly patients with AML in remission: a single-center experience. Cancer Immunology, Immunotherapy, 2022, 71, 2913-2928.	4.2	8
10	Preexisting comorbidities shape the immune response associated with severe COVID-19. Journal of Allergy and Clinical Immunology, 2022, 150, 312-324.	2.9	8
11	Proteomic Phosphosite Analysis Identified Crucial NPM-ALK-Mediated NIPA Serine and Threonine Residues. International Journal of Molecular Sciences, 2019, 20, 4060.	4.1	5
12	APC/CCdh1 regulates the balance between maintenance and differentiation of hematopoietic stem and progenitor cells. Cellular and Molecular Life Sciences, 2019, 76, 369-380.	5.4	5
13	Existence of reprogrammed lymphoma stem cells in a murine ALCL-like model. Leukemia, 2020, 34, 3242-3255.	7.2	4
14	First-in-Human Study of WT1 Recombinant Protein Vaccination in Elderly Patients with AML in Remission: A Single-Center Experience. Blood, 2021, 138, 1278-1278.	1.4	4
15	A novel conditional NPM-ALK-driven model of CD30+ T-cell lymphoma mediated by a translational stop cassette. Oncogene, 2020, 39, 1904-1913.	5.9	3
16	NIPA (Nuclear Interaction Partner of ALK) Is Crucial for Effective NPM-ALK Mediated Lymphomagenesis. Frontiers in Oncology, 2022, 12, .	2.8	1
17	The F-Box Protein NIPA Limits Hematopoietic Stem Cell Survival and Transplantation Efficiency. Blood, 2013, 122, 1175-1175.	1.4	Ο
18	Towards Improved Models of NPM-ALK Induced T-Cell Lymphoma. Blood, 2018, 132, 1342-1342.	1.4	0

#	Article	IF	CITATIONS
19	Checkpoint Inhibition in CSF3R Mutated Chronic Neutrophilic Leukemia. Blood, 2018, 132, 3056-3056.	1.4	Ο
20	Examining the Role of CD30 in an Anaplastic Large Cell Lymphoma Mouse Model. Blood, 2019, 134, 2542-2542.	1.4	0
21	The Fanconi Anemia-Associated Protein NIPA Is Essential for the Nuclear Abundance of FANCD2. Blood, 2019, 134, 3741-3741.	1.4	Ο
22	Response to Are NKT cells a useful predictor of COVID-19 severity?. Immunity, 2022, 55, 188-189.	14.3	0