
Sean A Martin

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1667109/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	The association between obstructive sleep apnea and sleep spindles in middle-aged and older men: a community-based cohort study. Sleep, 2022, 45, .	0.6	11
2	The bidirectional association between depression and lower urinary tract symptoms (LUTS) in men: A systematic review and metaâ€analysis of observational studies. Neurourology and Urodynamics, 2022, 41, 552-561.	0.8	5
3	The association between sleep microarchitecture and cognitive function in middle-aged and older men: a community-based cohort study. Journal of Clinical Sleep Medicine, 2022, 18, 1593-1608.	1.4	6
4	Participation in physical activity is associated with reduced nocturnal hypoxaemia in males. ERJ Open Research, 2021, 7, 00852-2020.	1.1	1
5	Effect of depression on health service utilisation in men: a prospective cohort study of Australian men aged 35 to 80 years. BMJ Open, 2021, 11, e044893.	0.8	8
6	Sleep macroarchitecture but not obstructive sleep apnea is independently associated with cognitive function in only older men of a populationâ€based cohort. Journal of Sleep Research, 2021, 30, e13370.	1.7	11
7	Analysis of major fatty acids from matched plasma and serum samples reveals highly comparable absolute and relative levels. Prostaglandins Leukotrienes and Essential Fatty Acids, 2021, 168, 102268.	1.0	13
8	The Association of Obstructive Sleep Apnea and Nocturnal Hypoxemia with Lipid Profiles in a Population-Based Study of Community-Dwelling Australian Men. Nature and Science of Sleep, 2021, Volume 13, 1771-1782.	1.4	12
9	<associations and="" body="" community<br="" composition="" hypoxemia="" in="" nocturnal="" of="" osa="" strength="" with="">Dwelling Middle Aged and Older Men. Nature and Science of Sleep, 2020, Volume 12, 959-968.</associations>	1.4	4
10	Obstructive sleep apnea is not an independent determinant of testosterone in men. European Journal of Endocrinology, 2020, 183, 31-39.	1.9	13
11	Measuring Masculinity in Men With Chronic Disease. American Journal of Men's Health, 2019, 13, 155798831985970.	0.7	8
12	Quantitative electroencephalography measures in rapid eye movement and nonrapid eye movement sleep are associated with apnea–hypopnea index and nocturnal hypoxemia in men. Sleep, 2019, 42, .	0.6	36
13	Re: Rhee et al. "Longitudinal study of the relationship between lower urinary tract symptoms and depressive symptomsâ€. Journal of Psychosomatic Research, 2019, 116, 113-114.	1.2	0
14	Higher Serum Sex Hormone–Binding Globulin Levels Are Associated With Incident Cardiovascular Disease in Men. Journal of Clinical Endocrinology and Metabolism, 2019, 104, 6301-6315.	1.8	31
15	Association of endogenous testosterone concentration with depression in men. JBI Database of Systematic Reviews and Implementation Reports, 2019, 17, 1894-1900.	1.7	1
16	Suitability of the Epworth Sleepiness Scale (ESS) for Economic Evaluation: An Assessment of Its Convergent and Discriminant Validity. Behavioral Sleep Medicine, 2018, 16, 448-470.	1.1	5
17	<scp>International Prostate Symptom Score</scp> Should Be Considered a Complement Rather Than a Substitute to Generic Preferenceâ€Based Measures for Measuring <scp>Lower Urinary Tract Symptoms</scp> Within Economic Evaluation. LUTS: Lower Urinary Tract Symptoms, 2018, 10, 45-56.	0.6	1
18	The role of sex hormone-binding globulin (SHBG), testosterone, and other sex steroids, on the development of type 2 diabetes in a cohort of community-dwelling middle-aged to elderly men. Acta Diabetologica, 2018, 55, 861-872.	1.2	42

SEAN A MARTIN

#	Article	IF	CITATIONS
19	Position statement: a clinical approach to the management of adult nonâ€neurogenic overactive bladder. Medical Journal of Australia, 2018, 208, 461-462.	0.8	1
20	Age-related changes in estradiol and longitudinal associations with fat mass in men. PLoS ONE, 2018, 13, e0201912.	1.1	12
21	Cross-sectional and longitudinal determinants of serum sex hormone binding globulin (SHBG) in a cohort of community-dwelling men. PLoS ONE, 2018, 13, e0200078.	1.1	21
22	Chronic Kidney Disease and Sleep Apnea Association of Kidney Disease With Obstructive Sleep Apnea in a Population Study of Men. Sleep, 2017, 40, .	0.6	26
23	Coâ€morbid <scp>OSA</scp> and insomnia increases depression prevalence and severity in men. Respirology, 2017, 22, 1407-1415.	1.3	67
24	Erectile dysfunction is independently associated with apnea-hypopnea index and oxygen desaturation index in elderly, but not younger, community-dwelling men. Sleep Health, 2017, 3, 250-256.	1.3	12
25	The association between total phthalate concentration and non-communicable diseases and chronic inflammation in South Australian urban dwelling men. Environmental Research, 2017, 158, 366-372.	3.7	35
26	The association between gastroesophageal reflux disease with sleep quality, depression, and anxiety in a cohort study of Australian men. Journal of Gastroenterology and Hepatology (Australia), 2017, 32, 1170-1177.	1.4	33
27	Associations of Undiagnosed Obstructive Sleep Apnea and Excessive Daytime Sleepiness With Depression: An Australian Population Study. Journal of Clinical Sleep Medicine, 2017, 13, 575-582.	1.4	33
28	Long-Term Effects of a Randomised Controlled Trial Comparing High Protein or High Carbohydrate Weight Loss Diets on Testosterone, SHBG, Erectile and Urinary Function in Overweight and Obese Men. PLoS ONE, 2016, 11, e0161297.	1.1	60
29	Association of daytime sleepiness with obstructive sleep apnoea and comorbidities varies by sleepiness definition in a population cohort of men. Respirology, 2016, 21, 1314-1321.	1.3	34
30	Elucidating the Biological Mechanisms Linking Depressive Symptoms With Type 2 Diabetes in Men. Psychosomatic Medicine, 2016, 78, 221-232.	1.3	8
31	Nocturia, Other Lower Urinary Tract Symptoms and Sleep Dysfunction in a Community-Dwelling Cohort of Men. Urology, 2016, 97, 219-226.	0.5	24
32	Predictive value of serum testosterone for type 2 diabetes risk assessment in men. BMC Endocrine Disorders, 2016, 16, 26.	0.9	31
33	The association of obstructive sleep apnea (OSA) and nocturnal hypoxemia with the development of abnormal HbA1c in a population cohort of men without diabetes. Diabetes Research and Clinical Practice, 2016, 114, 151-159.	1.1	16
34	Hypertension Is Associated With Undiagnosed OSA During Rapid Eye Movement Sleep. Chest, 2016, 150, 495-505.	0.4	96
35	Food Habits, Lifestyle Factors and Mortality among Oldest Old Chinese: The Chinese Longitudinal Healthy Longevity Survey (CLHLS). Nutrients, 2015, 7, 7562-7579.	1.7	68
36	The Association of Socio-Demographic Status, Lifestyle Factors and Dietary Patterns with Total Urinary Phthalates in Australian Men. PLoS ONE, 2015, 10, e0122140.	1.1	26

SEAN A MARTIN

#	Article	lF	CITATIONS
37	Lower Urinary Tract Symptoms, Depression, Anxiety and Systemic Inflammatory Factors in Men: A Population-Based Cohort Study. PLoS ONE, 2015, 10, e0137903.	1.1	43
38	Nocturnal Hypoxemia and Severe Obstructive Sleep Apnea are Associated with Incident Type 2 Diabetes in a Population Cohort of Men. Journal of Clinical Sleep Medicine, 2015, 11, 609-614.	1.4	47
39	The longitudinal association between inflammation and incident depressive symptoms in men: The effects of hs-CRP are independent of abdominal obesity and metabolic disturbances. Physiology and Behavior, 2015, 139, 328-335.	1.0	16
40	Undiagnosed obstructive sleep apnea is independently associated with reductions in quality of life in middle-aged, but not elderly men of a population cohort. Sleep and Breathing, 2015, 19, 1309-1316.	0.9	57
41	Predictors of Sexual Dysfunction Incidence and Remission in Men. Journal of Sexual Medicine, 2014, 11, 1136-1147.	0.3	79
42	Cohort Profile: The Men Androgen Inflammation Lifestyle Environment and Stress (MAILES) Study. International Journal of Epidemiology, 2014, 43, 1040-1053.	0.9	53
43	Testosterone is associated with self-employment among Australian men. Economics and Human Biology, 2014, 13, 76-84.	0.7	27
44	Risk Factors for Progression or Improvement of Lower Urinary Tract Symptoms in a Prospective Cohort of Men. Journal of Urology, 2014, 191, 130-137.	0.2	76
45	Clinical and Biopsychosocial Determinants of Sexual Dysfunction in Middleâ€Aged and Older Australian Men. Journal of Sexual Medicine, 2012, 9, 2093-2103.	0.3	26
46	Overactive bladder in men as a marker of cardiometabolic risk. Medical Journal of Australia, 2012, 197, 379-380.	0.8	1
47	Prevalence and factors associated with uncomplicated storage and voiding lower urinary tract symptoms in community-dwelling Australian men. World Journal of Urology, 2011, 29, 179-184.	1.2	116
48	Serum testosterone bioassay evaluation in a large male cohort. Clinical Endocrinology, 2010, 72, 87-98.	1.2	5
49	Demographic, physical and lifestyle factors associated with androgen status: the Florey Adelaide Male Ageing Study (FAMAS). Clinical Endocrinology, 2009, 71, 261-272.	1.2	41
50	Chronic disease prevalence and associations in a cohort of Australian men: The Florey Adelaide Male Ageing Study (FAMAS). BMC Public Health, 2008, 8, 261.	1.2	13
51	Cohort Profile: The Florey Adelaide Male Ageing Study (FAMAS). International Journal of Epidemiology, 2007, 36, 302-306.	0.9	39
52	The Florey Adelaide Male Ageing Study (FAMAS): Design, procedures & participants. BMC Public Health, 2007, 7, 126.	1.2	40