

Felix Kh Chun

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1670696/felix-kh-chun-publications-by-year.pdf>

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

173 papers	7,868 citations	54 h-index	80 g-index
191 ext. papers	8,894 ext. citations	5.4 avg, IF	5.02 L-index

#	Paper	IF	Citations
173	Super early detailed assessment of lower urinary tract symptoms after holmium laser enucleation of the prostate (HoLEP): a prospective study. <i>World Journal of Urology</i> , 2020 , 38, 3207-3217	4	1
172	Performance and Impact of Prostate Specific Membrane Antigen-Based Diagnostics in the Management of Men with Biochemical Recurrence of Prostate Cancer and its Role in Salvage Lymph Node Dissection. <i>World Journal of Men's Health</i> , 2020 , 38, 32-47	6.8	1
171	Prediction of Complications in Radical Prostatectomy Prostate Cancer Patients: Simulated Annealing versus Co-Morbidity Indexes. <i>Urologia Internationalis</i> , 2019 , 102, 51-59	1.9	2
170	Adherence to pelvic lymph node dissection recommendations according to the National Comprehensive Cancer Network pelvic lymph node dissection guideline and the D'Amico lymph node invasion risk stratification. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2018 , 36, 81.e17-81.e24	2.8	10
169	North American Population-Based Validation of the National Comprehensive Cancer Network Practice Guideline Recommendation of Pelvic Lymphadenectomy in Contemporary Prostate Cancer. <i>Prostate</i> , 2017 , 77, 542-548	4.2	13
168	Local Therapy Improves Survival in Metastatic Prostate Cancer. <i>European Urology</i> , 2017 , 72, 118-124	10.2	77
167	Population-Based External Validation of the Updated 2012 Partin Tables in Contemporary North American Prostate Cancer Patients. <i>Prostate</i> , 2017 , 77, 105-113	4.2	20
166	Wound dehiscence in a sample of 1 776 cystectomies: identification of predictors and implications for outcomes. <i>BJU International</i> , 2016 , 117, E95-E101	5.6	21
165	The impact of resident involvement in male one-stage anterior urethroplasties. <i>Urology</i> , 2015 , 85, 937-41	1.6	17
164	The Effect of Resident Involvement on Perioperative Outcomes in Transurethral Urologic Surgeries. <i>Journal of Surgical Education</i> , 2015 , 72, 1018-25	3.4	27
163	Prognostic impact of infiltration of the vagina and/or uterus in women undergoing anterior pelvic exenteration for urothelial carcinoma of the bladder: results of a contemporary multicentre series. <i>World Journal of Urology</i> , 2015 , 33, 343-50	4	4
162	Thermal Ablation of Renal Tumors: Indications, Techniques and Results. <i>Deutsches A&#x0308;rztblatt International</i> , 2015 , 112, 412-8	2.5	13
161	Natural history of surgically treated high-risk prostate cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015 , 33, 163.e7-13	2.8	77
160	Evidence from the 'PROspective MulticEnTer Radical Cystectomy Series 2011 (PROMETRICS 2011)' study: how are preoperative patient characteristics associated with urinary diversion type after radical cystectomy for bladder cancer?. <i>Annals of Surgical Oncology</i> , 2015 , 22, 1032-42	3.1	26
159	In Reply. <i>Deutsches A&#x0308;rztblatt International</i> , 2015 , 112, 758	2.5	
158	Prediction of metastatic status in non-seminomatous testicular cancer. <i>World Journal of Urology</i> , 2014 , 32, 1205-11	4	5
157	Holmium laser enucleation of the prostate is safe in patients with prostate cancer and lower urinary tract symptoms--a retrospective feasibility study. <i>Journal of Endourology</i> , 2014 , 28, 335-41	2.7	10

156	A comparative assessment of active surveillance for localized prostate cancer in the community versus tertiary care referral center. <i>World Journal of Urology</i> , 2014 , 32, 891-7	4	7
155	Concomitant seminal vesicle invasion in pT4a urothelial carcinoma of the bladder with contiguous prostatic infiltration is an adverse prognosticator for cancer-specific survival after radical cystectomy. <i>Annals of Surgical Oncology</i> , 2014 , 21, 4034-40	3.1	12
154	Reduced membranous MET expression is linked to bladder cancer progression. <i>Cancer Genetics</i> , 2014 , 207, 147-52	2.3	4
153	Pathologic nodal staging scores in patients treated with radical prostatectomy: a postoperative decision tool. <i>European Urology</i> , 2014 , 66, 439-46	10.2	18
152	Does increasing the nodal yield improve outcomes in contemporary patients without nodal metastasis undergoing radical prostatectomy?. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 47.e1-8	2.8	7
151	Prediction of 90-day mortality after radical cystectomy for bladder cancer in a prospective European multicenter cohort. <i>European Urology</i> , 2014 , 66, 156-63	10.2	114
150	The role of biomarkers in the assessment of prostate cancer risk prior to prostate biopsy: which markers matter and how should they be used?. <i>World Journal of Urology</i> , 2014 , 32, 871-80	4	7
149	Impact of preoperative anemia on oncologic outcomes of upper tract urothelial carcinoma treated with radical nephroureterectomy. <i>Journal of Urology</i> , 2014 , 191, 316-22	2.5	39
148	Re-assessment of 30-, 60- and 90-day mortality rates in non-metastatic prostate cancer patients treated either with radical prostatectomy or radiation therapy. <i>Canadian Urological Association Journal</i> , 2014 , 8, E75-80	1.2	4
147	Clinical nodal staging scores for prostate cancer: a proposal for preoperative risk assessment. <i>British Journal of Cancer</i> , 2014 , 111, 213-9	8.7	21
146	Prognosis of patients with pelvic lymph node (LN) metastasis after radical prostatectomy: value of extranodal extension and size of the largest LN metastasis. <i>BJU International</i> , 2014 , 114, 503-10	5.6	24
145	MALDI imaging-based identification of prognostically relevant signals in bladder cancer using large-scale tissue microarrays. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 1225-33	2.8	24
144	Predictors of 30-day acute kidney injury following radical and partial nephrectomy for renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 1259-66	2.8	42
143	256-MDCT for evaluation of urolithiasis: iterative reconstruction allows for a significant reduction of the applied radiation dose while maintaining high subjective and objective image quality. <i>Journal of Medical Imaging and Radiation Oncology</i> , 2014 , 58, 283-90	1.7	12
142	Predictors of survival in patients with disease recurrence after radical nephroureterectomy. <i>BJU International</i> , 2014 , 113, 911-7	5.6	17
141	External validation of the updated Briganti nomogram to predict lymph node invasion in prostate cancer patients undergoing extended lymph node dissection. <i>Prostate</i> , 2013 , 73, 211-8	4.2	45
140	Outcomes and prognostic factors in patients with a single lymph node metastasis at time of radical cystectomy. <i>BJU International</i> , 2013 , 111, 74-84	5.6	24
139	Accurate preoperative prediction of non-organ-confined bladder urothelial carcinoma at cystectomy. <i>BJU International</i> , 2013 , 111, 404-11	5.6	35

138	Female gender is associated with higher risk of disease recurrence in patients with primary T1 high-grade urothelial carcinoma of the bladder. <i>World Journal of Urology</i> , 2013 , 31, 1029-36	4	44
137	Prognostic value of alpha-methyl CoA racemase (AMACR) expression in renal cell carcinoma. <i>World Journal of Urology</i> , 2013 , 31, 847-53	4	10
136	Lymph node dissection during radical cystectomy for bladder cancer treatment: considerations on relevance and extent. <i>International Urology and Nephrology</i> , 2013 , 45, 1561-7	2.3	11
135	Reply: Inflammatory prognostic markers in clear cell renal cell carcinoma - preoperative C-reactive protein does not improve predictive accuracy. <i>BJU International</i> , 2013 , 111, E20-1	5.6	2
134	Impact of age and comorbidities on long-term survival of patients with high-risk prostate cancer treated with radical prostatectomy: a multi-institutional competing-risks analysis. <i>European Urology</i> , 2013 , 63, 693-701	10.2	78
133	Assays for prostate cancer : changing the screening paradigm?. <i>Molecular Diagnosis and Therapy</i> , 2013 , 17, 1-8	4.5	5
132	Epithelial cell adhesion molecule is an independent prognostic marker in clear cell renal carcinoma. <i>International Journal of Cancer</i> , 2013 , 132, 2948-55	7.5	22
131	Pathologic nodal staging score for bladder cancer: a decision tool for adjuvant therapy after radical cystectomy. <i>European Urology</i> , 2013 , 63, 371-8	10.2	39
130	Initial prostate biopsy: development and internal validation of a biopsy-specific nomogram based on the prostate cancer antigen 3 assay. <i>European Urology</i> , 2013 , 63, 201-9	10.2	96
129	Predictors of cancer-specific mortality after disease recurrence following radical cystectomy. <i>BJU International</i> , 2013 , 111, E30-6	5.6	60
128	Impact of smoking on oncologic outcomes of upper tract urothelial carcinoma after radical nephroureterectomy. <i>European Urology</i> , 2013 , 63, 1082-90	10.2	65
127	Loss of SPINK1 expression is associated with unfavorable outcomes in urothelial carcinoma of the bladder after radical cystectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 1716-24	2.8	13
126	The search for biomarkers of metastatic seminoma. <i>Journal of Urology</i> , 2013 , 190, 1046-51	2.5	8
125	Impact of smoking and smoking cessation on outcomes in bladder cancer patients treated with radical cystectomy. <i>European Urology</i> , 2013 , 64, 456-64	10.2	75
124	Impact of smoking and smoking cessation on oncologic outcomes in primary non-muscle-invasive bladder cancer. <i>European Urology</i> , 2013 , 63, 724-32	10.2	87
123	Obesity is associated with worse outcomes in patients with T1 high grade urothelial carcinoma of the bladder. <i>Journal of Urology</i> , 2013 , 190, 480-6	2.5	52
122	Impact of clinical and histopathological parameters on disease specific survival in patients with collecting duct renal cell carcinoma: development of a disease specific risk model. <i>Journal of Urology</i> , 2013 , 190, 458-63	2.5	24
121	The impact of intravesical gemcitabine and 1/3 dose Bacillus Calmette-Guérin instillation therapy on the quality of life in patients with nonmuscle invasive bladder cancer: results of a prospective, randomized, phase II trial. <i>Journal of Urology</i> , 2013 , 190, 857-62	2.5	37

120	PTEN deletions are related to disease progression and unfavourable prognosis in early bladder cancer. <i>Histopathology</i> , 2013 , 63, 670-7	7.3	19
119	Prediction of the risk of harboring prostate cancer by a prebiopsy nomogram based on extended biopsy protocol. <i>Urologia Internationalis</i> , 2013 , 90, 306-11	1.9	4
118	Short-term outcome and morbidity of different contemporary urethroplasty techniques--a preliminary comparison. <i>Journal of Endourology</i> , 2013 , 27, 925-9	2.7	16
117	The development of nomograms for stratification of men at risk of prostate cancer prior to prostate biopsy. <i>Biomarkers in Medicine</i> , 2013 , 7, 843-50	2.3	5
116	Gender-specific effect of smoking on upper tract urothelial carcinoma outcomes. <i>BJU International</i> , 2013 , 112, 623-37	5.6	22
115	Prostate Health Index (Phi) and Prostate Cancer Antigen 3 (PCA3) significantly improve prostate cancer detection at initial biopsy in a total PSA range of 2-10 ng/ml. <i>PLoS ONE</i> , 2013 , 8, e67687	3.7	71
114	Does low-risk prostate cancer detection change with repeat biopsies?. <i>European Urology</i> , 2012 , 61, 230-1	10.2	1
113	Clinical nodal staging scores for bladder cancer: a proposal for preoperative risk assessment. <i>European Urology</i> , 2012 , 61, 237-42	10.2	54
112	Death certificates are valid for the determination of cause of death in patients with upper and lower tract urothelial carcinoma. <i>European Urology</i> , 2012 , 61, 854-5	10.2	138
111	Prognostic role and HER2 expression of circulating tumor cells in peripheral blood of patients prior to radical cystectomy: a prospective study. <i>European Urology</i> , 2012 , 61, 810-7	10.2	119
110	Subclassification of pT3 urothelial carcinoma of the renal pelvicalyceal system is associated with recurrence-free and cancer-specific survival: proposal for a revision of the current TNM classification. <i>European Urology</i> , 2012 , 62, 224-31	10.2	33
109	Extended pelvic lymph node dissection does not affect erectile function recovery in patients treated with bilateral nerve-sparing radical prostatectomy. <i>Journal of Sexual Medicine</i> , 2012 , 9, 2187-94	1.1	17
108	A comparative performance analysis of total prostate-specific antigen, percentage free prostate-specific antigen, prostate-specific antigen velocity and urinary prostate cancer gene 3 in the first, second and third repeat prostate biopsy. <i>BJU International</i> , 2012 , 109, 1627-35	5.6	52
107	Higher perioperative morbidity and in-hospital mortality in patients with end-stage renal disease undergoing nephrectomy for non-metastatic kidney cancer: a population-based analysis. <i>BJU International</i> , 2012 , 110, E183-90	5.6	14
106	Risk assessment of metastatic recurrence in patients with prostate cancer by using the Cancer of the Prostate Risk Assessment score: results from 2937 European patients. <i>BJU International</i> , 2012 , 110, 1714-20	5.6	11
105	Does increasing the nodal yield improve outcomes in patients without nodal metastasis at radical cystectomy?. <i>World Journal of Urology</i> , 2012 , 30, 807-14	4	13
104	Transurethral holmium laser enucleation versus transurethral resection of the prostate and simple open prostatectomy--which procedure is faster?. <i>Journal of Urology</i> , 2012 , 187, 1608-13	2.5	44
103	Impact of smoking on outcomes of patients with a history of recurrent nonmuscle invasive bladder cancer. <i>Journal of Urology</i> , 2012 , 188, 2120-7	2.5	33

102	Risk of cancer-specific mortality following recurrence after radical nephroureterectomy. <i>Annals of Surgical Oncology</i> , 2012 , 19, 4337-44	3.1	42
101	Head to head comparison of nomograms predicting probability of lymph node invasion of prostate cancer in patients undergoing extended pelvic lymph node dissection. <i>Urology</i> , 2012 , 79, 546-51	1.6	32
100	Assessment of cancer control outcomes in patients with high-risk renal cell carcinoma treated with partial nephrectomy. <i>Urology</i> , 2012 , 80, 347-53	1.6	43
99	Protocol-based active surveillance for low-risk prostate cancer: anxiety levels in both men and their partners. <i>Urology</i> , 2012 , 80, 564-9	1.6	23
98	Predicting the risk of lymph node invasion during radical prostatectomy using the European Association of Urology guideline nomogram: a validation study. <i>European Journal of Surgical Oncology</i> , 2012 , 38, 624-9	3.6	6
97	Inflammatory prognostic markers in clear cell renal cell carcinoma - preoperative C-reactive protein does not improve predictive accuracy. <i>BJU International</i> , 2012 , 110, E771-7	5.6	14
96	Impact of histological variants on clinical outcomes of patients with upper urinary tract urothelial carcinoma. <i>Journal of Urology</i> , 2012 , 188, 398-404	2.5	78
95	National trends and disparities in the use of minimally invasive adult pyeloplasty. <i>Journal of Urology</i> , 2012 , 188, 913-8	2.5	31
94	Biomolecular predictors of urothelial cancer behavior and treatment outcomes. <i>Current Urology Reports</i> , 2012 , 13, 122-35	2.9	45
93	Reduced CD151 expression is related to advanced tumour stage in urothelial bladder cancer. <i>Pathology</i> , 2012 , 44, 448-52	1.6	10
92	Clinical evaluation of the PCA3 assay in guiding initial biopsy decisions. <i>Journal of Urology</i> , 2011 , 185, 2119-25	2.5	114
91	Detection of circulating tumour cells in peripheral blood of patients with advanced non-metastatic bladder cancer. <i>BJU International</i> , 2011 , 107, 1668-75	5.6	71
90	Pathological results and rates of treatment failure in high-risk prostate cancer patients after radical prostatectomy. <i>BJU International</i> , 2011 , 107, 765-770	5.6	99
89	Prediction of patient-specific risk and percentile cohort risk of pathological stage outcome using continuous prostate-specific antigen measurement, clinical stage and biopsy Gleason score. <i>BJU International</i> , 2011 , 107, 1562-9	5.6	31
88	Radical prostatectomy improves progression-free and cancer-specific survival in men with lymph node positive prostate cancer in the prostate-specific antigen era: a confirmatory study. <i>BJU International</i> , 2011 , 107, 1755-61	5.6	85
87	Older patients suffer from adverse histopathological features after radical cystectomy. <i>International Journal of Urology</i> , 2011 , 18, 576-84	2.3	12
86	Critical assessment of preoperative urinary prostate cancer antigen 3 on the accuracy of prostate cancer staging. <i>European Urology</i> , 2011 , 59, 96-105	10.2	112
85	Contemporary role of prostate cancer antigen 3 in the management of prostate cancer. <i>European Urology</i> , 2011 , 60, 1045-54	10.2	119

84	Genomic profiling of cell-free DNA in blood and bone marrow of prostate cancer patients. <i>Journal of Cancer Research and Clinical Oncology</i> , 2011 , 137, 811-9	4.9	23
83	Chronological age is not an independent predictor of clinical outcomes after radical nephroureterectomy. <i>World Journal of Urology</i> , 2011 , 29, 473-80	4	41
82	The presence of prostate cancer on saturation biopsy can be accurately predicted. <i>BJU International</i> , 2010 , 105, 636-41	5.6	16
81	Differences in histopathological and biochemical outcomes in patients with low Gleason score prostate cancer. <i>BJU International</i> , 2010 , 105, 818-23	5.6	5
80	[F(18)]-fluoroethylcholine combined in-line PET-CT scan for detection of lymph-node metastasis in high risk prostate cancer patients prior to radical prostatectomy: Preliminary results from a prospective histology-based study. <i>European Journal of Cancer</i> , 2010 , 46, 449-55	7.5	36
79	Biochemical recurrence after radical prostatectomy: multiplicative interaction between surgical margin status and pathological stage. <i>Journal of Urology</i> , 2010 , 184, 1341-6	2.5	68
78	Reply to Juan Morote Letter to the Editor re: Felix K. Chun, Alexandre de la Taille, Hendrik van Poppel, et al. Prostate Cancer Gene 3 (PCA3): Development and Internal Validation of a Novel Biopsy Nomogram. <i>Eur Urol</i> 2009;56:659-68. <i>European Urology</i> , 2010 , 57, e2-e3	10.2	0
77	Head-to-head comparison of the three most commonly used preoperative models for prediction of biochemical recurrence after radical prostatectomy. <i>European Urology</i> , 2010 , 57, 562-8	10.2	58
76	External validation of urinary PCA3-based nomograms to individually predict prostate biopsy outcome. <i>European Urology</i> , 2010 , 58, 727-32	10.2	88
75	Optimizing performance and interpretation of prostate biopsy: a critical analysis of the literature. <i>European Urology</i> , 2010 , 58, 851-64	10.2	83
74	External validation of a preoperative nomogram for prediction of the risk of recurrence after radical prostatectomy. <i>International Journal of Radiation Oncology Biology Physics</i> , 2010 , 77, 788-92	4	8
73	Unilateral prostate cancer cannot be accurately predicted in low-risk patients. <i>International Journal of Radiation Oncology Biology Physics</i> , 2010 , 77, 784-7	4	19
72	From gene to clinic: TMA-based clinical validation of molecular markers in prostate cancer. <i>Methods in Molecular Biology</i> , 2010 , 664, 177-89	1.4	8
71	Circulating prostate tumor cells detected by reverse transcription-PCR in men with localized or castration-refractory prostate cancer: concordance with CellSearch assay and association with bone metastases and with survival. <i>Clinical Chemistry</i> , 2009 , 55, 765-73	5.5	106
70	Assessment of pathological prostate cancer characteristics in men with favorable biopsy features on predominantly sextant biopsy. <i>European Urology</i> , 2009 , 55, 617-28-6	10.2	18
69	Molecular cancer phenotype in normal prostate tissue. <i>European Urology</i> , 2009 , 55, 885-90	10.2	21
68	Prevalence of a tertiary Gleason grade and its impact on adverse histopathologic parameters in a contemporary radical prostatectomy series. <i>European Urology</i> , 2009 , 55, 394-401	10.2	31
67	Risk-adjusted hazard rates of biochemical recurrence for prostate cancer patients after radical prostatectomy. <i>European Urology</i> , 2009 , 55, 412-19	10.2	16

66	Prostate cancer gene 3 (PCA3): development and internal validation of a novel biopsy nomogram. <i>European Urology</i> , 2009 , 56, 659-67	10.2	142
65	Prediction of sexual function after radical prostatectomy. <i>Cancer</i> , 2009 , 115, 3150-9	6.4	31
64	Effect of body mass index on histopathologic parameters: results of large European contemporary consecutive open radical prostatectomy series. <i>Urology</i> , 2009 , 73, 615-9	1.6	27
63	Predictive value of prostate-specific antigen expression in prostate cancer: a tissue microarray study. <i>Urology</i> , 2009 , 74, 1169-73	1.6	17
62	Tissue factor procoagulant activity of plasma microparticles is increased in patients with early-stage prostate cancer. <i>Thrombosis and Haemostasis</i> , 2009 , 101, 1147-1155	7	60
61	Nomogram predicting the probability of early recurrence after radical prostatectomy for prostate cancer. <i>Journal of Urology</i> , 2009 , 181, 601-7; discussion 607-8	2.5	111
60	A nomogram is more accurate than a regression tree in predicting lymph node invasion in prostate cancer. <i>BJU International</i> , 2008 , 101, 556-60	5.6	21
59	Comparison of stage migration patterns between Europe and the USA: an analysis of 11 350 men treated with radical prostatectomy for prostate cancer. <i>BJU International</i> , 2008 , 101, 1513-8	5.6	122
58	Microsatellite analysis of allelic imbalance in tumour and blood from patients with prostate cancer. <i>BJU International</i> , 2008 , 102, 253-8	5.6	35
57	Marked gene transcript level alterations occur early during radical prostatectomy. <i>European Urology</i> , 2008 , 53, 333-44	10.2	31
56	Contemporary prostate cancer prevalence among T1c biopsy-referred men with a prostate-specific antigen level European Urology, 2008 , 53, 750-7	10.2	19
55	Evaluation of prostate cancer detection with ultrasound real-time elastography: a comparison with step section pathological analysis after radical prostatectomy. <i>European Urology</i> , 2008 , 54, 1354-62	10.2	194
54	Editorial comment on: preliminary results of a novel method to estimate the probability of prostate cancer in men with elevated serum PSA values. <i>European Urology</i> , 2008 , 54, 702	10.2	
53	Impact of surgical volume on the rate of lymph node metastases in patients undergoing radical prostatectomy and extended pelvic lymph node dissection for clinically localized prostate cancer. <i>European Urology</i> , 2008 , 54, 794-802	10.2	52
52	Sequential use of the tyrosine kinase inhibitors sorafenib and sunitinib in metastatic renal cell carcinoma: a retrospective outcome analysis. <i>European Urology</i> , 2008 , 54, 1373-8	10.2	77
51	Biopsies performed at tertiary care centers are superior to referral biopsies in predicting pathologic Gleason sum. <i>Journal of Endourology</i> , 2008 , 22, 533-8	2.7	6
50	Critical assessment of tools to predict clinically insignificant prostate cancer at radical prostatectomy in contemporary men. <i>Cancer</i> , 2008 , 113, 701-9	6.4	74
49	Currently used criteria for active surveillance in men with low-risk prostate cancer: an analysis of pathologic features. <i>Cancer</i> , 2008 , 113, 2068-72	6.4	88

48	Percent free prostate-specific antigen (PSA) is an accurate predictor of prostate cancer risk in men with serum PSA 2.5 ng/mL and lower. <i>Cancer</i> , 2008 , 113, 2695-703	6.4	26
47	Development and external validation of an extended repeat biopsy nomogram. <i>Journal of Urology</i> , 2007 , 177, 510-5	2.5	72
46	Plasma tissue factor antigen in localized prostate cancer: Distribution, clinical significance and correlation with haemostatic activation markers. <i>Thrombosis and Haemostasis</i> , 2007 , 97, 464-470	7	30
45	Detection of tumor-specific DNA in blood and bone marrow plasma from patients with prostate cancer. <i>International Journal of Cancer</i> , 2007 , 120, 1465-71	7.5	48
44	Obesity does not predispose to more aggressive prostate cancer either at biopsy or radical prostatectomy in European men. <i>International Journal of Cancer</i> , 2007 , 121, 791-5	7.5	42
43	A critical appraisal of logistic regression-based nomograms, artificial neural networks, classification and regression-tree models, look-up tables and risk-group stratification models for prostate cancer. <i>BJU International</i> , 2007 , 99, 794-800	5.6	88
42	Health-insurance status is a determinant of the stage at presentation and of cancer control in European men treated with radical prostatectomy for clinically localized prostate cancer. <i>BJU International</i> , 2007 , 99, 1404-8	5.6	15
41	Distribution of prostate specific antigen (PSA) and percentage free PSA in a contemporary screening cohort with no evidence of prostate cancer. <i>BJU International</i> , 2007 , 100, 37-41	5.6	16
40	Differences in the rate of lymph node invasion in men with clinically localized prostate cancer might be related to the continent of origin. <i>BJU International</i> , 2007 , 100, 528-32	5.6	32
39	Effect of autologous blood transfusion on the rate of biochemical recurrence after radical prostatectomy. <i>BJU International</i> , 2007 , 100, 1249-53	5.6	13
38	A nomogram for staging of exclusive nonobturator lymph node metastases in men with localized prostate cancer. <i>European Urology</i> , 2007 , 51, 112-9; discussion 119-20	10.2	54
37	Zonal origin of localized prostate cancer does not affect the rate of biochemical recurrence after radical prostatectomy. <i>European Urology</i> , 2007 , 51, 949-55; discussion 955	10.2	16
36	Initial biopsy outcome prediction--head-to-head comparison of a logistic regression-based nomogram versus artificial neural network. <i>European Urology</i> , 2007 , 51, 1236-40; discussion 1241-3	10.2	66
35	Development and external validation of an extended 10-core biopsy nomogram. <i>European Urology</i> , 2007 , 52, 436-44	10.2	91
34	Reply to Carsten Stephan et al Letter to the Editor re: Felix K.-H. Chun, Markus Graefen, Alberto Briganti, Andrea Gallina, Julia Hopp, Michael W. Kattan, Hartwig Huland and Pierre I. Karakiewicz. Initial Biopsy Outcome Prediction Head-to-Head Comparison of a Logistic Regression-Based Nomogram versus Artificial Neural Network. <i>Eur Urol</i> 2007;51:1236-43. <i>European Urology</i> , 2007 , 51, 1448	10.2	
33	Prediction of pathological stage is inaccurate in men with PSA values above 20 ng/mL. <i>European Urology</i> , 2007 , 52, 1374-80	10.2	16
32	Percentage of positive biopsy cores can improve the ability to predict lymph node invasion in patients undergoing radical prostatectomy and extended pelvic lymph node dissection. <i>European Urology</i> , 2007 , 51, 1573-81	10.2	68
31	Systematic assessment of the ability of the number and percentage of positive biopsy cores to predict pathologic stage and biochemical recurrence after radical prostatectomy. <i>European Urology</i> , 2007 , 52, 733-43	10.2	29

30	Prostate-specific antigen improves the ability of clinical stage and biopsy Gleason sum to predict the pathologic stage at radical prostatectomy in the new millennium. <i>European Urology</i> , 2007 , 52, 1067-74	10.2	17
29	Reply to Ian Beckley and Masood A. Khan Letter to the Editor re: Felix K.-H. Chun, Thomas Steuber, Andreas Erbersdobler, et al. Development and Internal Validation of a Nomogram Predicting the Probability of Prostate Cancer Gleason Sum Upgrading Between Biopsy and Radical Prostatectomy Pathology. <i>Eur Urol</i> 2006;49:820-6. <i>European Urology</i> , 2007 , 52, 1271	10.2	
28	Role of nomograms for prostate cancer in 2007. <i>World Journal of Urology</i> , 2007 , 25, 131-42	4	39
27	Management of erectile dysfunction after radical prostatectomy in 2007. <i>World Journal of Urology</i> , 2007 , 25, 143-8	4	29
26	Critical assessment of ideal nodal yield at pelvic lymphadenectomy to accurately diagnose prostate cancer nodal metastasis in patients undergoing radical retropubic prostatectomy. <i>Urology</i> , 2007 , 69, 147-51	1.6	131
25	Body mass index does not improve the ability to predict biochemical recurrence after radical prostatectomy. <i>European Journal of Cancer</i> , 2007 , 43, 375-82	7.5	20
24	Tumour volume and high grade tumour volume are the best predictors of pathologic stage and biochemical recurrence after radical prostatectomy. <i>European Journal of Cancer</i> , 2007 , 43, 536-43	7.5	69
23	Prostate volume and adverse prostate cancer features: fact not artifact. <i>European Journal of Cancer</i> , 2007 , 43, 2669-77	7.5	66
22	Development and internal validation of a nomogram predicting the probability of prostate cancer Gleason sum upgrading between biopsy and radical prostatectomy pathology. <i>European Urology</i> , 2006 , 49, 820-6	10.2	157
21	The 2002 AJCC pT2 substages confer no prognostic information on the rate of biochemical recurrence after radical prostatectomy. <i>European Urology</i> , 2006 , 49, 273-8; discussion 278-9	10.2	41
20	Validation of a nomogram predicting the probability of lymph node invasion among patients undergoing radical prostatectomy and an extended pelvic lymphadenectomy. <i>European Urology</i> , 2006 , 49, 1019-26; discussion 1026-7	10.2	180
19	Tumor size improves the accuracy of TNM predictions in patients with renal cancer. <i>European Urology</i> , 2006 , 50, 521-8; discussion 529	10.2	50
18	Frozen section for the management of intraoperatively detected palpable tumor lesions during nerve-sparing scheduled radical prostatectomy. <i>European Urology</i> , 2006 , 49, 1011-6; discussion 1016-8	10.2	49
17	High radical prostatectomy surgical volume is related to lower radical prostatectomy total hospital charges. <i>European Urology</i> , 2006 , 50, 58-62; discussion 62-3	10.2	18
16	High incidence of prostate cancer detected by saturation biopsy after previous negative biopsy series. <i>European Urology</i> , 2006 , 50, 498-505	10.2	159
15	Outcomes research: a methodologic review. <i>European Urology</i> , 2006 , 50, 218-24	10.2	14
14	Prostate cancer nomograms: an update. <i>European Urology</i> , 2006 , 50, 914-26; discussion 926	10.2	65
13	Complications and other surgical outcomes associated with extended pelvic lymphadenectomy in men with localized prostate cancer. <i>European Urology</i> , 2006 , 50, 1006-13	10.2	266

12	Risk assessment for biochemical recurrence prior to radical prostatectomy: significant enhancement contributed by human glandular kallikrein 2 (hK2) and free prostate specific antigen (PSA) in men with moderate PSA-elevation in serum. <i>International Journal of Cancer</i> , 2006 , 118, 1234-40	7.5	44
11	A comparative review of apomorphine formulations for erectile dysfunction : recommendations for use in the elderly. <i>Drugs and Aging</i> , 2006 , 23, 309-19	4.7	3
10	25-year prostate cancer control and survival outcomes: a 40-year radical prostatectomy single institution series. <i>Journal of Urology</i> , 2006 , 176, 569-74	2.5	103
9	Validation of a nomogram for prediction of side specific extracapsular extension at radical prostatectomy. <i>Journal of Urology</i> , 2006 , 175, 939-44; discussion 944	2.5	128
8	Development and internal validation of preoperative transition zone prostate cancer nomogram. <i>Urology</i> , 2006 , 68, 1295-300	1.6	15
7	Prostate specific-antigen distribution in asymptomatic Canadian men with no clinical evidence of prostate cancer. <i>BJU International</i> , 2006 , 98, 50-3	5.6	10
6	Significant upgrading affects a third of men diagnosed with prostate cancer: predictive nomogram and internal validation. <i>BJU International</i> , 2006 , 98, 329-34	5.6	110
5	Validation of a nomogram predicting the probability of lymph node invasion based on the extent of pelvic lymphadenectomy in patients with clinically localized prostate cancer. <i>BJU International</i> , 2006 , 98, 788-93	5.6	137
4	Circulating tumour-associated plasma DNA represents an independent and informative predictor of prostate cancer. <i>BJU International</i> , 2006 , 98, 544-8	5.6	92
3	Surgical volume is related to the rate of positive surgical margins at radical prostatectomy in European patients. <i>BJU International</i> , 2006 , 98, 1204-9	5.6	56
2	Anatomic radical retropubic prostatectomy-long-term recurrence-free survival rates for localized prostate cancer. <i>World Journal of Urology</i> , 2006 , 24, 273-80	4	112
1	A multicenter clinical trial on the use of (-5, -7) pro prostate specific antigen. <i>Journal of Urology</i> , 2005 , 174, 2150-3	2.5	34