

# Felix Kh Chun

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1670696/publications.pdf>

Version: 2024-02-01

177  
papers

9,542  
citations

24978

57  
h-index

45213

90  
g-index

191  
all docs

191  
docs citations

191  
times ranked

7397  
citing authors

#	ARTICLE	IF	CITATIONS
1	Complications and Other Surgical Outcomes Associated with Extended Pelvic Lymphadenectomy in Men with Localized Prostate Cancer. <i>European Urology</i> , 2006, 50, 1006-1013.	0.9	341
2	Evaluation of Prostate Cancer Detection with Ultrasound Real-Time Elastography: A Comparison with Step Section Pathological Analysis after Radical Prostatectomy. <i>European Urology</i> , 2008, 54, 1354-1362.	0.9	226
3	Validation of a Nomogram Predicting the Probability of Lymph Node Invasion among Patients Undergoing Radical Prostatectomy and an Extended Pelvic Lymphadenectomy. <i>European Urology</i> , 2006, 49, 1019-1027.	0.9	215
4	Development and Internal Validation of a Nomogram Predicting the Probability of Prostate Cancer Gleason Sum Upgrading Between Biopsy and Radical Prostatectomy Pathology. <i>European Urology</i> , 2006, 49, 820-826.	0.9	188
5	High Incidence of Prostate Cancer Detected by Saturation Biopsy after Previous Negative Biopsy Series. <i>European Urology</i> , 2006, 50, 498-505.	0.9	178
6	Validation of a Nomogram for Prediction of Side Specific Extracapsular Extension at Radical Prostatectomy. <i>Journal of Urology</i> , 2006, 175, 939-944.	0.2	163
7	Prognostic Role and HER2 Expression of Circulating Tumor Cells in Peripheral Blood of Patients Prior to Radical Cystectomy: A Prospective Study. <i>European Urology</i> , 2012, 61, 810-817.	0.9	163
8	Validation of a nomogram predicting the probability of lymph node invasion based on the extent of pelvic lymphadenectomy in patients with clinically localized prostate cancer. <i>BJU International</i> , 2006, 98, 788-793.	1.3	162
9	Prostate Cancer Gene 3 (PCA3): Development and Internal Validation of a Novel Biopsy Nomogram. <i>European Urology</i> , 2009, 56, 659-668.	0.9	161
10	Critical Assessment of Ideal Nodal Yield at Pelvic Lymphadenectomy to Accurately Diagnose Prostate Cancer Nodal Metastasis in Patients Undergoing Radical Retropubic Prostatectomy. <i>Urology</i> , 2007, 69, 147-151.	0.5	156
11	Prediction of 90-day Mortality After Radical Cystectomy for Bladder Cancer in a Prospective European Multicenter Cohort. <i>European Urology</i> , 2014, 66, 156-163.	0.9	156
12	Death Certificates Are Valid for the Determination of Cause of Death in Patients With Upper and Lower Tract Urothelial Carcinoma. <i>European Urology</i> , 2012, 61, 854-855.	0.9	152
13	Contemporary Role of Prostate Cancer Antigen 3 in the Management of Prostate Cancer. <i>European Urology</i> , 2011, 60, 1045-1054.	0.9	148
14	Clinical Evaluation of the PCA3 Assay in Guiding Initial Biopsy Decisions. <i>Journal of Urology</i> , 2011, 185, 2119-2125.	0.2	136
15	Anatomic radical retropubic prostatectomyâ€™long-term recurrence-free survival rates for localized prostate cancer. <i>World Journal of Urology</i> , 2006, 24, 273-280.	1.2	134
16	Comparison of stage migration patterns between Europe and the USA: an analysis of 11Â350 men treated with radical prostatectomy for prostate cancer. <i>BJU International</i> , 2008, 101, 1513-1518.	1.3	134
17	Nomogram Predicting the Probability of Early Recurrence After Radical Prostatectomy for Prostate Cancer. <i>Journal of Urology</i> , 2009, 181, 601-608.	0.2	129
18	Critical Assessment of Preoperative Urinary Prostate Cancer Antigen 3 on the Accuracy of Prostate Cancer Staging. <i>European Urology</i> , 2011, 59, 96-105.	0.9	127

#	ARTICLE	IF	CITATIONS
19	Significant upgrading affects a third of men diagnosed with prostate cancer: predictive nomogram and internal validation. <i>BJU International</i> , 2006, 98, 329-334.	1.3	126
20	Circulating Prostate Tumor Cells Detected by Reverse Transcription-PCR in Men with Localized or Castration-Refractory Prostate Cancer: Concordance with CellSearch Assay and Association with Bone Metastases and with Survival. <i>Clinical Chemistry</i> , 2009, 55, 765-773.	1.5	122
21	Pathological results and rates of treatment failure in high-risk prostate cancer patients after radical prostatectomy. <i>BJU International</i> , 2011, 107, 765-770.	1.3	120
22	25-Year Prostate Cancer Control and Survival Outcomes: A 40-Year Radical Prostatectomy Single Institution Series. <i>Journal of Urology</i> , 2006, 176, 569-574.	0.2	119
23	Development and External Validation of an Extended 10-Core Biopsy Nomogram. <i>European Urology</i> , 2007, 52, 436-445.	0.9	114
24	Impact of Histological Variants on Clinical Outcomes of Patients with Upper Urinary Tract Urothelial Carcinoma. <i>Journal of Urology</i> , 2012, 188, 398-404.	0.2	114
25	Initial Prostate Biopsy: Development and Internal Validation of a Biopsy-specific Nomogram Based on the Prostate Cancer Antigen 3 Assay. <i>European Urology</i> , 2013, 63, 201-209.	0.9	114
26	A critical appraisal of logistic regression-based nomograms, artificial neural networks, classification and regression-tree models, look-up tables and risk-group stratification models for prostate cancer. <i>BJU International</i> , 2007, 99, 794-800.	1.3	111
27	Radical prostatectomy improves progression-free and cancer-specific survival in men with lymph node positive prostate cancer in the prostate-specific antigen era: a confirmatory study. <i>BJU International</i> , 2011, 107, 1755-1761.	1.3	105
28	Impact of Smoking and Smoking Cessation on Oncologic Outcomes in Primary Non-muscle-invasive Bladder Cancer. <i>European Urology</i> , 2013, 63, 724-732.	0.9	105
29	Circulating tumour-associated plasma DNA represents an independent and informative predictor of prostate cancer. <i>BJU International</i> , 2006, 98, 544-548.	1.3	104
30	Impact of Smoking and Smoking Cessation on Outcomes in Bladder Cancer Patients Treated with Radical Cystectomy. <i>European Urology</i> , 2013, 64, 456-464.	0.9	101
31	Natural history of surgically treated high-risk prostate cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015, 33, 163.e7-163.e13.	0.8	101
32	Local Therapy Improves Survival in Metastatic Prostate Cancer. <i>European Urology</i> , 2017, 72, 118-124.	0.9	100
33	Impact of Age and Comorbidities on Long-term Survival of Patients with High-risk Prostate Cancer Treated with Radical Prostatectomy: A Multi-institutional Competing-risks Analysis. <i>European Urology</i> , 2013, 63, 693-701.	0.9	98
34	Impact of Smoking on Oncologic Outcomes of Upper Tract Urothelial Carcinoma After Radical Nephroureterectomy. <i>European Urology</i> , 2013, 63, 1082-1090.	0.9	98
35	Currently used criteria for active surveillance in men with low-risk prostate cancer. <i>Cancer</i> , 2008, 113, 2068-2072.	2.0	96
36	External Validation of Urinary PCA3-Based Nomograms to Individually Predict Prostate Biopsy Outcome. <i>European Urology</i> , 2010, 58, 727-732.	0.9	96

#	ARTICLE	IF	CITATIONS
37	Optimizing Performance and Interpretation of Prostate Biopsy: A Critical Analysis of the Literature. <i>European Urology</i> , 2010, 58, 851-864.	0.9	96
38	Sequential Use of the Tyrosine Kinase Inhibitors Sorafenib and Sunitinib in Metastatic Renal Cell Carcinoma: A Retrospective Outcome Analysis. <i>European Urology</i> , 2008, 54, 1373-1378.	0.9	91
39	Prostate Cancer Nomograms: An Update. <i>European Urology</i> , 2006, 50, 914-926.	0.9	89
40	Detection of circulating tumour cells in peripheral blood of patients with advanced non-metastatic bladder cancer. <i>BJU International</i> , 2011, 107, 1668-1675.	1.3	89
41	Prostate Health Index (Phi) and Prostate Cancer Antigen 3 (PCA3) Significantly Improve Prostate Cancer Detection at Initial Biopsy in a Total PSA Range of 2-10 ng/ml. <i>PLoS ONE</i> , 2013, 8, e67687.	1.1	87
42	Critical assessment of tools to predict clinically insignificant prostate cancer at radical prostatectomy in contemporary men. <i>Cancer</i> , 2008, 113, 701-709.	2.0	86
43	Percentage of Positive Biopsy Cores Can Improve the Ability to Predict Lymph Node Invasion in Patients Undergoing Radical Prostatectomy and Extended Pelvic Lymph Node Dissection. <i>European Urology</i> , 2007, 51, 1573-1581.	0.9	84
44	Biochemical Recurrence After Radical Prostatectomy: Multiplicative Interaction Between Surgical Margin Status and Pathological Stage. <i>Journal of Urology</i> , 2010, 184, 1341-1346.	0.2	84
45	Prostate volume and adverse prostate cancer features: Fact not artifact. <i>European Journal of Cancer</i> , 2007, 43, 2669-2677.	1.3	82
46	Initial Biopsy Outcome Prediction—Head-to-Head Comparison of a Logistic Regression-Based Nomogram versus Artificial Neural Network. <i>European Urology</i> , 2007, 51, 1236-1243.	0.9	79
47	Tumour volume and high grade tumour volume are the best predictors of pathologic stage and biochemical recurrence after radical prostatectomy. <i>European Journal of Cancer</i> , 2007, 43, 536-543.	1.3	77
48	Predictors of cancer-specific mortality after disease recurrence following radical cystectomy. <i>BJU International</i> , 2013, 111, E30-6.	1.3	77
49	Development and External Validation of an Extended Repeat Biopsy Nomogram. <i>Journal of Urology</i> , 2007, 177, 510-515.	0.2	75
50	Head-to-Head Comparison of the Three Most Commonly Used Preoperative Models for Prediction of Biochemical Recurrence After Radical Prostatectomy. <i>European Urology</i> , 2010, 57, 562-568.	0.9	69
51	Clinical Nodal Staging Scores for Bladder Cancer: A Proposal for Preoperative Risk Assessment. <i>European Urology</i> , 2012, 61, 237-242.	0.9	69
52	Frozen Section for the Management of Intraoperatively Detected Palpable Tumor Lesions During Nerve-Sparing Scheduled Radical Prostatectomy. <i>European Urology</i> , 2006, 49, 1011-1018.	0.9	67
53	Tissue factor procoagulant activity of plasma microparticles is increased in patients with early-stage prostate cancer. <i>Thrombosis and Haemostasis</i> , 2009, 101, 1147-1155.	1.8	67
54	A Nomogram for Staging of Exclusive Nonobturator Lymph Node Metastases in Men with Localized Prostate Cancer. <i>European Urology</i> , 2007, 51, 112-120.	0.9	66

#	ARTICLE	IF	CITATIONS
55	Obesity is Associated with Worse Outcomes in Patients with T1 High Grade Urothelial Carcinoma of the Bladder. <i>Journal of Urology</i> , 2013, 190, 480-486.	0.2	66
56	Surgical volume is related to the rate of positive surgical margins at radical prostatectomy in European patients. <i>BJU International</i> , 2006, 98, 1204-1209.	1.3	62
57	Chronological age is not an independent predictor of clinical outcomes after radical nephroureterectomy. <i>World Journal of Urology</i> , 2011, 29, 473-480.	1.2	62
58	Impact of Surgical Volume on the Rate of Lymph Node Metastases in Patients Undergoing Radical Prostatectomy and Extended Pelvic Lymph Node Dissection for Clinically Localized Prostate Cancer. <i>European Urology</i> , 2008, 54, 794-804.	0.9	61
59	Tumor Size Improves the Accuracy of TNM Predictions in Patients with Renal Cancer. <i>European Urology</i> , 2006, 50, 521-529.	0.9	60
60	A comparative performance analysis of total prostate-specific antigen, percentage free prostate-specific antigen, prostate-specific antigen velocity and urinary prostate cancer gene 3 in the first, second and third repeat prostate biopsy. <i>BJU International</i> , 2012, 109, 1627-1635.	1.3	59
61	The Impact of Intravesical Gemcitabine and 1/3 Dose Bacillus Calmette-Guérin Instillation Therapy on the Quality of Life in Patients with Nonmuscle Invasive Bladder Cancer: Results of a Prospective, Randomized, Phase II Trial. <i>Journal of Urology</i> , 2013, 190, 857-862.	0.2	58
62	Female gender is associated with higher risk of disease recurrence in patients with primary T1 high-grade urothelial carcinoma of the bladder. <i>World Journal of Urology</i> , 2013, 31, 1029-1036.	1.2	55
63	Detection of tumor-specific DNA in blood and bone marrow plasma from patients with prostate cancer. <i>International Journal of Cancer</i> , 2007, 120, 1465-1471.	2.3	54
64	Risk of Cancer-specific Mortality following Recurrence After Radical Nephroureterectomy. <i>Annals of Surgical Oncology</i> , 2012, 19, 4337-4344.	0.7	53
65	Transurethral Holmium Laser Enucleation Versus Transurethral Resection of the Prostate and Simple Open Prostatectomy—Which Procedure is Faster?. <i>Journal of Urology</i> , 2012, 187, 1608-1613.	0.2	51
66	Biomolecular Predictors of Urothelial Cancer Behavior and Treatment Outcomes. <i>Current Urology Reports</i> , 2012, 13, 122-135.	1.0	51
67	External validation of the updated briganti nomogram to predict lymph node invasion in prostate cancer patients undergoing extended lymph node dissection. <i>Prostate</i> , 2013, 73, 211-218.	1.2	51
68	Predictors of 30-day acute kidney injury following radical and partial nephrectomy for renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014, 32, 1259-1266.	0.8	50
69	Assessment of Cancer Control Outcomes in Patients With High-risk Renal Cell Carcinoma Treated With Partial Nephrectomy. <i>Urology</i> , 2012, 80, 347-353.	0.5	49
70	Impact of Preoperative Anemia on Oncologic Outcomes of Upper Tract Urothelial Carcinoma Treated with Radical Nephroureterectomy. <i>Journal of Urology</i> , 2014, 191, 316-322.	0.2	49
71	Risk assessment for biochemical recurrence prior to radical prostatectomy: Significant enhancement contributed by human glandular kallikrein 2 (hk2) and free prostate specific antigen (PSA) in men with moderate PSA-elevation in serum. <i>International Journal of Cancer</i> , 2006, 118, 1234-1240.	2.3	48
72	Role of nomograms for prostate cancer in 2007. <i>World Journal of Urology</i> , 2007, 25, 131-142.	1.2	48

#	ARTICLE	IF	CITATIONS
73	Accurate preoperative prediction of non-organ-confined bladder urothelial carcinoma at cystectomy. <i>BJU International</i> , 2013, 111, 404-411.	1.3	48
74	Pathologic Nodal Staging Score for Bladder Cancer: A Decision Tool for Adjuvant Therapy After Radical Cystectomy. <i>European Urology</i> , 2013, 63, 371-378.	0.9	47
75	The 2002 AJCC pT2 Substages Confer No Prognostic Information on the Rate of Biochemical Recurrence After Radical Prostatectomy. <i>European Urology</i> , 2006, 49, 273-279.	0.9	45
76	Impact of Smoking on Outcomes of Patients with a History of Recurrent Nonmuscle Invasive Bladder Cancer. <i>Journal of Urology</i> , 2012, 188, 2120-2128.	0.2	45
77	Obesity does not predispose to more aggressive prostate cancer either at biopsy or radical prostatectomy in European men. <i>International Journal of Cancer</i> , 2007, 121, 791-795.	2.3	44
78	National Trends and Disparities in the Use of Minimally Invasive Adult Pyeloplasty. <i>Journal of Urology</i> , 2012, 188, 913-918.	0.2	44
79	Subclassification of pT3 Urothelial Carcinoma of the Renal Pelvicalyceal System is Associated With Recurrence-Free and Cancer-Specific Survival: Proposal for a Revision of the Current TNM Classification. <i>European Urology</i> , 2012, 62, 224-231.	0.9	44
80	A MULTICENTER CLINICAL TRIAL ON THE USE OF (â€“5, â€“7) PRO PROSTATE SPECIFIC ANTIGEN. <i>Journal of Urology</i> , 2005, 174, 2150-2153.	0.2	41
81	Marked Gene Transcript Level Alterations Occur Early During Radical Prostatectomy. <i>European Urology</i> , 2008, 53, 333-346.	0.9	40
82	[F18]-fluoroethylcholine combined in-line PET-CT scan for detection of lymph-node metastasis in high risk prostate cancer patients prior to radical prostatectomy: Preliminary results from a prospective histology-based study. <i>European Journal of Cancer</i> , 2010, 46, 449-455.	1.3	39
83	Microsatellite analysis of allelic imbalance in tumour and blood from patients with prostate cancer. <i>BJU International</i> , 2008, 102, 253-258.	1.3	38
84	Management of erectile dysfunction after radical prostatectomy in 2007. <i>World Journal of Urology</i> , 2007, 25, 143-148.	1.2	37
85	Prediction of patient-specific risk and percentile cohort risk of pathological stage outcome using continuous prostate-specific antigen measurement, clinical stage and biopsy Gleason score. <i>BJU International</i> , 2011, 107, 1562-1569.	1.3	36
86	The Effect of Resident Involvement on Perioperative Outcomes in Transurethral Urologic Surgeries. <i>Journal of Surgical Education</i> , 2015, 72, 1018-1025.	1.2	36
87	Plasma tissue factor antigen in localized prostate cancer: Distribution, clinical significance and correlation with haemostatic activation markers. <i>Thrombosis and Haemostasis</i> , 2007, 97, 464-470.	1.8	35
88	Systematic Assessment of the Ability of the Number and Percentage of Positive Biopsy Cores to Predict Pathologic Stage and Biochemical Recurrence after Radical Prostatectomy. <i>European Urology</i> , 2007, 52, 733-745.	0.9	35
89	Prevalence of a Tertiary Gleason Grade and Its Impact on Adverse Histopathologic Parameters in a Contemporary Radical Prostatectomy Series. <i>European Urology</i> , 2009, 55, 394-403.	0.9	35
90	Head to Head Comparison of Nomograms Predicting Probability of Lymph Node Invasion of Prostate Cancer in Patients Undergoing Extended Pelvic Lymph Node Dissection. <i>Urology</i> , 2012, 79, 546-551.	0.5	34

#	ARTICLE	IF	CITATIONS
91	Differences in the rate of lymph node invasion in men with clinically localized prostate cancer might be related to the continent of origin. <i>BJU International</i> , 2007, 100, 528-532.	1.3	33
92	Prediction of sexual function after radical prostatectomy. <i>Cancer</i> , 2009, 115, 3150-3159.	2.0	33
93	Prognosis of patients with pelvic lymph node (<sc>LN</sc>) metastasis after radical prostatectomy: Value of extranodal extension and size of the largest <sc>LN</sc> metastasis. <i>BJU International</i> , 2014, 114, 503-510.	1.3	33
94	Evidence from the â€˜PROspective MulticEnTer Radical Cystectomy Series 2011 (PROMETRICS 2011)â€™ Study: How are Preoperative Patient Characteristics Associated with Urinary Diversion Type After Radical Cystectomy for Bladder Cancer?. <i>Annals of Surgical Oncology</i> , 2015, 22, 1032-1042.	0.7	33
95	Percent free prostateâ€™specific antigen (PSA) is an accurate predictor of prostate cancer risk in men with serum PSA 2.5 ng/mL and lower. <i>Cancer</i> , 2008, 113, 2695-2703.	2.0	32
96	Impact of Clinical and Histopathological Parameters on Disease Specific Survival in Patients with Collecting Duct Renal Cell Carcinoma: Development of a Disease Specific Risk Model. <i>Journal of Urology</i> , 2013, 190, 458-463.	0.2	31
97	Genderâ€™specific effect of smoking on upper tract urothelial carcinoma outcomes. <i>BJU International</i> , 2013, 112, 623-637.	1.3	31
98	Effect of Body Mass Index on Histopathologic Parameters: Results of Large European Contemporary Consecutive Open Radical Prostatectomy Series. <i>Urology</i> , 2009, 73, 615-619.	0.5	30
99	<i><sc>PTEN</sc></i> deletions are related to disease progression and unfavourable prognosis in early bladder cancer. <i>Histopathology</i> , 2013, 63, 670-677.	1.6	30
100	Predictors of survival in patients with disease recurrence after radical nephroureterectomy. <i>BJU International</i> , 2014, 113, 911-917.	1.3	28
101	MALDI imagingâ€™based identification of prognostically relevant signals in bladder cancer using large-scale tissue microarrays1These authors contributed equally to this work.. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014, 32, 1225-1233.	0.8	27
102	Protocol-based Active Surveillance for Low-risk Prostate Cancer: Anxiety Levels in Both Men and Their Partners. <i>Urology</i> , 2012, 80, 564-569.	0.5	26
103	Outcomes and prognostic factors in patients with a single lymph node metastasis at time of radical cystectomy. <i>BJU International</i> , 2013, 111, 74-84.	1.3	26
104	Assessment of Pathological Prostate Cancer Characteristics in Men with Favorable Biopsy Features on Predominantly Sextant Biopsy. <i>European Urology</i> , 2009, 55, 617-628.	0.9	25
105	Genomic profiling of cell-free DNA in blood and bone marrow of prostate cancer patients. <i>Journal of Cancer Research and Clinical Oncology</i> , 2011, 137, 811-819.	1.2	25
106	Epithelial cell adhesion molecule is an independent prognostic marker in clear cell renal carcinoma. <i>International Journal of Cancer</i> , 2013, 132, 2948-2955.	2.3	25
107	A nomogram is more accurate than a regression tree in predicting lymph node invasion in prostate cancer. <i>BJU International</i> , 2008, 101, 556-560.	1.3	24
108	Contemporary Prostate Cancer Prevalence among T1c Biopsy-Referred Men with a Prostate-Specific Antigen Level â‰¥ 4.0ng per Milliliter. <i>European Urology</i> , 2008, 53, 750-757.	0.9	24

#	ARTICLE	IF	CITATIONS
109	Unilateral Prostate Cancer Cannot be Accurately Predicted in Low-Risk Patients. International Journal of Radiation Oncology Biology Physics, 2010, 77, 784-787.	0.4	24
110	Clinical nodal staging scores for prostate cancer: a proposal for preoperative risk assessment. British Journal of Cancer, 2014, 111, 213-219.	2.9	24
111	Pathologic Nodal Staging Scores in Patients Treated with Radical Prostatectomy: A Postoperative Decision Tool. European Urology, 2014, 66, 439-446.	0.9	24
112	High Radical Prostatectomy Surgical Volume is Related to Lower Radical Prostatectomy Total Hospital Charges. European Urology, 2006, 50, 58-63.	0.9	23
113	Zonal Origin of Localized Prostate Cancer Does not Affect the Rate of Biochemical Recurrence after Radical Prostatectomy. European Urology, 2007, 51, 949-955.	0.9	23
114	Molecular Cancer Phenotype in Normal Prostate Tissue. European Urology, 2009, 55, 885-891.	0.9	23
115	Thermal Ablation of Renal Tumors. Deutsches A&#x0308;rztblatt International, 2015, 112, 412-8.	0.6	23
116	Wound dehiscence in a sample of 1Â776 cystectomies: identification of predictors and implications for outcomes. BJU International, 2016, 117, E95-E101.	1.3	23
117	The Impact of Resident Involvement in Male One-stage Anterior Urethroplasties. Urology, 2015, 85, 937-941.	0.5	21
118	Populationâ€Based External Validation of the Updated 2012 Partin Tables in Contemporary North American Prostate Cancer Patients. Prostate, 2017, 77, 105-113.	1.2	21
119	Body mass index does not improve the ability to predict biochemical recurrence after radical prostatectomy. European Journal of Cancer, 2007, 43, 375-382.	1.3	20
120	Outcomes Research: A Methodologic Review. European Urology, 2006, 50, 218-224.	0.9	19
121	Distribution of prostate specific antigen (PSA) and percentage free PSA in a contemporary screening cohort with no evidence of prostate cancer. BJU International, 2007, 100, 37-41.	1.3	19
122	Prostate-Specific Antigen Improves the Ability of Clinical Stage and Biopsy Gleason Sum to Predict the Pathologic Stage at Radical Prostatectomy in the New Millennium. European Urology, 2007, 52, 1067-1075.	0.9	19
123	The presence of prostate cancer on saturation biopsy can be accurately predicted. BJU International, 2010, 105, 636-641.	1.3	19
124	Higher perioperative morbidity and inâ€hospital mortality in patients with endâ€stage renal disease undergoing nephrectomy for nonâ€metastatic kidney cancer: a populationâ€based analysis. BJU International, 2012, 110, E183-90.	1.3	19
125	Short-Term Outcome and Morbidity of Different Contemporary Urethroplasty Techniquesâ€A Preliminary Comparison. Journal of Endourology, 2013, 27, 925-929.	1.1	19
126	Risk-Adjusted Hazard Rates of Biochemical Recurrence for Prostate Cancer Patients after Radical Prostatectomy. European Urology, 2009, 55, 412-421.	0.9	18



#	ARTICLE	IF	CITATIONS
127	Predictive Value of Prostate-specific Antigen Expression in Prostate Cancer: A Tissue Microarray Study. <i>Urology</i> , 2009, 74, 1169-1173.	0.5	18
128	Adherence to pelvic lymph node dissection recommendations according to the National Comprehensive Cancer Network pelvic lymph node dissection guideline and the D'Amico lymph node invasion risk stratification. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2018, 36, 81.e17-81.e24.	0.8	18
129	Inflammatory prognostic markers in clear cell renal cell carcinoma "preoperative <sc>C</sc>" reactive protein does not improve predictive accuracy. <i>BJU International</i> , 2012, 110, E771-7.	1.3	17
130	Extended Pelvic Lymph Node Dissection Does Not Affect Erectile Function Recovery in Patients Treated with Bilateral Nerve-sparing Radical Prostatectomy. <i>Journal of Sexual Medicine</i> , 2012, 9, 2187-2194.	0.3	17
131	Development and internal validation of preoperative transition zone prostate cancer nomogram. <i>Urology</i> , 2006, 68, 1295-1300.	0.5	16
132	Health-insurance status is a determinant of the stage at presentation and of cancer control in European men treated with radical prostatectomy for clinically localized prostate cancer. <i>BJU International</i> , 2007, 99, 1404-1408.	1.3	16
133	Effect of autologous blood transfusion on the rate of biochemical recurrence after radical prostatectomy. <i>BJU International</i> , 2007, 100, 1249-1253.	1.3	16
134	Prediction of Pathological Stage is Inaccurate in Men with PSA Values above 20ng/mL. <i>European Urology</i> , 2007, 52, 1374-1380.	0.9	16
135	Does increasing the nodal yield improve outcomes in patients without nodal metastasis at radical cystectomy?. <i>World Journal of Urology</i> , 2012, 30, 807-814.	1.2	16
136	Holmium Laser Enucleation of the Prostate Is Safe in Patients with Prostate Cancer and Lower Urinary Tract Symptoms" A Retrospective Feasibility Study. <i>Journal of Endourology</i> , 2014, 28, 335-341.	1.1	16
137	Older patients suffer from adverse histopathological features after radical cystectomy. <i>International Journal of Urology</i> , 2011, 18, 576-584.	0.5	15
138	Loss of SPINK1 expression is associated with unfavorable outcomes in urothelial carcinoma of the bladder after radical cystectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013, 31, 1716-1724.	0.8	15
139	North American Population-Based Validation of the National Comprehensive Cancer Network Practice Guideline Recommendation of Pelvic Lymphadenectomy in Contemporary Prostate Cancer. <i>Prostate</i> , 2017, 77, 542-548.	1.2	15
140	256"MDCT" for evaluation of urolithiasis: Iterative reconstruction allows for a significant reduction of the applied radiation dose while maintaining high subjective and objective image quality. <i>Journal of Medical Imaging and Radiation Oncology</i> , 2014, 58, 283-290.	0.9	14
141	Prognostic value of alpha-methyl CoA racemase (AMACR) expression in renal cell carcinoma. <i>World Journal of Urology</i> , 2013, 31, 847-853.	1.2	13
142	Lymph node dissection during radical cystectomy for bladder cancer treatment: considerations on relevance and extent. <i>International Urology and Nephrology</i> , 2013, 45, 1561-1567.	0.6	13
143	Concomitant Seminal Vesicle Invasion in pT4a Urothelial Carcinoma of the Bladder with Contiguous Prostatic Infiltration is an Adverse Prognosticator for Cancer-Specific Survival after Radical Cystectomy. <i>Annals of Surgical Oncology</i> , 2014, 21, 4034-4040.	0.7	13
144	Risk assessment of metastatic recurrence in patients with prostate cancer by using the Cancer of the Prostate Risk Assessment score: results from 2937 European patients. <i>BJU International</i> , 2012, 110, 1714-1720.	1.3	12

#	ARTICLE	IF	CITATIONS
145	Prostate specific-antigen distribution in asymptomatic Canadian men with no clinical evidence of prostate cancer. <i>BJU International</i> , 2006, 98, 50-53.	1.3	11
146	Reduced CD151 expression is related to advanced tumour stage in urothelial bladder cancer. <i>Pathology</i> , 2012, 44, 448-452.	0.3	11
147	Re-assessment of 30-, 60- and 90-day mortality rates in non-metastatic prostate cancer patients treated either with radical prostatectomy or radiation therapy. <i>Canadian Urological Association Journal</i> , 2014, 8, 75.	0.3	11
148	External Validation of a Preoperative Nomogram for Prediction of the Risk of Recurrence After Radical Prostatectomy. <i>International Journal of Radiation Oncology Biology Physics</i> , 2010, 77, 788-792.	0.4	9
149	The Search for Biomarkers of Metastatic Seminoma. <i>Journal of Urology</i> , 2013, 190, 1046-1051.	0.2	9
150	From Gene to Clinic: TMA-Based Clinical Validation of Molecular Markers in Prostate Cancer. <i>Methods in Molecular Biology</i> , 2010, 664, 177-189.	0.4	9
151	Does increasing the nodal yield improve outcomes in contemporary patients without nodal metastasis undergoing radical prostatectomy?. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014, 32, 47.e1-47.e8.	0.8	8
152	The role of biomarkers in the assessment of prostate cancer risk prior to prostate biopsy: Which markers matter and how should they be used?. <i>World Journal of Urology</i> , 2014, 32, 871-880.	1.2	8
153	Biopsies Performed at Tertiary Care Centers are Superior to Referral Biopsies in Predicting Pathologic Gleason Sum. <i>Journal of Endourology</i> , 2008, 22, 533-538.	1.1	7
154	A comparative assessment of active surveillance for localized prostate cancer in the community versus tertiary care referral center. <i>World Journal of Urology</i> , 2014, 32, 891-897.	1.2	7
155	Super early detailed assessment of lower urinary tract symptoms after holmium laser enucleation of the prostate (HoLEP): a prospective study. <i>World Journal of Urology</i> , 2020, 38, 3207-3217.	1.2	7
156	Reply to Juan Morote's Letter to the Editor re: Felix K. Chun, Alexandre de la Taille, Hendrik van Poppel, et al. Prostate Cancer Gene 3 (PCA3): Development and Internal Validation of a Novel Biopsy Nomogram. <i>Eur Urol</i> 2009;56:659-68. <i>European Urology</i> , 2010, 57, e2-e3.	0.9	6
157	Differences in histopathological and biochemical outcomes in patients with low Gleason score prostate cancer. <i>BJU International</i> , 2010, 105, 818-823.	1.3	6
158	Predicting the risk of lymph node invasion during radical prostatectomy using the European association of urology guideline nomogram: A validation study. <i>European Journal of Surgical Oncology</i> , 2012, 38, 624-629.	0.5	6
159	Assays for Prostate Cancer. <i>Molecular Diagnosis and Therapy</i> , 2013, 17, 1-8.	1.6	6
160	Prediction of metastatic status in non-seminomatous testicular cancer. <i>World Journal of Urology</i> , 2014, 32, 1205-1211.	1.2	6
161	A Comparative Review of Apomorphine Formulations for Erectile Dysfunction. <i>Drugs and Aging</i> , 2006, 23, 309-319.	1.3	5
162	The development of nomograms for stratification of men at risk of prostate cancer prior to prostate biopsy. <i>Biomarkers in Medicine</i> , 2013, 7, 843-850.	0.6	5

#	ARTICLE	IF	CITATIONS
163	Reduced membranous MET expression is linked to bladder cancer progression. <i>Cancer Genetics</i> , 2014, 207, 147-152.	0.2	5
164	Prediction of the Risk of Harboring Prostate Cancer by a Prebiopsy Nomogram Based on Extended Biopsy Protocol. <i>Urologia Internationalis</i> , 2013, 90, 306-311.	0.6	4
165	Prognostic impact of infiltration of the vagina and/or uterus in women undergoing anterior pelvic exenteration for urothelial carcinoma of the bladder: results of a contemporary multicentre series. <i>World Journal of Urology</i> , 2015, 33, 343-350.	1.2	4
166	Prediction of Complications in Radical Prostatectomy Prostate Cancer Patients: Simulated Annealing versus Co-Morbidity Indexes. <i>Urologia Internationalis</i> , 2019, 102, 51-59.	0.6	4
167	Reply to Carsten Stephan et al's Letter to the Editor re: Felix K.-H. Chun, Markus Graefen, Alberto Briganti, Andrea Gallina, Julia Hopp, Michael W. Kattan, Hartwig Huland and Pierre I. Karakiewicz. Initial Biopsy Outcome Prediction—Head-to-Head Comparison of a Logistic Regression-Based Nomogram versus Artificial Neural Network. <i>Eur Urol</i> 2007;51:1236–43. <i>European Urology</i> , 2007, 51, 1448.	0.9	2
168	Reply. <i>BJU International</i> , 2013, 111, E20-1.	1.3	2
169	Prostate imaging—the future is now: current concepts and future potentials. <i>World Journal of Urology</i> , 2014, 32, 843-845.	1.2	2
170	Performance and Impact of Prostate Specific Membrane Antigen-Based Diagnostics in the Management of Men with Biochemical Recurrence of Prostate Cancer and its Role in Salvage Lymph Node Dissection. <i>World Journal of Men's Health</i> , 2020, 38, 32.	1.7	2
171	[18F]FLUOROCHOLINE COMBINED IN-LINE PET-CT SCAN FOR DETECTION OF LYMPH-NODE METASTASIS PRIOR TO RADICAL PROSTATECTOMY: RESULTS FROM A PROSPECTIVE HISTOLOGY BASED STUDY. <i>Journal of Urology</i> , 2008, 179, 49-49.	0.2	1
172	Does Low-Risk Prostate Cancer Detection Change With Repeat Biopsies?. <i>European Urology</i> , 2012, 61, 230-231.	0.9	1
173	Reply to Ian Beckley and Masood A. Khan's Letter to the Editor re: Felix K.-H. Chun, Thomas Steuber, Andreas Erbersdobler, et al. Development and Internal Validation of a Nomogram Predicting the Probability of Prostate Cancer Gleason Sum Upgrading Between Biopsy and Radical Prostatectomy Pathology. <i>Eur Urol</i> 2006;49:820–26. <i>European Urology</i> , 2007, 52, 1271.	0.9	0
174	Editorial Comment on: Preliminary Results of a Novel Method to Estimate the Probability of Prostate Cancer in Men with Elevated Serum PSA Values. <i>European Urology</i> , 2008, 54, 702.	0.9	0
175	NERVE-SPARING RADICAL PROSTATECTOMY DOES NOT UNDERMINE THE RATE OF BIOCHEMICAL RECURRENCE IN CAREFULLY SELECTED PATIENTS WITH PATHOLOGICALLY CONFIRMED EXTRACAPSULAR EXTENSION. <i>Journal of Urology</i> , 2008, 179, 646-647.	0.2	0
176	LONG TERM OUTCOME OF PATIENTS WITH POSITIVE LYMPH NODES DURING RADICAL PROSTATECTOMY: SURVIVAL BENEFIT OF PATIENTS WITH COMPLETED VS. ABANDONED SURGERY. <i>Journal of Urology</i> , 2008, 179, 252-252.	0.2	0
177	In Reply. <i>Deutsches Arzteblatt International</i> , 2015, 112, 758.	0.6	0