

# Bogdan Kolarz

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1675410/publications.pdf>

Version: 2024-02-01

29  
papers

557  
citations

840119

11  
h-index

676716

22  
g-index

29  
all docs

29  
docs citations

29  
times ranked

869  
citing authors

#	ARTICLE	IF	CITATIONS
1	The lack of association between PADI4_94 or PADI4_104 polymorphisms and RF, ACPA and anti-PAD4 in patients with rheumatoid arthritis. <i>Scientific Reports</i> , 2022, 12, .	1.6	3
2	Insights of rheumatoid arthritis biomarkers. <i>Biomarkers</i> , 2021, 26, 185-195.	0.9	14
3	The value of anti-CarP and anti-PAD4 as markers of rheumatoid arthritis in ACPA/RF negative rheumatoid arthritis patients. <i>Therapeutic Advances in Musculoskeletal Disease</i> , 2021, 13, 1759720X2198986.	1.2	14
4	Plasma micro-RNA-22 is associated with disease activity in well-established rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2021, , .	0.4	0
5	Hypermethylation of the miR-155 gene in the whole blood and decreased plasma level of miR-155 in rheumatoid arthritis. <i>PLoS ONE</i> , 2020, 15, e0233897.	1.1	12
6	Peptidyl Arginine Deiminase Type 4 Gene Promoter Hypo-Methylation in Rheumatoid Arthritis. <i>Journal of Clinical Medicine</i> , 2020, 9, 2049.	1.0	10
7	Methylation Pattern of the SOCS3 and IL6R Promoters in Rheumatoid Arthritis. <i>International Journal of Inflammation</i> , 2020, 2020, 1-7.	0.9	0
8	THU0139â€¦IS ADDING ANTI-CARP OR ANTI-PADI4 BENEFICIAL FOR DIAGNOSIS OF RHEUMATOID ARTHRITIS?. , 2019, , .		0
9	IRF5 promoter methylation as a new potential marker of rheumatoid arthritis. <i>Polish Archives of Internal Medicine</i> , 2019, 129, 370-376.	0.3	5
10	Epigenetic aspects of rheumatoid arthritis: contribution of non-coding RNAs. <i>Seminars in Arthritis and Rheumatism</i> , 2017, 46, 724-731.	1.6	28
11	The Role of Interleukin-17, Interleukin-23, and Transforming Growth Factor- $\beta$ in Pregnancy Complicated by Placental Insufficiency. <i>BioMed Research International</i> , 2017, 2017, 1-5.	0.9	41
12	Epigenetic determinants in rheumatoid arthritis: the influence of DNA methylation and histone modifications. <i>Postepy Higieny I Medycyny Doswiadczonej</i> , 2017, 71, 0-0.	0.1	0
13	A Prevention of Pre-eclampsia with the Use of Acetylsalicylic Acid and Low-molecular Weight Heparin â€“ Molecular Mechanisms. <i>Current Pharmaceutical Biotechnology</i> , 2016, 17, 624-628.	0.9	8
14	T CD3+CD8+Lymphocytes Are More Susceptible for Apoptosis in the First Trimester of Normal Human Pregnancy. <i>Journal of Immunology Research</i> , 2014, 2014, 1-9.	0.9	4
15	Antiphospholipid antibodies during 6-month treatment with infliximab: A preliminary report. <i>Medical Science Monitor</i> , 2014, 20, 1227-1231.	0.5	7
16	The expression of B7-H1 and B7-H4 co-stimulatory molecules on myeloid and plasmacytoid dendritic cells in pre-eclampsia and normal pregnancy. <i>Journal of Reproductive Immunology</i> , 2013, 99, 33-38.	0.8	22
17	The Expressions of Co-stimulatory Molecules are Altered on Putative Antigen-presenting Cells in Cord Blood. <i>American Journal of Reproductive Immunology</i> , 2013, 69, 180-187.	1.2	4
18	Apoptosis Signaling Is Altered in CD4+CD25+FoxP3+ T Regulatory Lymphocytes in Pre-Eclampsia. <i>International Journal of Molecular Sciences</i> , 2012, 13, 6548-6560.	1.8	26

#	ARTICLE	IF	CITATIONS
19	OS007. The apoptosis markers are altered in CD4+CD25+FOXP3+ T regulatory lymphocytes in pre-eclampsia. <i>Pregnancy Hypertension</i> , 2012, 2, 178.	0.6	0
20	PP069. The expressions of B7-H1 and B7-H4 co-stimulatory molecules on myeloid and lymphoid dendritic cells in pre-eclampsia and normal pregnancy. The expressions of B7-H1 and B7-H4 co-stimulatory molecules on myeloid and lymphoid dendritic cells in pre-eclampsia and normal pregnancy. <i>Pregnancy Hypertension</i> , 2012, 2, 278-279.	0.6	1
21	The Expressions of <sc>CD</sc>200 and <sc>CD</sc>200<sc>R</sc> Molecules on Myeloid and Lymphoid Dendritic Cells in Pre-Eclampsia and Normal Pregnancy. <i>American Journal of Reproductive Immunology</i> , 2012, 67, 474-481.	1.2	17
22	The predominance of Th17 lymphocytes and decreased number and function of Treg cells in preeclampsia. <i>Journal of Reproductive Immunology</i> , 2012, 93, 75-81.	0.8	199
23	The concentrations of soluble HLA-G protein are elevated during mid-gestation and decreased in pre-eclampsia. <i>Folia Histochemica Et Cytobiologica</i> , 2012, 50, 286-291.	0.6	31
24	O3. The predominance of Th17 lymphocytes and decreased number and function of Treg cells are present in preeclampsia. <i>Pregnancy Hypertension</i> , 2011, 1, 258.	0.6	0
25	Antibodies against cyclic citrullinated peptide donâ€™t decrease after 6 months of infliximab treatment in refractory rheumatoid arthritis. <i>Rheumatology International</i> , 2011, 31, 1439-1443.	1.5	2
26	Apoptosis of HeLa and CaSki cell lines incubated with All-trans retinoid acid.. <i>Folia Histochemica Et Cytobiologica</i> , 2010, 47, 599-603.	0.6	2
27	The expression and concentration of CD40 ligand in normal pregnancy and pre-eclampsia. <i>Journal of Reproductive Immunology</i> , 2009, 79, 215-219.	0.8	12
28	Activated T Lymphocytes in Pre-Eclampsia. <i>American Journal of Reproductive Immunology</i> , 2007, 58, 39-45.	1.2	92
29	Plasma micro-RNA-22 is associated with disease activity in well-established rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 0, , .	0.4	3