

Toshihiro Hasegawa

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1677696/toshihiro-hasegawa-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

168
papers

5,011
citations

35
h-index

65
g-index

174
ext. papers

5,915
ext. citations

4.5
avg, IF

5.25
L-index

#	Paper	IF	Citations
168	Low N level increases the susceptibility of PSI to photoinhibition induced by short repetitive flashes in leaves of different rice varieties.. <i>Physiologia Plantarum</i> , 2022 , e13644	4.6	0
167	Integration of Genomics with Crop Modeling for Predicting Rice Days to Flowering: A Multi-Model Analysis. <i>Field Crops Research</i> , 2022 , 276, 108394	5.5	0
166	Short-term high nighttime temperatures pose an emerging risk to rice grain failure. <i>Agricultural and Forest Meteorology</i> , 2022 , 314, 108779	5.8	0
165	Winter nocturnal warming affects the freeze-thaw frequency, soil aggregate distribution, and the contents and decomposability of C and N in paddy fields. <i>Science of the Total Environment</i> , 2022 , 802, 149870	10.2	2
164	A global dataset for the projected impacts of climate change on four major crops.. <i>Scientific Data</i> , 2022 , 9, 58	8.2	2
163	Monitoring canopy micrometeorology in diverse climates to improve the prediction of heat-induced spikelet sterility in rice under climate change. <i>Agricultural and Forest Meteorology</i> , 2022 , 316, 108860	5.8	1
162	A strong negative trade-off between seed number and 100-seed weight stalls genetic yield gains in northern Japanese soybean cultivars in comparison with Midwestern US cultivars. <i>Field Crops Research</i> , 2022 , 283, 108539	5.5	0
161	Evaluation of crop model prediction and uncertainty using Bayesian parameter estimation and Bayesian model averaging. <i>Agricultural and Forest Meteorology</i> , 2021 , 311, 108686	5.8	4
160	Effects of elevated atmospheric CO2 concentration on growth and photosynthesis in eddo at two different air temperatures. <i>Plant Production Science</i> , 2021 , 24, 363-373	2.4	
159	Heat-Mitigation Effects of Irrigated Rice-Paddy Fields Under Changing Atmospheric Carbon Dioxide Based on a Coupled Atmosphere and Crop Energy-Balance Model. <i>Boundary-Layer Meteorology</i> , 2021 , 179, 447-476	3.4	1
158	Factors determining the occurrence of floret sterility in rice in a hot and low-wind paddy field in Jiangnan Basin, China. <i>Field Crops Research</i> , 2021 , 267, 108161	5.5	1
157	Current rice models underestimate yield losses from short-term heat stresses. <i>Global Change Biology</i> , 2021 , 27, 402-416	11.4	7
156	Effect of foliar spray of kinetin on the enhancement of rice yield by elevated CO2. <i>Journal of Agronomy and Crop Science</i> , 2021 , 207, 535-543	3.9	0
155	Revision of estimates of climate change impacts on rice yield and quality in Japan by considering the combined effects of temperature and CO2 concentration. <i>J Agricultural Meteorology</i> , 2021 , 77, 139-149	11.9	1
154	Five-year soil warming changes soil C and N dynamics in a single rice paddy field in Japan. <i>Science of the Total Environment</i> , 2021 , 756, 143845	10.2	5
153	Development of a Simplified Growth and Yield Prediction Model for Supporting the Management of Paddy Rice Fertilization for Commercial Rice Varieties in the Tohoku Region. <i>Japanese Journal of Crop Science</i> , 2021 , 90, 430-443	0.1	1
152	Predicting biomass of rice with intermediate traits: Modeling method combining crop growth models and genomic prediction models. <i>PLoS ONE</i> , 2020 , 15, e0233951	3.7	9

151	Rice quality analysis using the Crop Survey database of Japan for climate impact assessment 2020 , 20, 1-8		0
150	Atmospheric CO Concentration and N Availability Affect the Balance of the Two Photosystems in Mature Leaves of Rice Plants Grown at a Free-Air CO Enrichment Site. <i>Frontiers in Plant Science</i> , 2020 , 11, 786	6.2	2
149	Dependence of pollination and fertilization in rice (<i>Oryza sativa</i> L.) on floret height within the canopy. <i>Field Crops Research</i> , 2020 , 249, 107741	5.5	4
148	Quantifying the Feedback Between Rice Architecture, Physiology, and Microclimate Under Current and Future CO ₂ Conditions. <i>Journal of Geophysical Research G: Biogeosciences</i> , 2020 , 125, e2019JG005452	3.7	3
147	Nitrogen Aspects of the Free-Air CO ₂ Enrichment (FACE) Study for Paddy Rice Ecosystems 2020 , 331-340		1
146	Is the yield change due to warming affected by photoperiod sensitivity? Effects of the soybean E4 locus. <i>Food and Energy Security</i> , 2020 , 9, e186	4.1	3
145	Analysis of factors related to varietal differences in the yield of rice (<i>Oryza sativa</i> L.) under Free-Air CO ₂ Enrichment (FACE) conditions. <i>Plant Production Science</i> , 2020 , 23, 19-27	2.4	6
144	Evaluation of the most appropriate spatial resolution of input data for assessing the impact of climate change on rice productivity in Japan. <i>J Agricultural Meteorology</i> , 2020 , 76, 61-68	1.1	4
143	Effects of free-air CO ₂ enrichment on heat-induced sterility and pollination in rice. <i>Plant Production Science</i> , 2019 , 22, 374-381	2.4	3
142	A High-Yielding Rice Cultivar "Takanari" Shows No N Constraints on CO Fertilization. <i>Frontiers in Plant Science</i> , 2019 , 10, 361	6.2	20
141	Oxalate contents in leaves of two rice cultivars grown at a free-air CO ₂ enrichment (FACE) site. <i>Plant Production Science</i> , 2019 , 22, 407-411	2.4	9
140	High mesophyll conductance in the high-yielding rice cultivar Takanari quantified with the combined gas exchange and chlorophyll fluorescence measurements under free-air CO ₂ enrichment. <i>Plant Production Science</i> , 2019 , 22, 395-406	2.4	10
139	Yield responses to elevated CO ₂ concentration among Japanese rice cultivars released since 1882. <i>Plant Production Science</i> , 2019 , 22, 352-366	2.4	17
138	Effects of free-air CO ₂ enrichment on flower opening time in rice. <i>Plant Production Science</i> , 2019 , 22, 367-373	2.4	5
137	Effect of long anther dehiscence on seed set at high temperatures during flowering in rice (<i>Oryza sativa</i> L.). <i>Scientific Reports</i> , 2019 , 9, 20363	4.9	8
136	Changes in grain protein and amino acids composition of wheat and rice under short-term increased [CO ₂] and temperature of canopy air in a paddy from East China. <i>New Phytologist</i> , 2019 , 222, 726-734	9.8	26
135	Stage-dependent temperature sensitivity function predicts seed-setting rates under short-term extreme heat stress in rice. <i>Agricultural and Forest Meteorology</i> , 2018 , 256-257, 196-206	5.8	18
134	Effects of Elevated Atmospheric CO ₂ on Respiratory Rates in Mature Leaves of Two Rice Cultivars Grown at a Free-Air CO ₂ Enrichment Site and Analyses of the Underlying Mechanisms. <i>Plant and Cell Physiology</i> , 2018 , 59, 637-649	4.9	8

133	Increasing canopy photosynthesis in rice can be achieved without a large increase in water use-A model based on free-air CO ₂ enrichment. <i>Global Change Biology</i> , 2018 , 24, 1321-1341	11.4	33
132	Combined drought and heat stress impact during flowering and grain filling in contrasting rice cultivars grown under field conditions. <i>Field Crops Research</i> , 2018 , 229, 66-77	5.5	35
131	Emerging research topics in agricultural meteorology and assessment of climate change adaptation. <i>J Agricultural Meteorology</i> , 2018 , 74, 54-59	1.1	8
130	Quantitative trait loci for large sink capacity enhance rice grain yield under free-air CO ₂ enrichment conditions. <i>Scientific Reports</i> , 2017 , 7, 1827	4.9	35
129	Nitrogen resorption in senescing leaf blades of rice exposed to free-air CO ₂ enrichment (FACE) under different N fertilization levels. <i>Plant and Soil</i> , 2017 , 418, 231-240	4.2	4
128	Taking account of water temperature effects on phenology improves the estimation of rice heading dates: Evidence from 758 field observations across Japan. <i>J Agricultural Meteorology</i> , 2017 , 73, 84-91	1.1	4
127	Causes of variation among rice models in yield response to CO ₂ examined with Free-Air CO ₂ Enrichment and growth chamber experiments. <i>Scientific Reports</i> , 2017 , 7, 14858	4.9	29
126	Nitrogen Distribution in Leaf Canopies of High-Yielding Rice Cultivar Takanari. <i>Crop Science</i> , 2017 , 57, 2080-2088	2.4	14
125	Large-scale evaluation of the effects of adaptation to climate change by shifting transplanting date on rice production and quality in Japan. <i>J Agricultural Meteorology</i> , 2017 , 73, 156-173	1.1	18
124	Characteristics of Atmosphere-rice Paddy Exchange of Gaseous and Particulate Reactive Nitrogen in Terms of Nitrogen Input to a Single-cropping Rice Paddy Area in Central Japan. <i>Asian Journal of Atmospheric Environment</i> , 2017 , 11, 202-216	1.3	2
123	Quantifying rice spikelet sterility in potential heat-vulnerable regions: Field surveys in Laos and southern India. <i>Field Crops Research</i> , 2016 , 190, 3-9	5.5	20
122	Effect of Elevated CO ₂ Concentration, Elevated Temperature and No Nitrogen Fertilization on Methanogenic Archaeal and Methane-Oxidizing Bacterial Community Structures in Paddy Soil. <i>Microbes and Environments</i> , 2016 , 31, 349-56	2.6	16
121	A taxonomy-based approach to shed light on the babel of mathematical models for rice simulation. <i>Environmental Modelling and Software</i> , 2016 , 85, 332-341	5.2	15
120	Inheritance analysis of anther dehiscence as a trait for the heat tolerance at flowering in japonica hybrid rice (<i>Oryza sativa</i> L.). <i>Euphytica</i> , 2016 , 211, 311-320	2.1	6
119	A meta-analysis of leaf nitrogen distribution within plant canopies. <i>Annals of Botany</i> , 2016 , 118, 239-47	4.1	45
118	Toward integration of genomic selection with crop modelling: the development of an integrated approach to predicting rice heading dates. <i>Theoretical and Applied Genetics</i> , 2016 , 129, 805-817	6	48
117	QTL mapping of dehiscence length at the basal part of thecae related to heat tolerance of rice (<i>Oryza sativa</i> L.). <i>Euphytica</i> , 2016 , 209, 715-723	2.1	7
116	Rice Free-Air Carbon Dioxide Enrichment Studies to Improve Assessment of Climate Change Effects on Rice Agriculture. <i>Advances in Agricultural Systems Modeling</i> , 2016 , 45-68	0.3	15

115	Rice grain yield and quality responses to free-air CO ₂ enrichment combined with soil and water warming. <i>Global Change Biology</i> , 2016 , 22, 1256-70	11.4	56
114	Effects of nitrogen input and climate trends on provincial rice yields in China between 1961 and 2003: quantitative evaluation using a crop model. <i>Paddy and Water Environment</i> , 2015 , 13, 529-543	1.6	5
113	Grain growth of different rice cultivars under elevated CO ₂ concentrations affects yield and quality. <i>Field Crops Research</i> , 2015 , 179, 72-80	5.5	29
112	Elevated atmospheric CO ₂ levels affect community structure of rice root-associated bacteria. <i>Frontiers in Microbiology</i> , 2015 , 6, 136	5.7	26
111	Response of soil, leaf endosphere and phyllosphere bacterial communities to elevated CO ₂ and soil temperature in a rice paddy. <i>Plant and Soil</i> , 2015 , 392, 27-44	4.2	32
110	A statistical analysis of three ensembles of crop model responses to temperature and CO ₂ concentration. <i>Agricultural and Forest Meteorology</i> , 2015 , 214-215, 483-493	5.8	25
109	Characterization of leaf blade- and leaf sheath-associated bacterial communities and assessment of their responses to environmental changes in CO ₂ temperature, and nitrogen levels under field conditions. <i>Microbes and Environments</i> , 2015 , 30, 51-62	2.6	17
108	Impacts of elevated atmospheric CO ₂ on nutrient content of important food crops. <i>Scientific Data</i> , 2015 , 2, 150036	8.2	50
107	A methodology for estimating phenological parameters of rice cultivars utilizing data from common variety trials. <i>J Agricultural Meteorology</i> , 2015 , 71, 77-89	1.1	17
106	Differential response of rice plants to high night temperatures imposed at varying developmental phases. <i>Agricultural and Forest Meteorology</i> , 2015 , 209-210, 69-77	5.8	31
105	Uncertainties in predicting rice yield by current crop models under a wide range of climatic conditions. <i>Global Change Biology</i> , 2015 , 21, 1328-41	11.4	260
104	Adaptation of rice to climate change through a cultivar-based simulation: a possible cultivar shift in eastern Japan. <i>Climate Research</i> , 2015 , 64, 275-290	1.6	14
103	Elevated temperature has stronger effects on the soil food web of a flooded paddy than does CO ₂ . <i>Soil Biology and Biochemistry</i> , 2014 , 70, 166-175	7.5	16
102	Increasing CO ₂ threatens human nutrition. <i>Nature</i> , 2014 , 510, 139-42	50.4	762
101	Elevated CO ₂ decreases the Photorespiratory NH ₃ production but does not decrease the NH ₃ compensation point in rice leaves. <i>Plant and Cell Physiology</i> , 2014 , 55, 1582-91	4.9	8
100	Free-air CO ₂ enrichment (FACE) net nitrogen fixation experiment at a paddy soil surface under submerged conditions. <i>Nutrient Cycling in Agroecosystems</i> , 2014 , 98, 57-69	3.3	6
99	Heat-tolerant rice cultivars retain grain appearance quality under free-air CO ₂ enrichment. <i>Rice</i> , 2014 , 7, 6	5.8	41
98	Isotopomer analysis of production, consumption and soil-to-atmosphere emission processes of N ₂ O at the beginning of paddy field irrigation. <i>Soil Biology and Biochemistry</i> , 2014 , 70, 66-78	7.5	29

97	Effects of elevated carbon dioxide, elevated temperature, and rice growth stage on the community structure of rice root-associated bacteria. <i>Microbes and Environments</i> , 2014 , 29, 184-90	2.6	35
96	Lower-Than-Expected Floret Sterility of Rice under Extremely Hot Conditions in a Flood-Irrigated Field in New South Wales, Australia. <i>Plant Production Science</i> , 2014 , 17, 245-252	2.4	21
95	Fully automated, high-throughput instrumentation for measuring the $\delta^{13}\text{C}$ value of methane and application of the instrumentation to rice paddy samples. <i>Rapid Communications in Mass Spectrometry</i> , 2014 , 28, 2315-24	2.2	12
94	Planting geometry as a pre-screening technique for identifying CO ₂ responsive rice genotypes: a case study of panicle number. <i>Physiologia Plantarum</i> , 2014 , 152, 520-8	4.6	12
93	Soil and water warming accelerates phenology and down-regulation of leaf photosynthesis of rice plants grown under free-air CO ₂ enrichment (FACE). <i>Plant and Cell Physiology</i> , 2014 , 55, 370-80	4.9	31
92	Do the rich always become richer? Characterizing the leaf physiological response of the high-yielding rice cultivar Takanari to free-air CO ₂ enrichment. <i>Plant and Cell Physiology</i> , 2014 , 55, 381-91	4.9	40
91	Temperature Difference between Meteorological Station and Nearby Farmland Case Study for Kumagaya City in Japan. <i>Scientific Online Letters on the Atmosphere</i> , 2014 , 10, 45-49	2.1	10
90	The contribution of entrapped gas bubbles to the soil methane pool and their role in methane emission from rice paddy soil in free-air [CO ₂] enrichment and soil warming experiments. <i>Plant and Soil</i> , 2013 , 364, 131-143	4.2	32
89	Lower responsiveness of canopy evapotranspiration rate than of leaf stomatal conductance to open-air CO ₂ elevation in rice. <i>Global Change Biology</i> , 2013 , 19, 2444-53	11.4	22
88	Effects of elevated [CO ₂] on stem and root lodging among rice cultivars. <i>Science Bulletin</i> , 2013 , 58, 1787-1794	11	11
87	Vulnerability of lodging risk to elevated CO ₂ and increased soil temperature differs between rice cultivars. <i>European Journal of Agronomy</i> , 2013 , 46, 20-24	5	16
86	Canopy-scale relationships between stomatal conductance and photosynthesis in irrigated rice. <i>Global Change Biology</i> , 2013 , 19, 2209-20	11.4	32
85	Amelioration of the reactive nitrogen flux calculation by a day/night separation in weekly mean air concentration measurements. <i>Atmospheric Environment</i> , 2013 , 79, 462-471	5.3	7
84	Rice cultivar responses to elevated CO ₂ at two free-air CO ₂ enrichment (FACE) sites in Japan. <i>Functional Plant Biology</i> , 2013 , 40, 148-159	2.7	174
83	The effects of free-air CO ₂ enrichment (FACE) on carbon and nitrogen accumulation in grains of rice (<i>Oryza sativa</i> L.). <i>Journal of Experimental Botany</i> , 2013 , 64, 3179-88	7	37
82	Responses of Eighteen Rice (<i>Oryza sativa</i> L.) Cultivars to Temperature Tested Using Two Types of Growth Chambers. <i>Plant Production Science</i> , 2013 , 16, 217-225	2.4	4
81	Application of a process-based biogeochemistry model, DNDC-Rice, to a rice field under free-air CO ₂ enrichment (FACE). <i>J Agricultural Meteorology</i> , 2013 , 69, 173-190	1.1	9
80	Nitrogen Uptake by Rice (<i>Oryza sativa</i> L.) Exposed to Low Water Temperatures at Different Growth Stages. <i>Journal of Agronomy and Crop Science</i> , 2012 , 198, 145-151	3.9	14

79	Microbial community composition controls the effects of climate change on methane emission from rice paddies. <i>Environmental Microbiology Reports</i> , 2012 , 4, 648-54	3.7	16
78	MINCERnet: A global research alliance to support the fight against heat stress in rice. <i>J Agricultural Meteorology</i> , 2012 , 68, 149-157	1.1	12
77	The temporal and species dynamics of photosynthetic acclimation in flag leaves of rice (<i>Oryza sativa</i>) and wheat (<i>Triticum aestivum</i>) under elevated carbon dioxide. <i>Physiologia Plantarum</i> , 2012 , 145, 395-405	4.6	52
76	Varietal Range in Transpiration Conductance of Flowering Rice Panicle and Its Impact on Panicle Temperature. <i>Plant Production Science</i> , 2012 , 15, 258-264	2.4	19
75	Performance of the enlarged Rice-FACE system using pure CO ₂ installed in Tsukuba, Japan. <i>J Agricultural Meteorology</i> , 2012 , 68, 15-23	1.1	41
74	MINCER: A novel instrument for monitoring the micrometeorology of rice canopies. <i>J Agricultural Meteorology</i> , 2012 , 68, 135-147	1.1	14
73	Spikelet sterility of rice observed in the record hot summer of 2007 and the factors associated with its variation. <i>J Agricultural Meteorology</i> , 2011 , 67, 225-232	1.1	42
72	Integrated micrometeorology model for panicle and canopy temperature (IM2PACT) for rice heat stress studies under climate change. <i>J Agricultural Meteorology</i> , 2011 , 67, 233-247	1.1	47
71	Microbial biomass carbon and methane oxidation influenced by rice cultivars and elevated CO ₂ in a Japanese paddy soil. <i>European Journal of Soil Science</i> , 2011 , 62, 69-73	3.4	18
70	Methane and soil CO ₂ production from current-season photosynthates in a rice paddy exposed to elevated CO ₂ concentration and soil temperature. <i>Global Change Biology</i> , 2011 , 17, 3327-3337	11.4	80
69	Gene expression profiling of rice grown in free air CO ₂ enrichment (FACE) and elevated soil temperature. <i>Field Crops Research</i> , 2011 , 121, 195-199	5.5	47
68	Potential ammonia emission from flag leaves of paddy rice (<i>Oryza sativa</i> L. cv. Koshihikari). <i>Agriculture, Ecosystems and Environment</i> , 2011 , 144, 117-123	5.7	7
67	Expected changes in future agro-climatological conditions in Northeast Thailand and their differences between general circulation models. <i>Theoretical and Applied Climatology</i> , 2011 , 106, 383-403		7
66	Genetic Adjustment to Changing Climates: Rice 2011 , 298-313		3
65	Varietal Difference in the Occurrence of Milky White Kernels in Response to Assimilate Supply in Rice Plants (<i>Oryza sativa</i> L.). <i>Plant Production Science</i> , 2011 , 14, 111-117	2.4	12
64	Spatial characterization of recent hot summers in Japan with agro-climatic indices related to rice production. <i>J Agricultural Meteorology</i> , 2011 , 67, 209-224	1.1	14
63	MeteoCrop DB: an agro-meteorological database coupled with crop models for studying climate change impacts on rice in Japan. <i>J Agricultural Meteorology</i> , 2011 , 67, 297-306	1.1	32
62	Diurnal and seasonal variations in stomatal conductance of rice at elevated atmospheric CO ₂ under fully open-air conditions. <i>Plant, Cell and Environment</i> , 2010 , 33, 322-31	8.4	36

61	Effects of free-air CO ₂ enrichment (FACE) and soil warming on CH ₄ emission from a rice paddy field: impact assessment and stoichiometric evaluation. <i>Biogeosciences</i> , 2010 , 7, 2639-2653	4.6	77
60	Effects of Temperature, Solar Radiation, and Vapor-Pressure Deficit on Flower Opening Time in Rice. <i>Plant Production Science</i> , 2010 , 13, 21-28	2.4	31
59	Genetic improvements for high yield and low soil nitrogen tolerance in rice (<i>Oryza Sativa</i> L.) under a cold environment. <i>Field Crops Research</i> , 2010 , 116, 38-45	5.5	22
58	Combined effects of elevated [CO ₂] and high night temperature on carbon assimilation, nitrogen absorption, and the allocations of C and N by rice (<i>Oryza sativa</i> L.). <i>Agricultural and Forest Meteorology</i> , 2010 , 150, 1174-1181	5.8	74
57	Heat-Induced Floret Sterility of Hybrid Rice (<i>Oryza sativa</i> L.) Cultivars under Humid and Low Wind Conditions in the Field of Jiangnan Basin, China. <i>Plant Production Science</i> , 2010 , 13, 243-251	2.4	66
56	The lowland paddy weed <i>Monochoria vaginalis</i> emits N ₂ O but not CH ₄ . <i>Agriculture, Ecosystems and Environment</i> , 2010 , 137, 219-221	5.7	5
55	Traits responsible for variation in pollination and seed set among six rice cultivars grown in a miniature paddy field with free air at a hot, humid spot in China. <i>Agriculture, Ecosystems and Environment</i> , 2010 , 139, 110-115	5.7	14
54	Response of the floating aquatic fern <i>Azolla filiculoides</i> to elevated CO ₂ , temperature, and phosphorus levels. <i>Hydrobiologia</i> , 2010 , 656, 5-14	2.4	31
53	Effect of panicle removal on photosynthetic acclimation under elevated CO ₂ in rice. <i>Photosynthetica</i> , 2010 , 48, 530-536	2.2	20
52	Estimation of water saturated areas in Northeast Thailand using a large-scale water balance model. <i>J Agricultural Meteorology</i> , 2010 , 66, 91-101	1.1	4
51	Genotypic variation in rice yield enhancement by elevated CO ₂ relates to growth before heading, and not to maturity group. <i>Journal of Experimental Botany</i> , 2009 , 60, 523-32	7	95
50	Rice plant response to long term CO ₂ enrichment: Gene expression profiling. <i>Plant Science</i> , 2009 , 177, 203-210	5.3	37
49	Interactions of elevated [CO ₂] and night temperature on rice growth and yield. <i>Agricultural and Forest Meteorology</i> , 2009 , 149, 51-58	5.8	140
48	Revising a process-based biogeochemistry model (DNDC) to simulate methane emission from rice paddy fields under various residue management and fertilizer regimes. <i>Global Change Biology</i> , 2008 , 14, 382-402	11.4	113
47	Rice yield enhancement by elevated CO ₂ is reduced in cool weather. <i>Global Change Biology</i> , 2008 , 14, 276-284	11.4	42
46	Increased night temperature reduces the stimulatory effect of elevated carbon dioxide concentration on methane emission from rice paddy soil. <i>Global Change Biology</i> , 2008 , 14, 644-656	11.4	35
45	CH ₄ emission with differences in atmospheric CO ₂ enrichment and rice cultivars in a Japanese paddy soil. <i>Global Change Biology</i> , 2008 , 14, 2678-2687	11.4	41
44	Effect of Elevated [CO ₂] on Soil Bubble and CH ₄ Emission from a Rice Paddy: A Test by ¹³ C Pulse-Labeling under Free-Air CO ₂ Enrichment. <i>Geomicrobiology Journal</i> , 2008 , 25, 396-403	2.5	10

43	A model driven by crop water use and nitrogen supply for simulating changes in the regional yield of rain-fed lowland rice in Northeast Thailand. <i>Paddy and Water Environment</i> , 2008 , 6, 73-82	1.6	32
42	Characteristics of water balance in a rainfed paddy field in Northeast Thailand. <i>Paddy and Water Environment</i> , 2008 , 6, 153-157	1.6	10
41	Modeling the dependence of the crop calendar for rain-fed rice on precipitation in Northeast Thailand. <i>Paddy and Water Environment</i> , 2008 , 6, 83-90	1.6	30
40	Features of the AFFRC model for evaluating the relationship between the water cycle and rice production. <i>Paddy and Water Environment</i> , 2008 , 6, 15-23	1.6	1
39	Design of sphingomonad-detecting probes for a DNA array, and its application to investigate the behavior, distribution, and source of Rhizospherous sphingomonas and other sphingomonads inhabiting an acid sulfate soil paddock in Kalimantan, Indonesia. <i>Bioscience, Biotechnology and Biochemistry</i> , 2007 , 71, 343-51	2.1	6
38	An empirical model of soil chemical properties that regulate methane production in Japanese rice paddy soils. <i>Journal of Environmental Quality</i> , 2007 , 36, 1920-5	3.4	33
37	Modeling the Effects of Water Temperature on Rice Growth and Yield under a Cool Climate: II. Model Application. <i>Agronomy Journal</i> , 2007 , 99, 1338-1344	2.2	21
36	Soil organic carbon stocks in China and changes from 1980s to 2000s. <i>Global Change Biology</i> , 2007 , 13, 1989-2007	11.4	262
35	Lodging in rice can be alleviated by atmospheric CO ₂ enrichment. <i>Agriculture, Ecosystems and Environment</i> , 2007 , 118, 223-230	5.7	26
34	Effect of elevated atmospheric CO ₂ concentration on soil and root respiration in winter wheat by using a respiration partitioning chamber. <i>Plant and Soil</i> , 2007 , 299, 237-249	4.2	27
33	Stability of Rice Pollination in The Field Under Hot And Dry Conditions in The Riverina Region of New South Wales, Australia. <i>Plant Production Science</i> , 2007 , 10, 57-63	2.4	64
32	Modeling the Effects of Water Temperature on Rice Growth and Yield under a Cool Climate: I. Model Development. <i>Agronomy Journal</i> , 2007 , 99, 1327-1337	2.2	21
31	CROWIS: A System for Sharing and Integrating Crop and Weather Data. <i>Agricultural Information Research</i> , 2007 , 16, 124-131	0.2	1
30	Seasonal changes in temperature dependence of photosynthetic rate in rice under a free-air CO ₂ enrichment. <i>Annals of Botany</i> , 2006 , 97, 549-57	4.1	47
29	Enhancement of rice canopy carbon gain by elevated CO ₂ is sensitive to growth stage and leaf nitrogen concentration. <i>New Phytologist</i> , 2006 , 170, 321-32	9.8	71
28	CH ₄ production potential in a paddy soil exposed to atmospheric CO ₂ enrichment. <i>Soil Science and Plant Nutrition</i> , 2006 , 52, 769-773	1.6	8
27	Frequent isolation of sphingomonads from local rice varieties and other weeds grown on acid sulfate soil in South Kalimantan, Indonesia. <i>Tropics</i> , 2006 , 15, 391-395	0.9	6
26	Soil nitrogen supply and nitrogen uptake for local rice grown in unfertilized acid sulfate soil in South Kalimantan. <i>Tropics</i> , 2006 , 15, 349-354	0.9	

25	Paddy Rice Responses to Free-Air [CO ₂] Enrichment 2006 , 87-104		12
24	Phosphorus Solubilizing Microorganisms in the Rhizosphere of Local Rice Varieties Grown without Fertilizer on Acid Sulfate Soils. <i>Soil Science and Plant Nutrition</i> , 2005 , 51, 679-681	1.6	13
23	Neutral rhizoplane pH of local rice and some predominant tree species in South and Central Kalimantan: A possible strategy of plant adaptation to acidic-soil. <i>Tropics</i> , 2005 , 14, 139-147	0.9	8
22	Modeling Spikelet Sterility Induced by Low Temperature in Rice. <i>Agronomy Journal</i> , 2005 , 97, 1524-1536	2.2	45
21	Rice morphogenesis and plant architecture: measurement, specification and the reconstruction of structural development by 3D architectural modelling. <i>Annals of Botany</i> , 2005 , 95, 1131-43	4.1	121
20	Structure and Function of Rice Root System under FACE Condition. <i>J Agricultural Meteorology</i> , 2005 , 60, 961-964	1.1	3
19	Genotypic difference in root penetration ability by durum wheat (<i>Triticum turgidum</i> L. var. durum) evaluated by a pot with paraffin-Vaseline discs. <i>Plant and Soil</i> , 2004 , 262, 169-177	4.2	21
18	Responses of leaf photosynthesis and plant water status in rice to low water temperature at different growth stages. <i>Field Crops Research</i> , 2004 , 89, 71-83	5.5	45
17	Seasonal Changes in Spectral Photon Flux Measured at Sapporo, Gifu and Naha. <i>Seibutsu Kankyo Chosetsu [Environment Control in Biology]</i> , 2004 , 42, 147-154		
16	Difference between Canopy Temperature and Air Temperature as a Criterion for Drought Avoidance in Crop Genotypes under Field Conditions in Japan. <i>Japanese Journal of Crop Science</i> , 2003 , 72, 461-470	0.1	3
15	Growth and yield of potato plants grown from microtubers in fields. <i>American Journal of Potato Research</i> , 2003 , 80, 371-378	2.1	17
14	Response of growth and grain yield in paddy rice to cool water at different growth stages. <i>Field Crops Research</i> , 2002 , 73, 67-79	5.5	96
13	Stability of Phenotypic Variation of Root Length over Environmental Conditions in the Seedling Generation of Potato.. <i>Japanese Journal of Crop Science</i> , 2000 , 69, 332-336	0.1	3
12	Changes in Vertical Distribution of Leaf Nitrogen with the Growth Stage and the Influence on Dry Matter Production in Rice. <i>Plant Production Science</i> , 1999 , 2, 37-46	2.4	2
11	Comparison of Rice Yield after Various Years of Cultivation by Natural Farming. <i>Plant Production Science</i> , 1999 , 2, 58-64	2.4	7
10	Analysis of Cool-weather Damage of Rice in 1993 Based on Data of Municipal Basis in Hokkaido.. <i>Japanese Journal of Crop Science</i> , 1998 , 67, 573-580	0.1	
9	Modelling the effect of nitrogen on rice growth and development. <i>Systems Approaches for Sustainable Agricultural Development</i> , 1997 , 243-257		17
8	Leaf nitrogen, plant age and crop dry matter production in rice. <i>Field Crops Research</i> , 1996 , 47, 107-116	5.5	21

7	Rice Leaf Photosynthesis as a Function of Nitrogen Content and Crop Developmental Stage.. <i>Japanese Journal of Crop Science</i> , 1996 , 65, 553-554	0.1	4
6	Response of Spikelet Number to Plant Nitrogen Concentration and Dry Weight in Paddy Rice. <i>Agronomy Journal</i> , 1994 , 86, 673-676	2.2	51
5	A Simplified Model for Estimating Nitrogen Mineralization in Paddy Soil.. <i>Japanese Journal of Crop Science</i> , 1994 , 63, 496-501	0.1	9
4	Improvement of yielding ability in Japonica rice cultivars and its impact on regional yield increase in Kinki District, Japan. <i>Agricultural Systems</i> , 1991 , 35, 173-187	6.1	5
3	Fertilizer-derived nitrogen use of two varieties of single-crop paddy rice: a free-air carbon dioxide enrichment study using polymer-coated ¹⁵ N-labeled urea. <i>Soil Science and Plant Nutrition</i> , 1-12	1.6	1
2	Effects of free-air CO ₂ enrichment (FACE) and soil warming on CH ₄ emission from a rice paddy field: impact assessment and stoichiometric evaluation ⁴		
1	A global dataset for the projected impacts of climate change on four major crops		1