

Junying Shi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/167917/junying-shi-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

100
papers

5,604
citations

39
h-index

73
g-index

105
ext. papers

7,210
ext. citations

6.4
avg, IF

5.58
L-index

#	Paper	IF	Citations
100	A Novel Role of Pipecolic Acid Biosynthetic Pathway in Drought Tolerance through the Antioxidant System in Tomato.. <i>Antioxidants</i> , 2021 , 10,	7.1	5
99	The Glutamate Receptor Plays a Role in Defense against <i>Botrytis cinerea</i> through Electrical Signaling in Tomato. <i>Applied Sciences (Switzerland)</i> , 2021 , 11, 11217	2.6	0
98	Strigolactones positively regulate abscisic acid-dependent heat and cold tolerance in tomato. <i>Horticulture Research</i> , 2021 , 8, 237	7.7	8
97	Crosstalk between Brassinosteroid and Redox Signaling Contributes to the Activation of CBF Expression during Cold Responses in Tomato. <i>Antioxidants</i> , 2021 , 10,	7.1	7
96	The protein kinase CPK28 phosphorylates ascorbate peroxidase and enhances thermotolerance in tomato. <i>Plant Physiology</i> , 2021 , 186, 1302-1317	6.6	9
95	Brassinosteroid signaling integrates multiple pathways to release apical dominance in tomato. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021 , 118,	11.5	9
94	The phyB-dependent induction of HY5 promotes iron uptake by systemically activating FER expression. <i>EMBO Reports</i> , 2021 , 22, e51944	6.5	8
93	High CO ₂ - and pathogen-driven expression of the carbonic anhydrase CA3 confers basal immunity in tomato. <i>New Phytologist</i> , 2021 , 229, 2827-2843	9.8	5
92	Combined genomic, transcriptomic, and metabolomic analyses provide insights into chayote (<i>Sechium edule</i>) evolution and fruit development. <i>Horticulture Research</i> , 2021 , 8, 35	7.7	9
91	Ethylene response factors 15 and 16 trigger jasmonate biosynthesis in tomato during herbivore resistance. <i>Plant Physiology</i> , 2021 , 185, 1182-1197	6.6	9
90	Nitrogen forms and metabolism affect plant defence to foliar and root pathogens in tomato. <i>Plant, Cell and Environment</i> , 2021 , 44, 1596-1610	8.4	8
89	Noncoding RNAs: functional regulatory factors in tomato fruit ripening. <i>Theoretical and Applied Genetics</i> , 2020 , 133, 1753-1762	6	5
88	Brassinosteroids act as a positive regulator of NBR1-dependent selective autophagy in response to chilling stress in tomato. <i>Journal of Experimental Botany</i> , 2020 , 71, 1092-1106	7	27
87	N-decanoyl-homoserine lactone alleviates elevated CO ₂ -induced defense suppression to <i>Botrytis cinerea</i> in tomato. <i>Scientia Horticulturae</i> , 2020 , 268, 109353	4.1	2
86	Transcriptomic and genetic approaches reveal an essential role of the NAC transcription factor SINAP1 in the growth and defense response of tomato. <i>Horticulture Research</i> , 2020 , 7, 209	7.7	13
85	Brassinosteroid-mediated reactive oxygen species are essential for tapetum degradation and pollen fertility in tomato. <i>Plant Journal</i> , 2020 , 102, 931-947	6.9	22
84	An Essential Role of Mitochondrial α -Ketoglutarate Dehydrogenase E2 in the Basal Immune Response Against Bacterial Pathogens in Tomato. <i>Frontiers in Plant Science</i> , 2020 , 11, 579772	6.2	4

83	The genome and transcriptome analysis of snake gourd provide insights into its evolution and fruit development and ripening. <i>Horticulture Research</i> , 2020 , 7, 199	7.7	6
82	Systemic Root-Shoot Signaling Drives Jasmonate-Based Root Defense against Nematodes. <i>Current Biology</i> , 2019 , 29, 3430-3438.e4	6.3	43
81	Role of ethylene biosynthesis and signaling in elevated CO ₂ -induced heat stress response in tomato. <i>Planta</i> , 2019 , 250, 563-572	4.7	22
80	Brassinosteroids Act as a Positive Regulator of Photoprotection in Response to Chilling Stress. <i>Plant Physiology</i> , 2019 , 180, 2061-2076	6.6	46
79	A novel CO ₂ -responsive systemic signaling pathway controlling plant mycorrhizal symbiosis. <i>New Phytologist</i> , 2019 , 224, 106-116	9.8	20
78	Natural variation for unusual host responses and flagellin-mediated immunity against <i>Pseudomonas syringae</i> in genetically diverse tomato accessions. <i>New Phytologist</i> , 2019 , 223, 447-461	9.8	12
77	Stomatal movements are involved in elevated CO ₂ -mitigated high temperature stress in tomato. <i>Physiologia Plantarum</i> , 2019 , 165, 569-583	4.6	17
76	Strigolactones positively regulate defense against root-knot nematodes in tomato. <i>Journal of Experimental Botany</i> , 2019 , 70, 1325-1337	7	27
75	SLHY5 Integrates Temperature, Light, and Hormone Signaling to Balance Plant Growth and Cold Tolerance. <i>Plant Physiology</i> , 2019 , 179, 749-760	6.6	39
74	BZR1 Mediates Brassinosteroid-Induced Autophagy and Nitrogen Starvation in Tomato. <i>Plant Physiology</i> , 2019 , 179, 671-685	6.6	53
73	A Plant Phytosulfokine Peptide Initiates Auxin-Dependent Immunity through Cytosolic Ca Signaling in Tomato. <i>Plant Cell</i> , 2018 , 30, 652-667	11.6	72
72	Heat Shock Factor HsfA1a Is Essential for Gene-Mediated Nematode Resistance and Triggers HO ₂ Production. <i>Plant Physiology</i> , 2018 , 176, 2456-2471	6.6	30
71	Induction of systemic resistance in tomato against <i>Botrytis cinerea</i> by N-decanoyl-homoserine lactone via jasmonic acid signaling. <i>Planta</i> , 2018 , 247, 1217-1227	4.7	21
70	Tomato photorespiratory glycolate-oxidase-derived H ₂ O ₂ production contributes to basal defence against <i>Pseudomonas syringae</i> . <i>Plant, Cell and Environment</i> , 2018 , 41, 1126-1138	8.4	14
69	Brassinosteroids act as a positive regulator for resistance against root-knot nematode involving RESPIRATORY BURST OXIDASE HOMOLOG-dependent activation of MAPKs in tomato. <i>Plant, Cell and Environment</i> , 2018 , 41, 1113-1125	8.4	35
68	Light Signaling-Dependent Regulation of Photoinhibition and Photoprotection in Tomato. <i>Plant Physiology</i> , 2018 , 176, 1311-1326	6.6	52
67	Brassinosteroid-mediated apoplastic H ₂ O ₂ -glutaredoxin 12/14 cascade regulates antioxidant capacity in response to chilling in tomato. <i>Plant, Cell and Environment</i> , 2018 , 41, 1052-1064	8.4	53
66	Elevated CO ₂ Improves Photosynthesis Under High Temperature by Attenuating the Functional Limitations to Energy Fluxes, Electron Transport and Redox Homeostasis in Tomato Leaves. <i>Frontiers in Plant Science</i> , 2018 , 9, 1739	6.2	34

65	HsfA1a upregulates melatonin biosynthesis to confer cadmium tolerance in tomato plants. <i>Journal of Pineal Research</i> , 2017 , 62, e12387	10.4	130
64	Nitric oxide is involved in the oxytetracycline-induced suppression of root growth through inhibiting hydrogen peroxide accumulation in the root meristem. <i>Scientific Reports</i> , 2017 , 7, 43096	4.9	13
63	24-Epibrassinolide alleviates organic pollutants-retarded root elongation by promoting redox homeostasis and secondary metabolism in <i>Cucumis sativus</i> L. <i>Environmental Pollution</i> , 2017 , 229, 922-931	9.3	38
62	Systemic Induction of Photosynthesis via Illumination of the Shoot Apex Is Mediated Sequentially by Phytochrome B, Auxin and Hydrogen Peroxide in Tomato. <i>Plant Physiology</i> , 2016 , 172, 1259-1272	6.6	46
61	Neglecting legumes has compromised human health and sustainable food production. <i>Nature Plants</i> , 2016 , 2, 16112	11.5	344
60	Interactions between 2-Cys peroxiredoxins and ascorbate in autophagosome formation during the heat stress response in <i>Solanum lycopersicum</i> . <i>Journal of Experimental Botany</i> , 2016 , 67, 1919-33	7	20
59	Phytochrome A and B Function Antagonistically to Regulate Cold Tolerance via Abscisic Acid-Dependent Jasmonate Signaling. <i>Plant Physiology</i> , 2016 , 170, 459-71	6.6	133
58	Genome-Wide Identification and Expression Analysis of Calcium-dependent Protein Kinase in Tomato. <i>Frontiers in Plant Science</i> , 2016 , 7, 469	6.2	36
57	Unraveling Main Limiting Sites of Photosynthesis under Below- and Above-Ground Heat Stress in Cucumber and the Alleviatory Role of Luffa Rootstock. <i>Frontiers in Plant Science</i> , 2016 , 7, 746	6.2	17
56	Redox Signaling and CBF-Responsive Pathway Are Involved in Salicylic Acid-Improved Photosynthesis and Growth under Chilling Stress in Watermelon. <i>Frontiers in Plant Science</i> , 2016 , 7, 1519	6.2	35
55	Melatonin mediates selenium-induced tolerance to cadmium stress in tomato plants. <i>Journal of Pineal Research</i> , 2016 , 61, 291-302	10.4	140
54	Apoplastic H ₂ O ₂ plays a critical role in axillary bud outgrowth by altering auxin and cytokinin homeostasis in tomato plants. <i>New Phytologist</i> , 2016 , 211, 1266-78	9.8	34
53	Microarray and genetic analysis reveals that csa-miR159b plays a critical role in abscisic acid-mediated heat tolerance in grafted cucumber plants. <i>Plant, Cell and Environment</i> , 2016 , 39, 1790-804	8.4	36
52	Grafting cucumber onto luffa improves drought tolerance by increasing ABA biosynthesis and sensitivity. <i>Scientific Reports</i> , 2016 , 6, 20212	4.9	37
51	Interplay between mitogen-activated protein kinase and nitric oxide in brassinosteroid-induced pesticide metabolism in <i>Solanum lycopersicum</i> . <i>Journal of Hazardous Materials</i> , 2016 , 316, 221-31	12.8	24
50	DWARF overexpression induces alteration in phytohormone homeostasis, development, architecture and carotenoid accumulation in tomato. <i>Plant Biotechnology Journal</i> , 2016 , 14, 1021-33	11.6	50
49	Overexpression of a brassinosteroid biosynthetic gene Dwarf enhances photosynthetic capacity through activation of Calvin cycle enzymes in tomato. <i>BMC Plant Biology</i> , 2016 , 16, 33	5.3	42
48	Melatonin enhances thermotolerance by promoting cellular protein protection in tomato plants. <i>Journal of Pineal Research</i> , 2016 , 61, 457-469	10.4	146

47	Salicylic acid binding of mitochondrial alpha-ketoglutarate dehydrogenase E2 affects mitochondrial oxidative phosphorylation and electron transport chain components and plays a role in basal defense against tobacco mosaic virus in tomato. <i>New Phytologist</i> , 2015 , 205, 1296-1307	9.8	45
46	Brassinosteroids play a critical role in the regulation of pesticide metabolism in crop plants. <i>Scientific Reports</i> , 2015 , 5, 9018	4.9	73
45	NPR1-dependent salicylic acid signaling is not involved in elevated CO ₂ -induced heat stress tolerance in <i>Arabidopsis thaliana</i> . <i>Plant Signaling and Behavior</i> , 2015 , 10, e1011944	2.5	12
44	RNA-seq analysis reveals the role of red light in resistance against <i>Pseudomonas syringae</i> pv. tomato DC3000 in tomato plants. <i>BMC Genomics</i> , 2015 , 16, 120	4.5	52
43	Interplay between reactive oxygen species and hormones in the control of plant development and stress tolerance. <i>Journal of Experimental Botany</i> , 2015 , 66, 2839-56	7	401
42	Enhanced photosynthetic capacity and antioxidant potential mediate brassinosteroid-induced phenanthrene stress tolerance in tomato. <i>Environmental Pollution</i> , 2015 , 201, 58-66	9.3	29
41	Antagonism between phytohormone signalling underlies the variation in disease susceptibility of tomato plants under elevated CO ₂ . <i>Journal of Experimental Botany</i> , 2015 , 66, 1951-63	7	77
40	High atmospheric carbon dioxide-dependent alleviation of salt stress is linked to RESPIRATORY BURST OXIDASE 1 (RBOH1)-dependent H ₂ O ₂ production in tomato (<i>Solanum lycopersicum</i>). <i>Journal of Experimental Botany</i> , 2015 , 66, 7391-404	7	34
39	Application of 24-epibrassinolide decreases the susceptibility to cucumber mosaic virus in zucchini (<i>Cucurbita pepo</i> L). <i>Scientia Horticulturae</i> , 2015 , 195, 116-123	4.1	13
38	Light-induced systemic resistance in tomato plants against root-knot nematode <i>Meloidogyne incognita</i> . <i>Plant Growth Regulation</i> , 2015 , 76, 167-175	3.2	14
37	Tomato- <i>Pseudomonas syringae</i> interactions under elevated CO ₂ concentration: the role of stomata. <i>Journal of Experimental Botany</i> , 2015 , 66, 307-16	7	24
36	Guard cell hydrogen peroxide and nitric oxide mediate elevated CO ₂ -induced stomatal movement in tomato. <i>New Phytologist</i> , 2015 , 208, 342-53	9.8	63
35	Involvement of nitric oxide in the jasmonate-dependent basal defense against root-knot nematode in tomato plants. <i>Frontiers in Plant Science</i> , 2015 , 6, 193	6.2	37
34	Melatonin mitigates cadmium phytotoxicity through modulation of phytochelatin biosynthesis, vacuolar sequestration, and antioxidant potential in <i>Solanum lycopersicum</i> L. <i>Frontiers in Plant Science</i> , 2015 , 6, 601	6.2	174
33	Tomato HsfA1a plays a critical role in plant drought tolerance by activating ATG genes and inducing autophagy. <i>Autophagy</i> , 2015 , 11, 2033-2047	10.2	109
32	Carbon dioxide enrichment alleviates heat stress by improving cellular redox homeostasis through an ABA-independent process in tomato plants. <i>Plant Biology</i> , 2015 , 17, 81-9	3.7	48
31	The relationship between the plant-encoded RNA-dependent RNA polymerase 1 and alternative oxidase in tomato basal defense against Tobacco mosaic virus. <i>Planta</i> , 2015 , 241, 641-50	4.7	14
30	Hydrogen peroxide mediates abscisic acid-induced HSP70 accumulation and heat tolerance in grafted cucumber plants. <i>Plant, Cell and Environment</i> , 2014 , 37, 2768-80	8.4	95

29	Effects of <i>Fusarium oxysporum</i> on rhizosphere microbial communities of two cucumber genotypes with contrasting Fusarium wilt resistance under hydroponic condition. <i>European Journal of Plant Pathology</i> , 2014 , 140, 643-653	2.1	10
28	H ₂ O ₂ mediates the crosstalk of brassinosteroid and abscisic acid in tomato responses to heat and oxidative stresses. <i>Journal of Experimental Botany</i> , 2014 , 65, 4371-83	7	204
27	Identification of multiple salicylic acid-binding proteins using two high throughput screens. <i>Frontiers in Plant Science</i> , 2014 , 5, 777	6.2	93
26	RBOH1-dependent H ₂ O ₂ production and subsequent activation of MPK1/2 play an important role in acclimation-induced cross-tolerance in tomato. <i>Journal of Experimental Botany</i> , 2014 , 65, 595-607	7	103
25	Chloroplastic thioredoxin-f and thioredoxin-m1/4 play important roles in brassinosteroids-induced changes in CO ₂ assimilation and cellular redox homeostasis in tomato. <i>Journal of Experimental Botany</i> , 2014 , 65, 4335-47	7	23
24	Glutathione biosynthesis and regeneration play an important role in the metabolism of chlorothalonil in tomato. <i>Chemosphere</i> , 2013 , 90, 2563-70	8.4	32
23	Flexible change and cooperation between mitochondrial electron transport and cytosolic glycolysis as the basis for chilling tolerance in tomato plants. <i>Planta</i> , 2013 , 237, 589-601	4.7	16
22	Silencing of tomato RBOH1 and MPK2 abolishes brassinosteroid-induced H ₂ O ₂ generation and stress tolerance. <i>Plant, Cell and Environment</i> , 2013 , 36, 789-803	8.4	100
21	Stimulated leaf dark respiration in tomato in an elevated carbon dioxide atmosphere. <i>Scientific Reports</i> , 2013 , 3, 3433	4.9	57
20	The role of hydrogen peroxide and nitric oxide in the induction of plant-encoded RNA-dependent RNA polymerase 1 in the basal defense against Tobacco mosaic virus. <i>PLoS ONE</i> , 2013 , 8, e76090	3.7	39
19	Temperature effects on the reactive oxygen species formation and antioxidant defence in roots of two cucurbit species with contrasting root zone temperature optima. <i>Acta Physiologiae Plantarum</i> , 2012 , 34, 713-720	2.6	13
18	The reduction of reactive oxygen species formation by mitochondrial alternative respiration in tomato basal defense against TMV infection. <i>Planta</i> , 2012 , 235, 225-38	4.7	43
17	Brassinosteroid improves seed germination and early development of tomato seedling under phenanthrene stress. <i>Plant Growth Regulation</i> , 2012 , 68, 87-96	3.2	25
16	Cellular glutathione redox homeostasis plays an important role in the brassinosteroid-induced increase in CO ₂ assimilation in <i>Cucumis sativus</i> . <i>New Phytologist</i> , 2012 , 194, 932-943	9.8	106
15	Hydrogen peroxide is involved in the cold acclimation-induced chilling tolerance of tomato plants. <i>Plant Physiology and Biochemistry</i> , 2012 , 60, 141-9	5.4	106
14	Induction of systemic stress tolerance by brassinosteroid in <i>Cucumis sativus</i> . <i>New Phytologist</i> , 2011 , 191, 706-720	9.8	113
13	Systemic induction and role of mitochondrial alternative oxidase and nitric oxide in a compatible tomato-Tobacco mosaic virus interaction. <i>Molecular Plant-Microbe Interactions</i> , 2010 , 23, 39-48	3.6	75
12	Effects of aqueous root extracts and hydrophobic root exudates of cucumber (<i>Cucumis sativus</i> L.) on nuclei DNA content and expression of cell cycle-related genes in cucumber radicles. <i>Plant and Soil</i> , 2010 , 327, 455-463	4.2	32

11	Light quality affects incidence of powdery mildew, expression of defence-related genes and associated metabolism in cucumber plants. <i>European Journal of Plant Pathology</i> , 2010 , 127, 125-135	2.1	88
10	Photoinhibition-induced reduction in photosynthesis is alleviated by abscisic acid, cytokinin and brassinosteroid in detached tomato leaves. <i>Plant Growth Regulation</i> , 2010 , 60, 175-182	3.2	34
9	Microbial community responses associated with the development of <i>Fusarium oxysporum</i> f. sp. cucumerinum after 24-epibrassinolide applications to shoots and roots in cucumber. <i>European Journal of Plant Pathology</i> , 2009 , 124, 141-150	2.1	14
8	Effects of cucumber mosaic virus infection on electron transport and antioxidant system in chloroplasts and mitochondria of cucumber and tomato leaves. <i>Physiologia Plantarum</i> , 2009 , 135, 246-574.6	4.6	66
7	Effects of calcium cyanamide on soil microbial communities and <i>Fusarium oxysporum</i> f. sp. cucumerinum. <i>Chemosphere</i> , 2009 , 75, 872-7	8.4	28
6	Detached leaves of tomato differ in their photosynthetic physiological response to moderate high and low temperature stress. <i>Scientia Horticulturae</i> , 2009 , 123, 17-22	4.1	40
5	Reactive oxygen species are involved in brassinosteroid-induced stress tolerance in cucumber. <i>Plant Physiology</i> , 2009 , 150, 801-14	6.6	507
4	Putrescine enhancement of tolerance to root-zone hypoxia in <i>Cucumis sativus</i> : a role for increased nitrate reduction. <i>Functional Plant Biology</i> , 2008 , 35, 337-345	2.7	17
3	Brassinosteroids Alleviate Heat-Induced Inhibition of Photosynthesis by Increasing Carboxylation Efficiency and Enhancing Antioxidant Systems in <i>Lycopersicon esculentum</i> . <i>Journal of Plant Growth Regulation</i> , 2008 , 27, 49-57	4.7	207
2	Decreased energy synthesis is partially compensated by a switch to sucrose synthase pathway of sucrose degradation in restricted root of tomato plants. <i>Plant Physiology and Biochemistry</i> , 2008 , 46, 1040-4	5.4	13
1	The Response of Antioxidant Enzymes in Cellular Organelles in Cucumber (<i>Cucumis sativus</i> L.) Leaves to Methyl Viologen-induced Photo-oxidative Stress. <i>Plant Growth Regulation</i> , 2006 , 49, 85-93	3.2	14