

Judith Haendeler

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1680551/judith-haendeler-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

123
papers

11,755
citations

57
h-index

108
g-index

135
ext. papers

12,656
ext. citations

7.9
avg, IF

5.78
L-index

#	Paper	IF	Citations
123	Flavanol Consumption in Healthy Men Preserves Integrity of Immunological-Endothelial Barrier Cell Functions: Nutri(epi)genomic Analysis.. <i>Molecular Nutrition and Food Research</i> , 2022 , e2100991	5.9	2
122	Protective Effects of Curcumin in Cardiovascular Diseases-Impact on Oxidative Stress and Mitochondria.. <i>Cells</i> , 2022 , 11,	7.9	7
121	Aryl Hydrocarbon Receptor-Dependent and -Independent Pathways Mediate Curcumin Anti-Aging Effects.. <i>Antioxidants</i> , 2022 , 11,	7.1	1
120	Mitochondrial Telomerase Reverse Transcriptase Protects From Myocardial Ischemia/Reperfusion Injury by Improving Complex I Composition and Function. <i>Circulation</i> , 2021 , 144, 1876-1890	16.7	8
119	Telomerase as a Therapeutic Target in Cardiovascular Disease. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2021 , 41, 1047-1061	9.4	8
118	Accessing Mitochondrial Protein Import in Living Cells by Protein Microinjection. <i>Frontiers in Cell and Developmental Biology</i> , 2021 , 9, 698658	5.7	0
117	Endothelial hyaluronan synthase 3 aggravates acute colitis in an experimental model of inflammatory bowel disease. <i>Matrix Biology</i> , 2021 , 102, 20-36	11.4	1
116	Non-canonical functions of Telomerase Reverse Transcriptase - Impact on redox homeostasis. <i>Redox Biology</i> , 2020 , 34, 101543	11.3	24
115	Local Peroxynitrite Impairs Endothelial Transient Receptor Potential Vanilloid 4 Channels and Elevates Blood Pressure in Obesity. <i>Circulation</i> , 2020 , 141, 1318-1333	16.7	34
114	Molekularer Wirkmechanismus von Koffein entschlüsselt. <i>Biologie in Unserer Zeit</i> , 2019 , 49, 12-13	0.1	
113	4-Methylumbelliferone improves the thermogenic capacity of brown adipose tissue. <i>Nature Metabolism</i> , 2019 , 1, 546-559	14.6	14
112	High Concentration of Low-Density Lipoprotein Results in Disturbances in Mitochondrial Transcription and Functionality in Endothelial Cells. <i>Oxidative Medicine and Cellular Longevity</i> , 2019 , 2019, 7976382	6.7	7
111	The Aryl Hydrocarbon Receptor (AhR) in the Aging Process: Another Puzzling Role for This Highly Conserved Transcription Factor. <i>Frontiers in Physiology</i> , 2019 , 10, 1561	4.6	22
110	Induction of a senescent like phenotype and loss of gap junctional intercellular communication by carbon nanoparticle exposure of lung epithelial cells. <i>Experimental Gerontology</i> , 2019 , 117, 106-112	4.5	2
109	Interventions to slow cardiovascular aging: Dietary restriction, drugs and novel molecules. <i>Experimental Gerontology</i> , 2018 , 109, 108-118	4.5	15
108	Mouse cardiac mitochondria do not separate in subsarcolemmal and interfibrillar subpopulations. <i>Mitochondrion</i> , 2018 , 38, 1-5	4.9	9
107	Increased Protein Tyrosine Phosphatase 1B (PTP1B) Activity and Cardiac Insulin Resistance Precede Mitochondrial and Contractile Dysfunction in Pressure-Overloaded Hearts. <i>Journal of the American Heart Association</i> , 2018 , 7,	6	16

106	CDKN1B/p27 is localized in mitochondria and improves respiration-dependent processes in the cardiovascular system-New mode of action for caffeine. <i>PLoS Biology</i> , 2018 , 16, e2004408	9.7	15
105	Non-Canonical Activation of the Epidermal Growth Factor Receptor by Carbon Nanoparticles. <i>Nanomaterials</i> , 2018 , 8,	5.4	2
104	Intra- and Interorgan Communication in the Cardiovascular System: A Special View on Redox Regulation. <i>Antioxidants and Redox Signaling</i> , 2017 , 26, 613-615	8.4	2
103	Nuclear Factor (Erythroid-Derived 2)-Like 2 and Thioredoxin-1 in Atherosclerosis and Ischemia/Reperfusion Injury in the Heart. <i>Antioxidants and Redox Signaling</i> , 2017 , 26, 630-644	8.4	39
102	The Anti-Apoptotic Properties of APEX1 in the Endothelium Require the First 20 Amino Acids and Converge on Thioredoxin-1. <i>Antioxidants and Redox Signaling</i> , 2017 , 26, 616-629	8.4	4
101	The aryl hydrocarbon receptor promotes aging phenotypes across species. <i>Scientific Reports</i> , 2016 , 6, 19618	4.9	39
100	MicroRNA-15b regulates mitochondrial ROS production and the senescence-associated secretory phenotype through sirtuin 4/SIRT4. <i>Aging</i> , 2016 , 8, 484-505	5.6	82
99	Role of Telomerase in the Cardiovascular System. <i>Genes</i> , 2016 , 7,	4.2	19
98	Critical regulators of endothelial cell functions: for a change being alternative. <i>Antioxidants and Redox Signaling</i> , 2015 , 22, 1212-29	8.4	4
97	Activation of the aryl hydrocarbon receptor by the widely used Src family kinase inhibitor 4-amino-5-(4-chlorophenyl)-7-(dimethylethyl)pyrazolo[3,4-d]pyrimidine (PP2). <i>Archives of Toxicology</i> , 2015 , 89, 1329-36	5.8	15
96	The hallmarks of fibroblast ageing. <i>Mechanisms of Ageing and Development</i> , 2014 , 138, 26-44	5.6	126
95	Measurement of endothelium-dependent vasodilation in mice--brief report. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2014 , 34, 2651-7	9.4	40
94	Cellular functions of the dual-targeted catalytic subunit of telomerase, telomerase reverse transcriptase--potential role in senescence and aging. <i>Experimental Gerontology</i> , 2014 , 56, 189-93	4.5	33
93	The imbalanced redox status in senescent endothelial cells is due to dysregulated Thioredoxin-1 and NADPH oxidase 4. <i>Experimental Gerontology</i> , 2014 , 56, 45-52	4.5	17
92	Unhealthy diet and ultrafine carbon black particles induce senescence and disease associated phenotypic changes. <i>Experimental Gerontology</i> , 2013 , 48, 8-16	4.5	35
91	Redox balance in the aged endothelium. <i>Zeitschrift Fur Gerontologie Und Geriatrie</i> , 2013 , 46, 635-8	2.7	9
90	Interacting with thioredoxin-1--disease or no disease?. <i>Antioxidants and Redox Signaling</i> , 2013 , 18, 1053-62	6.4	30
89	Two isoforms of Sister-Of-Mammalian Grainyhead have opposing functions in endothelial cells and in vivo. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2013 , 33, 1639-46	9.4	12

88	Redox modification of cell signaling in the cardiovascular system. <i>Journal of Molecular and Cellular Cardiology</i> , 2012 , 52, 550-8	5.8	77
87	Carbon nanoparticles induce ceramide- and lipid raft-dependent signalling in lung epithelial cells: a target for a preventive strategy against environmentally-induced lung inflammation. <i>Particle and Fibre Toxicology</i> , 2012 , 9, 48	8.4	38
86	The transcription factor Grainyhead like 3 (GRHL3) affects endothelial cell apoptosis and migration in a NO-dependent manner. <i>Biochemical and Biophysical Research Communications</i> , 2011 , 412, 648-53	3.4	11
85	Pioglitazone activates aortic telomerase and prevents stress-induced endothelial apoptosis. <i>Atherosclerosis</i> , 2011 , 216, 23-34	3.1	48
84	Oxidative stress-induced degradation of thioredoxin-1 and apoptosis is inhibited by thioredoxin-1-actin interaction in endothelial cells. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011 , 31, 650-6	9.4	22
83	c-Src-mediated activation of Erk1/2 is a reaction of epithelial cells to carbon nanoparticle treatment and may be a target for a molecular preventive strategy. <i>Biological Chemistry</i> , 2010 , 391, 1327-32	4.5	29
82	Well-known signaling proteins exert new functions in the nucleus and mitochondria. <i>Antioxidants and Redox Signaling</i> , 2010 , 13, 551-8	8.4	5
81	Nuclear redox signaling. <i>Antioxidants and Redox Signaling</i> , 2010 , 12, 713-42	8.4	65
80	Inhibition of the p38 MAP kinase in vivo improves number and functional activity of vasculogenic cells and reduces atherosclerotic disease progression. <i>Basic Research in Cardiology</i> , 2010 , 105, 389-97	11.8	54
79	Downregulation of mitochondrial telomerase reverse transcriptase induced by H ₂ O ₂ is Src kinase dependent. <i>Experimental Gerontology</i> , 2010 , 45, 558-62	4.5	36
78	Silver ion-induced suicidal erythrocyte death. <i>Journal of Applied Toxicology</i> , 2009 , 29, 531-6	4.1	28
77	Physical exercise prevents cellular senescence in circulating leukocytes and in the vessel wall. <i>Circulation</i> , 2009 , 120, 2438-47	16.7	248
76	Mitochondrial telomerase reverse transcriptase binds to and protects mitochondrial DNA and function from damage. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2009 , 29, 929-35	9.4	216
75	Thioredoxin-1 and endothelial cell aging: role in cardiovascular diseases. <i>Antioxidants and Redox Signaling</i> , 2009 , 11, 1733-40	8.4	48
74	Downregulation of ETS rescues diabetes-induced reduction of endothelial progenitor cells. <i>PLoS ONE</i> , 2009 , 4, e4529	3.7	16
73	Wnt5a increases cardiac gene expressions of cultured human circulating progenitor cells via a PKC delta activation. <i>PLoS ONE</i> , 2009 , 4, e5765	3.7	23
72	"Shping 2" different cellular localizations - a potential new player in aging processes. <i>Aging</i> , 2009 , 1, 664-86	3.8	7
71	Changes of MMP-1 and collagen type Ialpha1 by UVA, UVB and IRA are differentially regulated by Trx-1. <i>Experimental Gerontology</i> , 2008 , 43, 633-637	4.5	70

70	The role of near infrared radiation in photoaging of the skin. <i>Experimental Gerontology</i> , 2008 , 43, 629-632.	4.5	94
69	Effects of physical exercise on myocardial telomere-regulating proteins, survival pathways, and apoptosis. <i>Journal of the American College of Cardiology</i> , 2008 , 52, 470-82	15.1	169
68	Gene trapping identifies a putative tumor suppressor and a new inducer of cell migration. <i>Biochemical and Biophysical Research Communications</i> , 2008 , 376, 748-52	3.4	18
67	Caffeine enhances endothelial repair by an AMPK-dependent mechanism. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2008 , 28, 1967-74	9.4	44
66	Telomere gap between granulocytes and lymphocytes is a determinant for hematopoietic progenitor cell impairment in patients with previous myocardial infarction. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2008 , 28, 968-74	9.4	57
65	Nuclear protein tyrosine phosphatase Shp-2 is one important negative regulator of nuclear export of telomerase reverse transcriptase. <i>Journal of Biological Chemistry</i> , 2008 , 283, 33155-61	5.4	63
64	A new kid on the block: PKD1: a promising target for antiangiogenic therapy?. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2008 , 28, 1689-90	9.4	11
63	Inhibition of suicidal erythrocyte death by nitric oxide. <i>Pflugers Archiv European Journal of Physiology</i> , 2008 , 456, 293-305	4.6	50
62	Molecular mechanisms involved in endothelial cell aging: role of telomerase reverse transcriptase. <i>Zeitschrift Fur Gerontologie Und Geriatrie</i> , 2007 , 40, 334-8	2.7	15
61	SIRT1 controls endothelial angiogenic functions during vascular growth. <i>Genes and Development</i> , 2007 , 21, 2644-58	12.6	464
60	Sphingosine-1-phosphate stimulates the functional capacity of progenitor cells by activation of the CXCR4-dependent signaling pathway via the S1P3 receptor. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007 , 27, 275-82	9.4	146
59	Epoxyeicosatrienoic acids regulate Trp channel dependent Ca ²⁺ signaling and hyperpolarization in endothelial cells. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007 , 27, 2612-8	9.4	143
58	Nuclear redox-signaling is essential for apoptosis inhibition in endothelial cells--important role for nuclear thioredoxin-1. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007 , 27, 2325-31	9.4	54
57	Nitric oxide and endothelial cell aging. <i>European Journal of Clinical Pharmacology</i> , 2006 , 62, 137-140	2.8	17
56	Effects of granulocyte colony stimulating factor on functional activities of endothelial progenitor cells in patients with chronic ischemic heart disease. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2006 , 26, 2238-43	9.4	105
55	Inseparably tied: functional and antioxidative capacity of endothelial progenitor cells. <i>Circulation Research</i> , 2006 , 98, 157-8	15.7	18
54	Thioredoxin-1 and posttranslational modifications. <i>Antioxidants and Redox Signaling</i> , 2006 , 8, 1723-8	8.4	45
53	Differentiation of circulating endothelial progenitor cells to a cardiomyogenic phenotype depends on E-cadherin. <i>FEBS Letters</i> , 2005 , 579, 6060-6	3.8	34

52	Protective role of thioredoxin-1 in cardiovascular systems. <i>Signal Transduction</i> , 2005 , 5, 314-321		2
51	Cell-to-cell connection of endothelial progenitor cells with cardiac myocytes by nanotubes: a novel mechanism for cell fate changes?. <i>Circulation Research</i> , 2005 , 96, 1039-41	15.7	246
50	p38 mitogen-activated protein kinase downregulates endothelial progenitor cells. <i>Circulation</i> , 2005 , 111, 1184-91	16.7	186
49	The role of junctional adhesion molecule-C (JAM-C) in oxidized LDL-mediated leukocyte recruitment. <i>FASEB Journal</i> , 2005 , 19, 2078-80	0.9	75
48	Cathepsin D and H ₂ O ₂ stimulate degradation of thioredoxin-1: implication for endothelial cell apoptosis. <i>Journal of Biological Chemistry</i> , 2005 , 280, 42945-51	5.4	61
47	Non-canonical Wnt signaling enhances differentiation of human circulating progenitor cells to cardiomyogenic cells. <i>Journal of Biological Chemistry</i> , 2005 , 280, 16838-42	5.4	109
46	Impaired CXCR4 signaling contributes to the reduced neovascularization capacity of endothelial progenitor cells from patients with coronary artery disease. <i>Circulation Research</i> , 2005 , 97, 1142-51	15.7	278
45	Antioxidant effects of statins via S-nitrosylation and activation of thioredoxin in endothelial cells: a novel vasculoprotective function of statins. <i>Circulation</i> , 2004 , 110, 856-61	16.7	179
44	GIT1 mediates thrombin signaling in endothelial cells: role in turnover of RhoA-type focal adhesions. <i>Circulation Research</i> , 2004 , 94, 1041-9	15.7	60
43	Statins enhance migratory capacity by upregulation of the telomere repeat-binding factor TRF2 in endothelial progenitor cells. <i>Circulation</i> , 2004 , 110, 3136-42	16.7	197
42	GIT1 functions as a scaffold for MEK1-extracellular signal-regulated kinase 1 and 2 activation by angiotensin II and epidermal growth factor. <i>Molecular and Cellular Biology</i> , 2004 , 24, 875-85	4.8	77
41	p21 ^{Cip1} levels differentially regulate turnover of mature endothelial cells, endothelial progenitor cells, and in vivo neovascularization. <i>Circulation Research</i> , 2004 , 94, 686-92	15.7	31
40	Antioxidants inhibit nuclear export of telomerase reverse transcriptase and delay replicative senescence of endothelial cells. <i>Circulation Research</i> , 2004 , 94, 768-75	15.7	296
39	Low doses of reactive oxygen species protect endothelial cells from apoptosis by increasing thioredoxin-1 expression. <i>FEBS Letters</i> , 2004 , 577, 427-33	3.8	73
38	Gas6 inhibits apoptosis in vascular smooth muscle: role of Axl kinase and Akt. <i>Journal of Molecular and Cellular Cardiology</i> , 2004 , 37, 881-7	5.8	96
37	GIT1 mediates Src-dependent activation of phospholipase C γ by angiotensin II and epidermal growth factor. <i>Journal of Biological Chemistry</i> , 2003 , 278, 49936-44	5.4	75
36	Regulation of telomerase activity and anti-apoptotic function by protein-protein interaction and phosphorylation. <i>FEBS Letters</i> , 2003 , 536, 180-6	3.8	115
35	Shear stress increases the amount of S-nitrosylated molecules in endothelial cells: important role for signal transduction. <i>FEBS Letters</i> , 2003 , 551, 153-8	3.8	51

34	Hydrogen peroxide triggers nuclear export of telomerase reverse transcriptase via Src kinase family-dependent phosphorylation of tyrosine 707. <i>Molecular and Cellular Biology</i> , 2003 , 23, 4598-610	4.8	188
33	HMG-CoA reductase inhibitors reduce senescence and increase proliferation of endothelial progenitor cells via regulation of cell cycle regulatory genes. <i>Circulation Research</i> , 2003 , 92, 1049-55	15.7	345
32	Thioredoxin: a key regulator of cardiovascular homeostasis. <i>Circulation Research</i> , 2003 , 93, 1029-33	15.7	195
31	The vascular NADPH oxidase subunit p47phox is involved in redox-mediated gene expression. <i>Free Radical Biology and Medicine</i> , 2002 , 32, 1116-22	7.8	83
30	Angiotensin II-induced upregulation of MAP kinase phosphatase-3 mRNA levels mediates endothelial cell apoptosis. <i>Basic Research in Cardiology</i> , 2002 , 97, 1-8	11.8	26
29	Redox regulatory and anti-apoptotic functions of thioredoxin depend on S-nitrosylation at cysteine 69. <i>Nature Cell Biology</i> , 2002 , 4, 743-9	23.4	341
28	Dephosphorylation of endothelial nitric oxide synthase contributes to the anti-angiogenic effects of endostatin. <i>FASEB Journal</i> , 2002 , 16, 706-8	0.9	118
27	Fluid shear stress attenuates hydrogen peroxide-induced c-Jun NH2-terminal kinase activation via a glutathione reductase-mediated mechanism. <i>Circulation Research</i> , 2002 , 91, 712-8	15.7	65
26	Hypoxic induction of the hypoxia-inducible factor is mediated via the adaptor protein Shc in endothelial cells. <i>Circulation Research</i> , 2002 , 91, 38-45	15.7	43
25	Regulation of endothelial cell apoptosis in atherothrombosis. <i>Current Opinion in Lipidology</i> , 2002 , 13, 531-6	4.4	66
24	TNFalpha and oxLDL reduce protein S-nitrosylation in endothelial cells. <i>Journal of Biological Chemistry</i> , 2001 , 276, 41383-7	5.4	68
23	Aging enhances the sensitivity of endothelial cells toward apoptotic stimuli: important role of nitric oxide. <i>Circulation Research</i> , 2001 , 89, 709-15	15.7	314
22	Receptor heterodimerization: essential mechanism for platelet-derived growth factor-induced epidermal growth factor receptor transactivation. <i>Molecular and Cellular Biology</i> , 2001 , 21, 6387-94	4.8	149
21	Src and Cas mediate JNK activation but not ERK1/2 and p38 kinases by reactive oxygen species. <i>Journal of Biological Chemistry</i> , 2000 , 275, 11706-12	5.4	203
20	Inhibition of endogenous nitric oxide synthase potentiates ischemia-reperfusion-induced myocardial apoptosis via a caspase-3 dependent pathway. <i>Cardiovascular Research</i> , 2000 , 45, 671-8	9.9	56
19	Posttranslational modification of Bcl-2 facilitates its proteasome-dependent degradation: molecular characterization of the involved signaling pathway. <i>Molecular and Cellular Biology</i> , 2000 , 20, 1886-96	4.8	279
18	Cyclophilin A is a secreted growth factor induced by oxidative stress. <i>Circulation Research</i> , 2000 , 87, 789-96	15.7	331
17	Nitric oxide down-regulates MKP-3 mRNA levels: involvement in endothelial cell protection from apoptosis. <i>Journal of Biological Chemistry</i> , 2000 , 275, 25502-7	5.4	101

16	The third cytoplasmic loop of the angiotensin II type 1 receptor exerts differential effects on extracellular signal-regulated kinase (ERK1/ERK2) and apoptosis via Ras- and Rap1-dependent pathways. <i>Circulation Research</i> , 2000 , 86, 729-36	15.7	41
15	Angiotensin II induces transactivation of two different populations of the platelet-derived growth factor beta receptor. Key role for the p66 adaptor protein Shc. <i>Journal of Biological Chemistry</i> , 2000 , 275, 15926-32	5.4	136
14	Angiotensin II mediated signal transduction. Important role of tyrosine kinases. <i>Regulatory Peptides</i> , 2000 , 95, 1-7		48
13	Congestive heart failure induces endothelial cell apoptosis: protective role of carvedilol. <i>Journal of the American College of Cardiology</i> , 2000 , 36, 2081-9	15.1	114
12	Nitric oxide and apoptosis. <i>Vitamins and Hormones</i> , 1999 , 57, 49-77	2.5	37
11	Nitric oxide inhibits caspase-3 by S-nitrosation in vivo. <i>Journal of Biological Chemistry</i> , 1999 , 274, 6823-6	5.4	333
10	Dephosphorylation targets Bcl-2 for ubiquitin-dependent degradation: a link between the apoptosome and the proteasome pathway. <i>Journal of Experimental Medicine</i> , 1999 , 189, 1815-22	16.6	284
9	Cyclosporin A inhibits apoptosis of human endothelial cells by preventing release of cytochrome C from mitochondria. <i>Circulation</i> , 1998 , 98, 1153-7	16.7	144
8	Fluid shear stress stimulates phosphorylation of Akt in human endothelial cells: involvement in suppression of apoptosis. <i>Circulation Research</i> , 1998 , 83, 334-41	15.7	360
7	Suppression of apoptosis by nitric oxide via inhibition of interleukin-1beta-converting enzyme (ICE)-like and cysteine protease protein (CPP)-32-like proteases. <i>Journal of Experimental Medicine</i> , 1997 , 185, 601-7	16.6	766
6	Effects of redox-related congeners of NO on apoptosis and caspase-3 activity. <i>Nitric Oxide - Biology and Chemistry</i> , 1997 , 1, 282-93	5	89
5	Oxidized low-density lipoprotein induces apoptosis of human endothelial cells by activation of CPP32-like proteases. A mechanistic clue to the Response to injury hypothesis. <i>Circulation</i> , 1997 , 95, 1760-3	16.7	242
4	Angiotensin II induces apoptosis of human endothelial cells. Protective effect of nitric oxide. <i>Circulation Research</i> , 1997 , 81, 970-6	15.7	240
3	Shear stress inhibits apoptosis of human endothelial cells. <i>FEBS Letters</i> , 1996 , 399, 71-4	3.8	252
2	Vitamin C and E prevent lipopolysaccharide-induced apoptosis in human endothelial cells by modulation of Bcl-2 and Bax. <i>European Journal of Pharmacology</i> , 1996 , 317, 407-11	5.3	108
1	Endotoxic shock leads to apoptosis in vivo and reduces Bcl-2. <i>Shock</i> , 1996 , 6, 405-9	3.4	49