Yuan Gao

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/1682235/yuan-gao-publications-by-year.pdf

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

25 1,505 16 25 g-index

25 2,063 8.4 4.86 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
25	Degradation of antibiotic pollutants by persulfate activated with various carbon materials. <i>Chemical Engineering Journal</i> , 2022 , 429, 132387	14.7	23
24	High-yield and high-performance porous biochar produced from pyrolysis of peanut shell with low-dose ammonium polyphosphate for chloramphenicol adsorption. <i>Journal of Cleaner Production</i> , 2020 , 264, 121516	10.3	32
23	Insight into activated carbon from different kinds of chemical activating agents: A review. <i>Science of the Total Environment</i> , 2020 , 746, 141094	10.2	99
22	Evaluation of pyrolysis residue of oil sludge for recycling as bed material. <i>Canadian Journal of Chemical Engineering</i> , 2020 , 98, 465-474	2.3	5
21	Edge defects-enriched porous carbon derived from food waste for high-performance supercapacitors. <i>Materials Letters</i> , 2019 , 253, 74-77	3.3	8
20	Synthesis of honeycomb-like hierarchical porous carbon via molten salt pyrolysis in a novel sequencing integration system for high-performance supercapacitors. <i>Microporous and Mesoporous Materials</i> , 2019 , 278, 195-205	5.3	13
19	Self-activation of biochar from furfural residues by recycled pyrolysis gas. <i>Waste Management</i> , 2018 , 77, 312-321	8.6	24
18	Facile one-step synthesis of functionalized biochar from sustainable prolifera-green-tide source for enhanced adsorption of copper ions. <i>Journal of Environmental Sciences</i> , 2018 , 73, 185-194	6.4	14
17	Chlorination of bisphenol S: Kinetics, products, and effect of humic acid. Water Research, 2018, 131, 208	8-21.7	39
16	Facile synthesis of hierarchical porous carbon material by potassium tartrate activation for chloramphenicol removal. <i>Journal of the Taiwan Institute of Chemical Engineers</i> , 2018 , 85, 141-148	5.3	13
15	Transformation of phenolic compounds by peroxymonosulfate in the presence of iodide and formation of iodinated aromatic products. <i>Chemical Engineering Journal</i> , 2018 , 335, 855-864	14.7	31
14	Facile synthesis of high-surface area mesoporous biochar for energy storage via in-situ template strategy. <i>Materials Letters</i> , 2018 , 230, 183-186	3.3	20
13	Application for oxytetracycline wastewater pretreatment by Fe-C-Ni catalytic cathodic-anodic-electrolysis granular fillers from rare-earth tailings. <i>Ecotoxicology and Environmental Safety</i> , 2018 , 164, 641-647	7	4
12	Facile synthesis of nano ZnO/ZnS modified biochar by directly pyrolyzing of zinc contaminated corn stover for Pb(II), Cu(II) and Cr(VI) removals. <i>Waste Management</i> , 2018 , 79, 625-637	8.6	63
11	Enhanced degradation of ciprofloxacin by graphitized mesoporous carbon (GMC)-TiO nanocomposite: Strong synergy of adsorption-photocatalysis and antibiotics degradation mechanism. <i>Journal of Colloid and Interface Science</i> , 2018 , 527, 202-213	9.3	104
10	Kinetics of Oxidation of Iodide (IDand Hypoiodous Acid (HOI) by Peroxymonosulfate (PMS) and Formation of Iodinated Products in the PMS/IDNOM System. <i>Environmental Science and Technology Letters</i> , 2017 , 4, 76-82	11	49
9	Activation of peroxymonosulfate by phenols: Important role of quinone intermediates and involvement of singlet oxygen. <i>Water Research</i> , 2017 , 125, 209-218	12.5	152

LIST OF PUBLICATIONS

8	Iodine Atom or Hypoiodous Acid? Comment on "Rapid Selective Circumneutral Degradation of Phenolic Pollutants Using Peroxymonosulfate-Iodide Metal-Free Oxidation: Role of Iodine Atoms". <i>Environmental Science & Discounty</i> , 2017, 51, 9410-9411	10.3	7
7	Preparation of well-developed mesoporous activated carbon with high yield by ammonium polyphosphate activation. <i>Journal of the Taiwan Institute of Chemical Engineers</i> , 2016 , 66, 394-399	5.3	15
6	Transformation of Flame Retardant Tetrabromobisphenol A by Aqueous Chlorine and the Effect of Humic Acid. <i>Environmental Science & Environmental Scien</i>	10.3	52
5	Activation of Peroxymonosulfate by Benzoquinone: A Novel Nonradical Oxidation Process. <i>Environmental Science & Environmental </i>	10.3	602
4	Activated carbons with well-developed mesoporosity prepared by activation with different alkali salts. <i>Materials Letters</i> , 2015 , 146, 34-36	3.3	14
3	Optimization of high surface area activated carbon production from Enteromorpha prolifra with low-dose activating agent. <i>Fuel Processing Technology</i> , 2015 , 132, 180-187	7.2	20
2	Preparation of highly developed mesoporous activated carbon by H4P2O7 activation and its adsorption behavior for oxytetracycline. <i>Powder Technology</i> , 2013 , 249, 54-62	5.2	37
1	Comparisons of porous, surface chemistry and adsorption properties of carbon derived from Enteromorpha prolifera activated by H4P2O7 and KOH. <i>Chemical Engineering Journal</i> , 2013 , 232, 582-59	9 ó 4·7	65