Toms Zima

List of Publications by Citations

Source: https://exaly.com/author-pdf/1685976/tomas-zima-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

146 papers

3,163 citations

31 h-index

4/ g-index

165 ext. papers

3,468 ext. citations

3.5 avg, IF

4.75 L-index

#	Paper	IF	Citations
146	Advanced glycoxidation end products in chronic diseases-clinical chemistry and genetic background. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2005 , 579, 37-	4 6 3	142
145	Soluble receptor for advanced glycation end products in patients with decreased renal function. <i>American Journal of Kidney Diseases</i> , 2006 , 47, 406-11	7.4	128
144	Oxidative stress and signal transduction pathways in alcoholic liver disease. <i>Alcoholism: Clinical and Experimental Research</i> , 2005 , 29, 110S-115S	3.7	98
143	On-line haemodiafiltration versus haemodialysis: stable haematocrit with less erythropoietin and improvement of other relevant blood parameters. <i>Blood Purification</i> , 2006 , 24, 163-73	3.1	89
142	Receptor for advanced glycation end products (RAGE)soluble form (sRAGE) and gene polymorphisms in patients with breast cancer. <i>Cancer Investigation</i> , 2007 , 25, 720-5	2.1	82
141	Depression, traumatic stress and interleukin-6. <i>Journal of Affective Disorders</i> , 2010 , 120, 231-4	6.6	80
140	The effect of quinolinate on rat brain lipid peroxidation is dependent on iron. <i>Neurochemistry International</i> , 1997 , 30, 233-237	4.4	79
139	Significantly higher procalcitonin levels could differentiate Gram-negative sepsis from Gram-positive and fungal sepsis. <i>Clinical and Experimental Medicine</i> , 2013 , 13, 165-70	4.9	76
138	Reference intervals in evaluation of maternal thyroid function during the first trimester of pregnancy. <i>European Journal of Endocrinology</i> , 2009 , 160, 791-7	6.5	62
137	Receptor for advanced glycation end productssoluble form and gene polymorphisms in chronic haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2007 , 22, 2020-6	4.3	62
136	Randomized prospective comparative study of ursodeoxycholic acid and S-adenosyl-L-methionine in the treatment of intrahepatic cholestasis of pregnancy. <i>Journal of Perinatal Medicine</i> , 2006 , 34, 383-9	1^{2.7}	61
135	Thyroid in pregnancy: From physiology to screening. <i>Critical Reviews in Clinical Laboratory Sciences</i> , 2017 , 54, 102-116	9.4	56
134	Human galectins induce conversion of dermal fibroblasts into myofibroblasts and production of extracellular matrix: potential application in tissue engineering and wound repair. <i>Cells Tissues Organs</i> , 2011 , 194, 469-80	2.1	52
133	Relationship of soluble RAGE and RAGE ligands HMGB1 and EN-RAGE to endothelial dysfunction in type 1 and type 2 diabetes mellitus. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2012 , 120, 277-81	2.3	51
132	Influence of plasma exchange on serum levels of cytokines and adhesion molecules in ANCA-positive renal vasculitis. <i>Blood Purification</i> , 1998 , 16, 72-80	3.1	51
131	Trace elements in hemodialysis and continuous ambulatory peritoneal dialysis patients. <i>Blood Purification</i> , 1998 , 16, 253-60	3.1	50
130	Oxidative stress and inflammation in pregnancy. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2006 , 66, 121-7	2	49

(2011-2002)

129	Advanced glycation end products and advanced oxidation protein products in hemodialyzed patients. <i>Blood Purification</i> , 2002 , 20, 531-6	3.1	49	
128	Qualitative and quantitative detection of urinary human complement factor H-related protein (BTA stat and BTA TRAK) and fragments of cytokeratins 8, 18 (UBC rapid and UBC IRMA) as markers for transitional cell carcinoma of the bladder. <i>European Urology</i> , 2002 , 41, 34-9	10.2	47	
127	Trace elements in end-stage renal disease. 2. Clinical implication of trace elements. <i>Blood Purification</i> , 1999 , 17, 187-98	3.1	46	
126	Marked increase of procalcitonin after the administration of anti-thymocyte globulin in patients before hematopoietic stem cell transplantation does not indicate sepsis: a prospective study. <i>Critical Care</i> , 2009 , 13, R37	10.8	45	
125	Soluble receptor for advanced glycation end-products (sRAGE) and polymorphisms of RAGE and glyoxalase I genes in patients with pancreas cancer. <i>Clinical Biochemistry</i> , 2010 , 43, 882-6	3.5	42	
124	Advanced glycation end products in clinical nephrology. <i>Kidney and Blood Pressure Research</i> , 2004 , 27, 18-28	3.1	42	
123	Glycoxidation and inflammation in chronic haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2003 , 18, 2577-81	4.3	40	
122	Performance characteristics of seven neuron-specific enolase assays. <i>Tumor Biology</i> , 2007 , 28, 84-92	2.9	36	
121	Relationship of pregnancy-associated plasma protein-a to renal function and dialysis modalities. <i>Kidney and Blood Pressure Research</i> , 2004 , 27, 88-95	3.1	36	
120	Ferritin as an independent mortality predictor in patients with pancreas cancer. Results of a pilot study. <i>Tumor Biology</i> , 2012 , 33, 1695-700	2.9	35	
119	Culture methods of glomerular podocytes. Kidney and Blood Pressure Research, 2007, 30, 162-74	3.1	35	
118	Advanced glycation end-products in patients with chronic alcohol misuse. <i>Alcohol and Alcoholism</i> , 2004 , 39, 316-20	3.5	35	
117	The relationship between the cervical and anal HPV infection in women with cervical intraepithelial neoplasia. <i>Journal of Clinical Virology</i> , 2014 , 59, 18-23	14.5	33	
116	Dynamic genetic architecture of metabolic syndrome attributes in the rat. <i>Physiological Genomics</i> , 2005 , 21, 243-52	3.6	32	
115	Oxidative stress and endothelium influenced by metformin in type 2 diabetes mellitus. <i>European Journal of Clinical Pharmacology</i> , 2007 , 63, 1107-14	2.8	31	
114	Serum trough infliximab levels: A comparison of three different immunoassays for the monitoring of CT-P13 (infliximab) treatment in patients with inflammatory bowel disease. <i>Biologicals</i> , 2016 , 44, 33-	-6 ^{1.8}	30	
113	Panel of Urinary Diagnostic Markers for Non-Invasive Detection of Primary and Recurrent Urothelial Urinary Bladder Carcinoma. <i>Urologia Internationalis</i> , 2015 , 95, 56-64	1.9	29	
112	Leptin and adiponectin in pancreatic cancer: connection with diabetes mellitus. <i>Neoplasma</i> , 2011 , 58, 58-64	3.3	28	

111	Soluble receptor for advanced glycation end products in physiological and pathological pregnancy. <i>Clinical Biochemistry</i> , 2010 , 43, 442-6	3.5	28
110	Reference intervals for thyroid markers in early pregnancy determined by 7 different analytical systems. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2014 , 74, 95-101	2	27
109	Comparison of the Suitability of Various Atomic SpectroscopicTechinques for the Determination of Selenium in Human Whole Blood. <i>Journal of Analytical Atomic Spectrometry</i> , 1997 , 12, 85-89	3.7	26
108	Vitamin D Binding Protein Is Not Involved in Vitamin D Deficiency in Patients with Chronic Kidney Disease. <i>BioMed Research International</i> , 2015 , 2015, 492365	3	25
107	Increased uric acid and glucose concentrations in vitreous and serum of patients with diabetic macular oedema. <i>Ophthalmic Research</i> , 2011 , 46, 73-9	2.9	25
106	No benefit of hemodiafiltration over hemodialysis in lowering elevated levels of asymmetric dimethylarginine in ESRD patients. <i>Blood Purification</i> , 2006 , 24, 439-44	3.1	25
105	Lower retinol levels as an independent predictor of mortality in long-term hemodialysis patients: a prospective observational cohort study. <i>American Journal of Kidney Diseases</i> , 2010 , 56, 513-21	7.4	24
104	The brighter (and evolutionarily older) face of the metabolic syndrome: evidence from Trypanosoma cruzi infection in CD-1 mice. <i>Diabetes/Metabolism Research and Reviews</i> , 2015 , 31, 346-35	97.5	23
103	Fibroblast growth factor 23 and matrix-metalloproteinases in patients with chronic kidney disease: are they associated with cardiovascular disease?. <i>Kidney and Blood Pressure Research</i> , 2009 , 32, 276-83	3.1	23
102	The quality of the extra-analytical phase of laboratory practice in some developing European countries and Mexico - a multicentric study. <i>Clinical Chemistry and Laboratory Medicine</i> , 2011 , 49, 215-28	3 ^{5.9}	23
101	Increased levels of pregnancy-associated plasma protein A are associated with mortality in hemodialysis patients: preliminary results. <i>Blood Purification</i> , 2004 , 22, 298-300	3.1	23
100	Pregnancy-associated plasma protein A (PAPP-A) and preeclampsia. <i>Advances in Clinical Chemistry</i> , 2014 , 63, 169-209	5.8	22
99	Low levels of circulating T-regulatory lymphocytes and short cervical length are associated with preterm labor. <i>Journal of Reproductive Immunology</i> , 2014 , 106, 110-7	4.2	22
98	A419C (E111A) polymorphism of the glyoxalase I gene and vascular complications in chronic hemodialysis patients. <i>Annals of the New York Academy of Sciences</i> , 2008 , 1126, 268-71	6.5	22
97	HMGB1, S100 proteins and other RAGE ligands in cancer - markers, mediators and putative therapeutic targets. <i>Biomedical Papers of the Medical Faculty of the University Palacky&#x0301;, Olomouc, Czechoslovakia</i> , 2016 , 160, 1-10	1.7	22
96	Detection of circulating tumor cells during follow-up of patients with early breast cancer: Clinical utility for monitoring of therapy efficacy. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2014 , 74, 132-42	2	21
95	Pregnancy-associated plasma protein a and soluble receptor for advanced glycation end products after kidney transplantation. <i>Kidney and Blood Pressure Research</i> , 2007 , 30, 31-7	3.1	21
94	ICRF-187 (dexrazoxan) protects from adriamycin-induced nephrotic syndrome in rats. <i>Nephrology Dialysis Transplantation</i> , 1998 , 13, 1975-9	4.3	21

(2008-2008)

93	Detection of feto-maternal infection/inflammation by the soluble receptor for advanced glycation end products (sRAGE): results of a pilot study. <i>Journal of Perinatal Medicine</i> , 2008 , 36, 399-404	2.7	20	
92	Comparison of different immunoassays for CA 19-9. <i>Clinical Chemistry and Laboratory Medicine</i> , 2001 , 39, 1278-82	5.9	20	
91	Antioxidant enzymessuperoxide dismutase and glutathione peroxidasein haemodialyzed patients. <i>Blood Purification</i> , 1996 , 14, 257-61	3.1	20	
90	Polymorphisms of the receptor for advanced glycation end-products and glyoxalase I in patients with renal cancer. <i>Tumor Biology</i> , 2015 , 36, 2121-6	2.9	19	
89	Surface plasmon resonance biosensor for detection of pregnancy associated plasma protein A2 in clinical samples. <i>Analytical and Bioanalytical Chemistry</i> , 2016 , 408, 7265-9	4.4	19	
88	The effect of silibinin on experimental cyclosporine nephrotoxicity. <i>Renal Failure</i> , 1998 , 20, 471-9	2.9	19	
87	Influence of oral vitamin E therapy on micro-inflammation and cardiovascular disease markers in chronic hemodialysis patients. <i>Renal Failure</i> , 2006 , 28, 395-9	2.9	19	
86	Impact of risk factors on prevalence of anal HPV infection in women with simultaneous cervical lesion. <i>Neoplasma</i> , 2015 , 62, 308-14	3.3	18	
85	Pregnancy-associated plasma protein A as an independent mortality predictor in long-term hemodialysis patients. <i>Kidney and Blood Pressure Research</i> , 2012 , 35, 192-201	3.1	18	
84	Study of urinary proteomes in Anderson-Fabry disease. <i>Renal Failure</i> , 2010 , 32, 1202-9	2.9	18	
83	Matrix metalloproteinases in serum and the follicular fluid of women treated by in vitro fertilization. <i>Journal of Assisted Reproduction and Genetics</i> , 2012 , 29, 1207-12	3.4	17	
82	Immunological and biochemical markers in preeclampsia. <i>Journal of Reproductive Immunology</i> , 2012 , 96, 90-4	4.2	17	
81	RAGE and its ligands in cancer - culprits, biomarkers, or therapeutic targets?. <i>Neoplasma</i> , 2015 , 62, 353-	6 4 .3	16	
80	Trefoil factor family (TFF) proteins as potential serum biomarkers in patients with metastatic colorectal cancer. <i>Neoplasma</i> , 2015 , 62, 470-7	3.3	16	
79	Galactose-deficient IgA1 and the corresponding IgG autoantibodies predict IgA nephropathy progression. <i>PLoS ONE</i> , 2019 , 14, e0212254	3.7	15	
78	Correlation of Vitreous Vascular Endothelial Growth Factor and Uric Acid Concentration Using Optical Coherence Tomography in Diabetic Macular Edema. <i>Journal of Ophthalmology</i> , 2015 , 2015, 478	569	15	
77	Genetic predisposition to advanced glycation end products toxicity is related to prognosis of chronic hemodialysis patients. <i>Kidney and Blood Pressure Research</i> , 2010 , 33, 30-6	3.1	15	
76	Recent progress in the pathogenesis of nephrotic proteinuria. <i>Critical Reviews in Clinical Laboratory Sciences</i> , 2008 , 45, 139-220	9.4	14	

75	Placental growth factor, pregnancy-associated plasma protein-A, soluble receptor for advanced glycation end products, extracellular newly identified receptor for receptor for advanced glycation end products binding protein and high mobility group box 1 levels in patients with acute kidney	2.7	13
74	High-risk human papillomavirus DNA in the primary tumor, sentinel, and nonsentinel pelvic lymph nodes in patients with early-stage cervical cancer: a correlation with histopathology. <i>International Journal of Gynecological Cancer</i> , 2009 , 19, 703-7	3.5	13
73	Serum S100A12 (EN-RAGE) levels in patients with decreased renal function and subclinical chronic inflammatory disease. <i>Kidney and Blood Pressure Research</i> , 2011 , 34, 457-64	3.1	12
72	Low maternal serum matrix metalloproteinase (MMP)-2 concentrations are associated with preterm labor and fetal inflammatory response. <i>Journal of Perinatal Medicine</i> , 2010 , 38, 589-96	2.7	12
71	Anti-inflammatory effect of biological treatment in patients with inflammatory bowel diseases: calprotectin and IL-6 changes do not correspond to sRAGE changes. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2010 , 70, 294-9	2	12
70	Glyoxalase I Glu111Ala polymorphism in patients with breast cancer. Cancer Investigation, 2009, 27, 655	- 6 0	12
69	Retinol and alpha-tocopherol in hemodialysis patients. <i>Renal Failure</i> , 1998 , 20, 505-12	2.9	12
68	Resting energy expenditure and thermal balance during isothermic and thermoneutral haemodialysisheat production does not explain increased body temperature during haemodialysis. <i>Nephrology Dialysis Transplantation</i> , 2007 , 22, 3553-60	4.3	12
67	A randomised controlled trial of roller versus centrifugal cardiopulmonary bypass pumps in patients undergoing pulmonary endarterectomy. <i>Perfusion (United Kingdom)</i> , 2015 , 30, 520-8	1.9	11
66	Diagnostic Importance of Selected Protein Serum Markers in the Primary Diagnostics of Prostate Cancer. <i>Urologia Internationalis</i> , 2015 , 95, 429-35	1.9	10
65	Pregnancy-associated plasma protein A associates with cardiovascular events in diabetic hemodialysis patients. <i>Atherosclerosis</i> , 2014 , 236, 263-9	3.1	10
64	Placental growth factor may predict increased left ventricular mass index in patients with mild to moderate chronic kidney diseasea prospective observational study. <i>BMC Nephrology</i> , 2013 , 14, 142	2.7	10
63	EN-RAGE (extracellular newly identified receptor for advanced glycation end-products binding protein) and mortality of long-term hemodialysis patients: A prospective observational cohort study. Clinical Biochemistry, 2012 , 45, 556-60	3.5	10
62	Expression of Biliverdin Reductase A in peripheral blood leukocytes is associated with treatment response in HCV-infected patients. <i>PLoS ONE</i> , 2013 , 8, e57555	3.7	10
61	Skin autofluorescence relates to soluble receptor for advanced glycation end-products and albuminuria in diabetes mellitus. <i>Journal of Diabetes Research</i> , 2013 , 2013, 650694	3.9	10
60	Impact of essential hypertension and primary aldosteronism on plasma brain natriuretic peptide concentration. <i>Blood Pressure</i> , 2006 , 15, 302-7	1.7	10
59	Oxidative stress, metabolism of ethanol and alcohol-related diseases 2001 , 8, 59		10
58	Placental growth factor in patients with decreased renal function. <i>Renal Failure</i> , 2011 , 33, 291-7	2.9	9

(2015-2003)

57	Pathobiochemistry of nephrotic syndrome. Advances in Clinical Chemistry, 2003, 37, 173-218	5.8	9
56	Effects of Doxorubicin (Adriamycin) and [(+)-1,2-bis(3,5-dioxopiperazinyl-1-yl)]propane (ICRF-187) on skeletal muscle protease activities. <i>Toxicology and Applied Pharmacology</i> , 2001 , 171, 135-40	4.6	9
55	Trace elements in end-stage renal disease. 1. Methodological aspects and the influence of water treatment and dialysis equipment. <i>Blood Purification</i> , 1999 , 17, 182-6	3.1	9
54	Early detection of pancreatic cancer: impact of high-resolution imaging methods and biomarkers. <i>European Journal of Gastroenterology and Hepatology</i> , 2016 , 28, e33-e43	2.2	9
53	Protective effects of inhaled carbon monoxide in endotoxin-induced cholestasis is dependent on its kinetics. <i>Biochimie</i> , 2014 , 97, 173-80	4.6	8
52	Association of pregnancy-associated plasma protein A polymorphism with preeclampsia - a pilot study. <i>Clinical Biochemistry</i> , 2011 , 44, 1380-4	3.5	8
51	RAGE polymorphisms, renal function and histological finding at 12 months after renal transplantation. <i>Clinical Biochemistry</i> , 2009 , 42, 347-52	3.5	8
50	Associations of serum levels of advanced glycation end products with nutrition markers and anemia in patients with chronic kidney disease. <i>Renal Failure</i> , 2011 , 33, 131-7	2.9	8
49	Does renal function influence plasma levels of advanced glycation and oxidation protein products in patients with chronic rheumatic diseases complicated by secondary amyloidosis?. <i>Kidney and Blood Pressure Research</i> , 2007 , 30, 1-7	3.1	8
48	Effect of hemodiafiltration on pregnancy-associated plasma protein A (PAPP-A) and related parameters. <i>Renal Failure</i> , 2006 , 28, 715-21	2.9	8
47	Lipid peroxidation and antioxidant enzymes in CAPD patients. Renal Failure, 1996, 18, 113-9	2.9	8
46	Receptor for advanced glycation end products (RAGE) and glyoxalase I gene polymorphisms in pathological pregnancy. <i>Clinical Biochemistry</i> , 2012 , 45, 1409-14	3.5	7
45	Pregnancy-associated plasma protein A: spotlight on kidney diseases. <i>Clinical Chemistry and Laboratory Medicine</i> , 2012 , 50, 1183-90	5.9	7
44	Plasma calprotectin in chronically dialyzed end-stage renal disease patients. <i>Inflammation Research</i> , 2010 , 59, 299-305	7.2	7
43	Quantification of trace elements in human serum fractions by liquid chromatography and inductively coupled plasma mass spectrometry. <i>Applied Organometallic Chemistry</i> , 2007 , 21, 5-14	3.1	7
42	Diagnosis and treatment of carpal tunnel syndrome. <i>Renal Failure</i> , 1993 , 15, 533-7	2.9	7
41	Relationship between serum bilirubin and uric acid to oxidative stress markers in Italian and Czech populations. <i>Journal of Applied Biomedicine</i> , 2013 , 11, 209-221	0.6	6
40	Predictive factors for the presence of tumor cells in bone marrow and peripheral blood in breast cancer patients. <i>Neoplasma</i> , 2015 , 62, 259-68	3.3	6

39	Dissociative symptoms reflect levels of tumor necrosis factor alpha in patients with unipolar depression. <i>Neuropsychiatric Disease and Treatment</i> , 2014 , 10, 675-9	3.1	6
38	Endotelial activation and flow-mediated vasodilation in young patients with breast cancer. <i>Neoplasma</i> , 2013 , 60, 690-7	3.3	6
37	Breath alcohol level and plasma amino acids: a comparison between older and younger chronic alcohol-dependent patients. <i>Alcohol and Alcoholism</i> , 2008 , 43, 653-7	3.5	6
36	PAPP-A, a novel marker of unstable plaque, is not influenced by hypolipidemic treatment in contrast to CRP. <i>Atherosclerosis</i> , 2003 , 166, 195-6	3.1	6
35	Acute doxorubicin (adriamycin) dosage does not reduce cardiac protein synthesis in vivo, but decreases diaminopeptidase I and proline endopeptidase activities. <i>Experimental and Molecular Pathology</i> , 2001 , 70, 154-61	4.4	6
34	The characterization of four gene expression analysis in circulating tumor cells made by Multiplex-PCR from the AdnaTest kit on the lab-on-a-chip Agilent DNA 1000 platform. <i>Biochemia Medica</i> , 2016 , 26, 103-13	2.5	6
33	Improvements in colorectal cancer screening programmes - quantitative immunochemical faecal occult blood testing - how to set the cut-off for a particular population. <i>Biomedical Papers of the Medical Faculty of the University Palacky&#x0301;, Olomouc, Czechoslovakia</i> , 2012 , 156, 143-50	1.7	6
32	Prolonged progesterone administration is associated with less frequent cervicovaginal colonization by Ureaplasma urealyticum during pregnancy - Results of a pilot study. <i>Journal of Reproductive Immunology</i> , 2016 , 116, 35-41	4.2	5
31	Biomarkers demonstrate increased consumption, but not abuse, of ethanol in essential tremor. Journal of Neurology, 2012 , 259, 1177-80	5.5	5
30	High-risk human papillomavirus DNA in paraaortic lymph nodes in advanced stages of cervical carcinoma. <i>Journal of Clinical Virology</i> , 2011 , 50, 46-9	14.5	5
29	Interlaboratory study of free monoclonal immunoglobulin light chain quantification. <i>Clinical Chemistry and Laboratory Medicine</i> , 2011 , 49, 89-92	5.9	5
28	Cell-free DNA is higher and more fragmented in intrahepatic cholestasis of pregnancy. <i>Prenatal Diagnosis</i> , 2016 , 36, 1156-1158	3.2	4
27	Early vancomycin, amikacin and gentamicin concentrations in pulmonary artery and pulmonary tissue are not affected by VA ECMO (venoarterial extracorporeal membrane oxygenation) in a pig model of prolonged cardiac arrest. <i>Pulmonary Pharmacology and Therapeutics</i> , 2013 , 26, 655-60	3.5	4
26	Low glucose degradation product peritoneal dialysis regimen is associated with lower plasma EN-RAGE and HMGB-1 proinflammatory ligands of receptor for advanced glycation end products. <i>Therapeutic Apheresis and Dialysis</i> , 2014 , 18, 309-16	1.9	4
25	Advanced glycation end products in myocardial reperfusion injury. <i>Heart and Vessels</i> , 2012 , 27, 208-15	2.1	4
24	Selenium: an important trace element and therapeutic adjunct in critical care. <i>Trace Elements and Electrolytes</i> , 2012 , 29, 246-255	1.8	4
23	Advanced glycation end products in hemodialyzed patients with diabetes mellitus correlate with leptin and leptin/body fat ratio. <i>Renal Failure</i> , 2003 , 25, 277-86	2.9	4
22	Antiphospholipid antibodies in patients with lupus nephritis. <i>Renal Failure</i> , 2003 , 25, 747-58	2.9	4

Modulation of Oxidative Stress by Alcohol. Alcoholism: Clinical and Experimental Research, 2005, 29, 1060, 1065 4. 21 Acute dosage with dexrazoxane, but not doxorubicin, is associated with increased rates of hepatic 20 2.1 4 protein synthesis in vivo. Toxicologic Pathology, 2001, 29, 591-9 Changes in levels of matrix metalloproteinase-2 and -9, pregnancy-associated plasma protein-A in 4.8 19 4 patients with various nephropathies. Journal of Nephrology, 2013, 26, 502-9 Diagnostic and prognostic value of presepsin in preterm deliveries. Journal of Maternal-Fetal and 18 Neonatal Medicine, 2015, 28, 1032-7 Gene Expression Analysis of Immunomagnetically Enriched Circulating Tumor Cell Fraction in 17 4.5 3 Castration-Resistant Prostate Cancer. Molecular Diagnosis and Therapy, 2018, 22, 381-390 Cys327Cys polymorphism of the PAPP-A gene (pregnancy associated plasma protein A) is related to 3.5 mortality of long term hemodialysis patients. Clinical Biochemistry, 2014, 47, 578-83 Fetuin-A early after renal transplantation. Kidney and Blood Pressure Research, 2009, 32, 217-22 15 3.1 3 Recommendations on prenatal screening and the connections to other diseases such as thyroid 14 5.9 dysfunction. Clinical Chemistry and Laboratory Medicine, 2012, 50, 1211-20 Influence of Parenteral Iron Therapy and Oral Vitamin E Supplementation on Neutrophil 13 2.9 3 Respiratory Burst in Chronic Hemodialysis Patients. Renal Failure, 2005, 27, 135-141 Continuing professional development crediting system for specialists in laboratory medicine within 2.5 28 EFLM national societies. Biochemia Medica, 2013, 23, 332-41 Procalcitonin interference in an immunometric calcitonin assay. Scandinavian Journal of Clinical and 11 2 Laboratory Investigation, 2011, 71, 157-62 Intravenous iron gluconate administration increases circulating PAPP-A in hemodialysis patients. 10 2.9 Renal Failure, 2005, 27, 707-11 Comparability of eight immunoassay procedures for the determination of CA 15-3 and related 5.9 2 markers. Clinical Chemistry and Laboratory Medicine, 2003, 41, 1087-94 The Bank of Biological Material (BBM) of the First Faculty of Medicine of Charles University in 2.1 1 Prague, Czech Republic. Biopreservation and Biobanking, 2015, 13, 299-300 The protective effects of erythropoietin on rat glomerular podocytes in culture are modulated by 7 3.1 1 extracellular matrix proteins. Kidney and Blood Pressure Research, 2013, 38, 142-55 Advanced glycation end products and acute myocardial infarction. Medical Principles and Practice, 2.1 2010, 19, 244-6 Pregnancy-associated plasma protein A (PAPP-A) and soluble receptor for advanced glycation end products (sRAGE)--intra- and inter-individual variability in chronic hemodialysis patients. 5 2 1

Scandinavian Journal of Clinical and Laboratory Investigation, 2012, 72, 296-303

393, 45-50

Studies of biochemistry and clinical biochemistry. Studies at sample medical schools in 13 EU countries regarding biochemistry and clinical biochemistry teaching. *Clinica Chimica Acta*, **2008**,

6.2



3	The influence of pefloxacine on experimental adriamycin-induced nephrotic syndrome in rats. <i>Renal Failure</i> , 1996 , 18, 195-9	2.9	1
2	FP124MATRIX METALLOPROTEINASES (MMP-2, 3, 7, 9) AND THEIR TISSUE INHIBITORS (TIMP-1, 2) IN PATIENTS WITH SYSTEMIC LUPUS ERYTHEMATOSUS. <i>Nephrology Dialysis Transplantation</i> , 2015 , 30, iii108-iii108	4.3	
1	Laboratory screening markers in gastroenterologystate of the art. <i>Biomedical Papers of the Medical Faculty of the University Palacky&#x0301;, Olomouc, Czechoslovakia</i> , 2013 , 157, 91-7	1.7	