Toms Zima

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/1685976/tomas-zima-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

146
papers3,163
citations31
h-index47
g-index165
ext. papers3,468
ext. citations3.5
avg, IF4.75
L-index

#	Paper	IF	Citations
146	Galactose-deficient IgA1 and the corresponding IgG autoantibodies predict IgA nephropathy progression. <i>PLoS ONE</i> , 2019 , 14, e0212254	3.7	15
145	Gene Expression Analysis of Immunomagnetically Enriched Circulating Tumor Cell Fraction in Castration-Resistant Prostate Cancer. <i>Molecular Diagnosis and Therapy</i> , 2018 , 22, 381-390	4.5	3
144	Thyroid in pregnancy: From physiology to screening. <i>Critical Reviews in Clinical Laboratory Sciences</i> , 2017 , 54, 102-116	9.4	56
143	Cell-free DNA is higher and more fragmented in intrahepatic cholestasis of pregnancy. <i>Prenatal Diagnosis</i> , 2016 , 36, 1156-1158	3.2	4
142	Prolonged progesterone administration is associated with less frequent cervicovaginal colonization by Ureaplasma urealyticum during pregnancy - Results of a pilot study. <i>Journal of Reproductive Immunology</i> , 2016 , 116, 35-41	4.2	5
141	Surface plasmon resonance biosensor for detection of pregnancy associated plasma protein A2 in clinical samples. <i>Analytical and Bioanalytical Chemistry</i> , 2016 , 408, 7265-9	4.4	19
140	Serum trough infliximab levels: A comparison of three different immunoassays for the monitoring of CT-P13 (infliximab) treatment in patients with inflammatory bowel disease. <i>Biologicals</i> , 2016 , 44, 33	-6 ^{1.8}	30
139	The characterization of four gene expression analysis in circulating tumor cells made by Multiplex-PCR from the AdnaTest kit on the lab-on-a-chip Agilent DNA 1000 platform. <i>Biochemia Medica</i> , 2016 , 26, 103-13	2.5	6
138	HMGB1, S100 proteins and other RAGE ligands in cancer - markers, mediators and putative therapeutic targets. <i>Biomedical Papers of the Medical Faculty of the University Palacky&#x0301;, Olomouc, Czechoslovakia</i> , 2016 , 160, 1-10	1.7	22
137	Early detection of pancreatic cancer: impact of high-resolution imaging methods and biomarkers. <i>European Journal of Gastroenterology and Hepatology</i> , 2016 , 28, e33-e43	2.2	9
136	Polymorphisms of the receptor for advanced glycation end-products and glyoxalase I in patients with renal cancer. <i>Tumor Biology</i> , 2015 , 36, 2121-6	2.9	19
135	Panel of Urinary Diagnostic Markers for Non-Invasive Detection of Primary and Recurrent Urothelial Urinary Bladder Carcinoma. <i>Urologia Internationalis</i> , 2015 , 95, 56-64	1.9	29
134	Diagnostic Importance of Selected Protein Serum Markers in the Primary Diagnostics of Prostate Cancer. <i>Urologia Internationalis</i> , 2015 , 95, 429-35	1.9	10
133	The Bank of Biological Material (BBM) of the First Faculty of Medicine of Charles University in Prague, Czech Republic. <i>Biopreservation and Biobanking</i> , 2015 , 13, 299-300	2.1	1
132	Diagnostic and prognostic value of presepsin in preterm deliveries. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2015 , 28, 1032-7	2	3
131	Predictive factors for the presence of tumor cells in bone marrow and peripheral blood in breast cancer patients. <i>Neoplasma</i> , 2015 , 62, 259-68	3.3	6
130	Correlation of Vitreous Vascular Endothelial Growth Factor and Uric Acid Concentration Using Optical Coherence Tomography in Diabetic Macular Edema. <i>Journal of Ophthalmology</i> , 2015 , 2015, 478	s5 6 9	15

129	RAGE and its ligands in cancer - culprits, biomarkers, or therapeutic targets?. <i>Neoplasma</i> , 2015 , 62, 353-	6 4 .3	16	
128	FP124MATRIX METALLOPROTEINASES (MMP-2, 3, 7, 9) AND THEIR TISSUE INHIBITORS (TIMP-1, 2) IN PATIENTS WITH SYSTEMIC LUPUS ERYTHEMATOSUS. <i>Nephrology Dialysis Transplantation</i> , 2015 , 30, iii108-iii108	4.3		
127	Impact of risk factors on prevalence of anal HPV infection in women with simultaneous cervical lesion. <i>Neoplasma</i> , 2015 , 62, 308-14	3.3	18	
126	Trefoil factor family (TFF) proteins as potential serum biomarkers in patients with metastatic colorectal cancer. <i>Neoplasma</i> , 2015 , 62, 470-7	3.3	16	
125	Vitamin D Binding Protein Is Not Involved in Vitamin D Deficiency in Patients with Chronic Kidney Disease. <i>BioMed Research International</i> , 2015 , 2015, 492365	3	25	
124	A randomised controlled trial of roller versus centrifugal cardiopulmonary bypass pumps in patients undergoing pulmonary endarterectomy. <i>Perfusion (United Kingdom)</i> , 2015 , 30, 520-8	1.9	11	
123	The brighter (and evolutionarily older) face of the metabolic syndrome: evidence from Trypanosoma cruzi infection in CD-1 mice. <i>Diabetes/Metabolism Research and Reviews</i> , 2015 , 31, 346-35	97.5	23	
122	The relationship between the cervical and anal HPV infection in women with cervical intraepithelial neoplasia. <i>Journal of Clinical Virology</i> , 2014 , 59, 18-23	14.5	33	
121	Pregnancy-associated plasma protein A (PAPP-A) and preeclampsia. <i>Advances in Clinical Chemistry</i> , 2014 , 63, 169-209	5.8	22	
120	Reference intervals for thyroid markers in early pregnancy determined by 7 different analytical systems. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2014 , 74, 95-101	2	27	
119	Pregnancy-associated plasma protein A associates with cardiovascular events in diabetic hemodialysis patients. <i>Atherosclerosis</i> , 2014 , 236, 263-9	3.1	10	
118	Cys327Cys polymorphism of the PAPP-A gene (pregnancy associated plasma protein A) is related to mortality of long term hemodialysis patients. <i>Clinical Biochemistry</i> , 2014 , 47, 578-83	3.5	3	
117	Protective effects of inhaled carbon monoxide in endotoxin-induced cholestasis is dependent on its kinetics. <i>Biochimie</i> , 2014 , 97, 173-80	4.6	8	
116	Dissociative symptoms reflect levels of tumor necrosis factor alpha in patients with unipolar depression. <i>Neuropsychiatric Disease and Treatment</i> , 2014 , 10, 675-9	3.1	6	
115	Low glucose degradation product peritoneal dialysis regimen is associated with lower plasma EN-RAGE and HMGB-1 proinflammatory ligands of receptor for advanced glycation end products. <i>Therapeutic Apheresis and Dialysis</i> , 2014 , 18, 309-16	1.9	4	
114	Detection of circulating tumor cells during follow-up of patients with early breast cancer: Clinical utility for monitoring of therapy efficacy. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2014 , 74, 132-42	2	21	
113	Low levels of circulating T-regulatory lymphocytes and short cervical length are associated with preterm labor. <i>Journal of Reproductive Immunology</i> , 2014 , 106, 110-7	4.2	22	
112	Significantly higher procalcitonin levels could differentiate Gram-negative sepsis from Gram-positive and fungal sepsis. <i>Clinical and Experimental Medicine</i> , 2013 , 13, 165-70	4.9	76	

111	Relationship between serum bilirubin and uric acid to oxidative stress markers in Italian and Czech populations. <i>Journal of Applied Biomedicine</i> , 2013 , 11, 209-221	0.6	6
110	Placental growth factor may predict increased left ventricular mass index in patients with mild to moderate chronic kidney diseasea prospective observational study. <i>BMC Nephrology</i> , 2013 , 14, 142	2.7	10
109	Placental growth factor, pregnancy-associated plasma protein-A, soluble receptor for advanced glycation end products, extracellular newly identified receptor for receptor for advanced glycation end products binding protein and high mobility group box 1 levels in patients with acute kidney	2.7	13
108	Early vancomycin, amikacin and gentamicin concentrations in pulmonary artery and pulmonary tissue are not affected by VA ECMO (venoarterial extracorporeal membrane oxygenation) in a pig model of prolonged cardiac arrest. <i>Pulmonary Pharmacology and Therapeutics</i> , 2013 , 26, 655-60	3.5	4
107	Endotelial activation and flow-mediated vasodilation in young patients with breast cancer. <i>Neoplasma</i> , 2013 , 60, 690-7	3.3	6
106	Expression of Biliverdin Reductase A in peripheral blood leukocytes is associated with treatment response in HCV-infected patients. <i>PLoS ONE</i> , 2013 , 8, e57555	3.7	10
105	Skin autofluorescence relates to soluble receptor for advanced glycation end-products and albuminuria in diabetes mellitus. <i>Journal of Diabetes Research</i> , 2013 , 2013, 650694	3.9	10
104	The protective effects of erythropoietin on rat glomerular podocytes in culture are modulated by extracellular matrix proteins. <i>Kidney and Blood Pressure Research</i> , 2013 , 38, 142-55	3.1	1
103	Continuing professional development crediting system for specialists in laboratory medicine within 28 EFLM national societies. <i>Biochemia Medica</i> , 2013 , 23, 332-41	2.5	2
102	Changes in levels of matrix metalloproteinase-2 and -9, pregnancy-associated plasma protein-A in patients with various nephropathies. <i>Journal of Nephrology</i> , 2013 , 26, 502-9	4.8	4
101	Laboratory screening markers in gastroenterologystate of the art. <i>Biomedical Papers of the Medical Faculty of the University Palacky&#x0301;, Olomouc, Czechoslovakia</i> , 2013 , 157, 91-7	1.7	
100	Matrix metalloproteinases in serum and the follicular fluid of women treated by in vitro fertilization. <i>Journal of Assisted Reproduction and Genetics</i> , 2012 , 29, 1207-12	3.4	17
99	EN-RAGE (extracellular newly identified receptor for advanced glycation end-products binding protein) and mortality of long-term hemodialysis patients: A prospective observational cohort study. <i>Clinical Biochemistry</i> , 2012 , 45, 556-60	3.5	10
98	Receptor for advanced glycation end products (RAGE) and glyoxalase I gene polymorphisms in pathological pregnancy. <i>Clinical Biochemistry</i> , 2012 , 45, 1409-14	3.5	7
97	Immunological and biochemical markers in preeclampsia. <i>Journal of Reproductive Immunology</i> , 2012 , 96, 90-4	4.2	17
96	Ferritin as an independent mortality predictor in patients with pancreas cancer. Results of a pilot study. <i>Tumor Biology</i> , 2012 , 33, 1695-700	2.9	35
95	Pregnancy-associated plasma protein A as an independent mortality predictor in long-term hemodialysis patients. <i>Kidney and Blood Pressure Research</i> , 2012 , 35, 192-201	3.1	18
94	Advanced glycation end products in myocardial reperfusion injury. <i>Heart and Vessels</i> , 2012 , 27, 208-15	2.1	4

(2011-2012)

93	Biomarkers demonstrate increased consumption, but not abuse, of ethanol in essential tremor. Journal of Neurology, 2012 , 259, 1177-80	5.5	5
92	Pregnancy-associated plasma protein A (PAPP-A) and soluble receptor for advanced glycation end products (sRAGE)intra- and inter-individual variability in chronic hemodialysis patients. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2012 , 72, 296-303	2	1
91	Selenium: an important trace element and therapeutic adjunct in critical care. <i>Trace Elements and Electrolytes</i> , 2012 , 29, 246-255	1.8	4
90	Recommendations on prenatal screening and the connections to other diseases such as thyroid dysfunction. <i>Clinical Chemistry and Laboratory Medicine</i> , 2012 , 50, 1211-20	5.9	3
89	Relationship of soluble RAGE and RAGE ligands HMGB1 and EN-RAGE to endothelial dysfunction in type 1 and type 2 diabetes mellitus. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2012 , 120, 277-81	2.3	51
88	Pregnancy-associated plasma protein A: spotlight on kidney diseases. <i>Clinical Chemistry and Laboratory Medicine</i> , 2012 , 50, 1183-90	5.9	7
87	Improvements in colorectal cancer screening programmes - quantitative immunochemical faecal occult blood testing - how to set the cut-off for a particular population. <i>Biomedical Papers of the Medical Faculty of the University Palacky&#x0301;, Olomouc, Czechoslovakia</i> , 2012 , 156, 143-50	1.7	6
86	Procalcitonin interference in an immunometric calcitonin assay. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2011 , 71, 157-62	2	2
85	Placental growth factor in patients with decreased renal function. Renal Failure, 2011, 33, 291-7	2.9	9
84	High-risk human papillomavirus DNA in paraaortic lymph nodes in advanced stages of cervical carcinoma. <i>Journal of Clinical Virology</i> , 2011 , 50, 46-9	14.5	5
83	Serum S100A12 (EN-RAGE) levels in patients with decreased renal function and subclinical chronic inflammatory disease. <i>Kidney and Blood Pressure Research</i> , 2011 , 34, 457-64	3.1	12
82	Association of pregnancy-associated plasma protein A polymorphism with preeclampsia - a pilot study. <i>Clinical Biochemistry</i> , 2011 , 44, 1380-4	3.5	8
81	Leptin and adiponectin in pancreatic cancer: connection with diabetes mellitus. <i>Neoplasma</i> , 2011 , 58, 58-64	3.3	28
80	The quality of the extra-analytical phase of laboratory practice in some developing European countries and Mexico - a multicentric study. <i>Clinical Chemistry and Laboratory Medicine</i> , 2011 , 49, 215-28	₃ 5.9	23
79	Increased uric acid and glucose concentrations in vitreous and serum of patients with diabetic macular oedema. <i>Ophthalmic Research</i> , 2011 , 46, 73-9	2.9	25
78	Human galectins induce conversion of dermal fibroblasts into myofibroblasts and production of extracellular matrix: potential application in tissue engineering and wound repair. <i>Cells Tissues Organs</i> , 2011 , 194, 469-80	2.1	52
77	Interlaboratory study of free monoclonal immunoglobulin light chain quantification. <i>Clinical Chemistry and Laboratory Medicine</i> , 2011 , 49, 89-92	5.9	5
76	Associations of serum levels of advanced glycation end products with nutrition markers and anemia in patients with chronic kidney disease. <i>Renal Failure</i> , 2011 , 33, 131-7	2.9	8

75	Genetic predisposition to advanced glycation end products toxicity is related to prognosis of chronic hemodialysis patients. <i>Kidney and Blood Pressure Research</i> , 2010 , 33, 30-6	3.1	15
74	Low maternal serum matrix metalloproteinase (MMP)-2 concentrations are associated with preterm labor and fetal inflammatory response. <i>Journal of Perinatal Medicine</i> , 2010 , 38, 589-96	2.7	12
73	Advanced glycation end products and acute myocardial infarction. <i>Medical Principles and Practice</i> , 2010 , 19, 244-6	2.1	1
7 2	Anti-inflammatory effect of biological treatment in patients with inflammatory bowel diseases: calprotectin and IL-6 changes do not correspond to sRAGE changes. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2010 , 70, 294-9	2	12
71	Study of urinary proteomes in Anderson-Fabry disease. <i>Renal Failure</i> , 2010 , 32, 1202-9	2.9	18
70	Plasma calprotectin in chronically dialyzed end-stage renal disease patients. <i>Inflammation Research</i> , 2010 , 59, 299-305	7.2	7
69	Lower retinol levels as an independent predictor of mortality in long-term hemodialysis patients: a prospective observational cohort study. <i>American Journal of Kidney Diseases</i> , 2010 , 56, 513-21	7.4	24
68	Depression, traumatic stress and interleukin-6. <i>Journal of Affective Disorders</i> , 2010 , 120, 231-4	6.6	80
67	Soluble receptor for advanced glycation end products in physiological and pathological pregnancy. <i>Clinical Biochemistry</i> , 2010 , 43, 442-6	3.5	28
66	Soluble receptor for advanced glycation end-products (sRAGE) and polymorphisms of RAGE and glyoxalase I genes in patients with pancreas cancer. <i>Clinical Biochemistry</i> , 2010 , 43, 882-6	3.5	42
65	Glyoxalase I Glu111Ala polymorphism in patients with breast cancer. Cancer Investigation, 2009, 27, 655	-60	12
64	Fetuin-A early after renal transplantation. <i>Kidney and Blood Pressure Research</i> , 2009 , 32, 217-22	3.1	3
63	Reference intervals in evaluation of maternal thyroid function during the first trimester of pregnancy. <i>European Journal of Endocrinology</i> , 2009 , 160, 791-7	6.5	62
62	RAGE polymorphisms, renal function and histological finding at 12 months after renal transplantation. <i>Clinical Biochemistry</i> , 2009 , 42, 347-52	3.5	8
61	Fibroblast growth factor 23 and matrix-metalloproteinases in patients with chronic kidney disease: are they associated with cardiovascular disease?. <i>Kidney and Blood Pressure Research</i> , 2009 , 32, 276-83	3.1	23
60	Marked increase of procalcitonin after the administration of anti-thymocyte globulin in patients before hematopoietic stem cell transplantation does not indicate sepsis: a prospective study. <i>Critical Care</i> , 2009 , 13, R37	10.8	45
59	High-risk human papillomavirus DNA in the primary tumor, sentinel, and nonsentinel pelvic lymph nodes in patients with early-stage cervical cancer: a correlation with histopathology. <i>International Journal of Gynecological Cancer</i> , 2009 , 19, 703-7	3.5	13
58	Recent progress in the pathogenesis of nephrotic proteinuria. <i>Critical Reviews in Clinical Laboratory Sciences</i> , 2008 , 45, 139-220	9.4	14

(2006-2008)

57	Studies of biochemistry and clinical biochemistry. Studies at sample medical schools in 13 EU countries regarding biochemistry and clinical biochemistry teaching. <i>Clinica Chimica Acta</i> , 2008 , 393, 45-50	6.2	1
56	Detection of feto-maternal infection/inflammation by the soluble receptor for advanced glycation end products (sRAGE): results of a pilot study. <i>Journal of Perinatal Medicine</i> , 2008 , 36, 399-404	2.7	20
55	Breath alcohol level and plasma amino acids: a comparison between older and younger chronic alcohol-dependent patients. <i>Alcohol and Alcoholism</i> , 2008 , 43, 653-7	3.5	6
54	A419C (E111A) polymorphism of the glyoxalase I gene and vascular complications in chronic hemodialysis patients. <i>Annals of the New York Academy of Sciences</i> , 2008 , 1126, 268-71	6.5	22
53	Receptor for advanced glycation end products (RAGE)soluble form (sRAGE) and gene polymorphisms in patients with breast cancer. <i>Cancer Investigation</i> , 2007 , 25, 720-5	2.1	82
52	Quantification of trace elements in human serum fractions by liquid chromatography and inductively coupled plasma mass spectrometry. <i>Applied Organometallic Chemistry</i> , 2007 , 21, 5-14	3.1	7
51	Oxidative stress and endothelium influenced by metformin in type 2 diabetes mellitus. <i>European Journal of Clinical Pharmacology</i> , 2007 , 63, 1107-14	2.8	31
50	Performance characteristics of seven neuron-specific enolase assays. <i>Tumor Biology</i> , 2007 , 28, 84-92	2.9	36
49	Culture methods of glomerular podocytes. <i>Kidney and Blood Pressure Research</i> , 2007 , 30, 162-74	3.1	35
48	Receptor for advanced glycation end productssoluble form and gene polymorphisms in chronic haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2007 , 22, 2020-6	4.3	62
47	Does renal function influence plasma levels of advanced glycation and oxidation protein products in patients with chronic rheumatic diseases complicated by secondary amyloidosis?. <i>Kidney and Blood Pressure Research</i> , 2007 , 30, 1-7	3.1	8
46	Pregnancy-associated plasma protein a and soluble receptor for advanced glycation end products after kidney transplantation. <i>Kidney and Blood Pressure Research</i> , 2007 , 30, 31-7	3.1	21
45	Resting energy expenditure and thermal balance during isothermic and thermoneutral haemodialysisheat production does not explain increased body temperature during haemodialysis. <i>Nephrology Dialysis Transplantation</i> , 2007 , 22, 3553-60	4.3	12
44	No benefit of hemodiafiltration over hemodialysis in lowering elevated levels of asymmetric dimethylarginine in ESRD patients. <i>Blood Purification</i> , 2006 , 24, 439-44	3.1	25
43	On-line haemodiafiltration versus haemodialysis: stable haematocrit with less erythropoietin and improvement of other relevant blood parameters. <i>Blood Purification</i> , 2006 , 24, 163-73	3.1	89
42	Randomized prospective comparative study of ursodeoxycholic acid and S-adenosyl-L-methionine in the treatment of intrahepatic cholestasis of pregnancy. <i>Journal of Perinatal Medicine</i> , 2006 , 34, 383-	91 ^{2.7}	61
41	Impact of essential hypertension and primary aldosteronism on plasma brain natriuretic peptide concentration. <i>Blood Pressure</i> , 2006 , 15, 302-7	1.7	10
40	Influence of oral vitamin E therapy on micro-inflammation and cardiovascular disease markers in chronic hemodialysis patients. <i>Renal Failure</i> , 2006 , 28, 395-9	2.9	19

39	Effect of hemodiafiltration on pregnancy-associated plasma protein A (PAPP-A) and related parameters. <i>Renal Failure</i> , 2006 , 28, 715-21	2.9	8
38	Soluble receptor for advanced glycation end products in patients with decreased renal function. <i>American Journal of Kidney Diseases</i> , 2006 , 47, 406-11	7.4	128
37	Oxidative stress and inflammation in pregnancy. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2006 , 66, 121-7	2	49
36	Intravenous iron gluconate administration increases circulating PAPP-A in hemodialysis patients. <i>Renal Failure</i> , 2005 , 27, 707-11	2.9	2
35	Modulation of Oxidative Stress by Alcohol. Alcoholism: Clinical and Experimental Research, 2005, 29, 106	50 5.1/ 06	55 ₄
34	Oxidative stress and signal transduction pathways in alcoholic liver disease. <i>Alcoholism: Clinical and Experimental Research</i> , 2005 , 29, 110S-115S	3.7	98
33	Advanced glycoxidation end products in chronic diseases-clinical chemistry and genetic background. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2005 , 579, 37	-4 ² 6 ³	142
32	Dynamic genetic architecture of metabolic syndrome attributes in the rat. <i>Physiological Genomics</i> , 2005 , 21, 243-52	3.6	32
31	Influence of Parenteral Iron Therapy and Oral Vitamin E Supplementation on Neutrophil Respiratory Burst in Chronic Hemodialysis Patients. <i>Renal Failure</i> , 2005 , 27, 135-141	2.9	3
30	Advanced glycation end products in clinical nephrology. <i>Kidney and Blood Pressure Research</i> , 2004 , 27, 18-28	3.1	42
29	Relationship of pregnancy-associated plasma protein-a to renal function and dialysis modalities. <i>Kidney and Blood Pressure Research</i> , 2004 , 27, 88-95	3.1	36
28	Increased levels of pregnancy-associated plasma protein A are associated with mortality in hemodialysis patients: preliminary results. <i>Blood Purification</i> , 2004 , 22, 298-300	3.1	23
27	Advanced glycation end-products in patients with chronic alcohol misuse. <i>Alcohol and Alcoholism</i> , 2004 , 39, 316-20	3.5	35
26	Advanced glycation end products in hemodialyzed patients with diabetes mellitus correlate with leptin and leptin/body fat ratio. <i>Renal Failure</i> , 2003 , 25, 277-86	2.9	4
25	Glycoxidation and inflammation in chronic haemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2003 , 18, 2577-81	4.3	40
24	PAPP-A, a novel marker of unstable plaque, is not influenced by hypolipidemic treatment in contrast to CRP. <i>Atherosclerosis</i> , 2003 , 166, 195-6	3.1	6
23	Antiphospholipid antibodies in patients with lupus nephritis. <i>Renal Failure</i> , 2003 , 25, 747-58	2.9	4
22	Comparability of eight immunoassay procedures for the determination of CA 15-3 and related markers. Clinical Chemistry and Laboratory Medicine, 2003, 41, 1087-94	5.9	2

21	Pathobiochemistry of nephrotic syndrome. Advances in Clinical Chemistry, 2003, 37, 173-218	5.8	9
20	Qualitative and quantitative detection of urinary human complement factor H-related protein (BTA stat and BTA TRAK) and fragments of cytokeratins 8, 18 (UBC rapid and UBC IRMA) as markers for transitional cell carcinoma of the bladder. <i>European Urology</i> , 2002 , 41, 34-9	10.2	47
19	Advanced glycation end products and advanced oxidation protein products in hemodialyzed patients. <i>Blood Purification</i> , 2002 , 20, 531-6	3.1	49
18	Acute doxorubicin (adriamycin) dosage does not reduce cardiac protein synthesis in vivo, but decreases diaminopeptidase I and proline endopeptidase activities. <i>Experimental and Molecular Pathology</i> , 2001 , 70, 154-61	4.4	6
17	Effects of Doxorubicin (Adriamycin) and [(+)-1,2-bis(3,5-dioxopiperazinyl-1-yl)]propane (ICRF-187) on skeletal muscle protease activities. <i>Toxicology and Applied Pharmacology</i> , 2001 , 171, 135-40	4.6	9
16	Comparison of different immunoassays for CA 19-9. <i>Clinical Chemistry and Laboratory Medicine</i> , 2001 , 39, 1278-82	5.9	20
15	Acute dosage with dexrazoxane, but not doxorubicin, is associated with increased rates of hepatic protein synthesis in vivo. <i>Toxicologic Pathology</i> , 2001 , 29, 591-9	2.1	4
14	Oxidative stress, metabolism of ethanol and alcohol-related diseases 2001 , 8, 59		10
13	Trace elements in end-stage renal disease. 2. Clinical implication of trace elements. <i>Blood Purification</i> , 1999 , 17, 187-98	3.1	46
12	Trace elements in end-stage renal disease. 1. Methodological aspects and the influence of water treatment and dialysis equipment. <i>Blood Purification</i> , 1999 , 17, 182-6	3.1	9
11	Retinol and alpha-tocopherol in hemodialysis patients. <i>Renal Failure</i> , 1998 , 20, 505-12	2.9	12
10	The effect of silibinin on experimental cyclosporine nephrotoxicity. Renal Failure, 1998, 20, 471-9	2.9	19
9	Trace elements in hemodialysis and continuous ambulatory peritoneal dialysis patients. <i>Blood Purification</i> , 1998 , 16, 253-60	3.1	50
8	ICRF-187 (dexrazoxan) protects from adriamycin-induced nephrotic syndrome in rats. <i>Nephrology Dialysis Transplantation</i> , 1998 , 13, 1975-9	4.3	21
7	Influence of plasma exchange on serum levels of cytokines and adhesion molecules in ANCA-positive renal vasculitis. <i>Blood Purification</i> , 1998 , 16, 72-80	3.1	51
6	Comparison of the Suitability of Various Atomic SpectroscopicTechinques for the Determination of Selenium in Human Whole Blood. <i>Journal of Analytical Atomic Spectrometry</i> , 1997 , 12, 85-89	3.7	26
5	The effect of quinolinate on rat brain lipid peroxidation is dependent on iron. <i>Neurochemistry International</i> , 1997 , 30, 233-237	4.4	79
4	Antioxidant enzymessuperoxide dismutase and glutathione peroxidasein haemodialyzed patients. <i>Blood Purification</i> , 1996 , 14, 257-61	3.1	20

3	Lipid peroxidation and antioxidant enzymes in CAPD patients. <i>Renal Failure</i> , 1996 , 18, 113-9	2.9	8
2	The influence of pefloxacine on experimental adriamycin-induced nephrotic syndrome in rats. <i>Renal Failure</i> , 1996 , 18, 195-9	2.9	1
1	Diagnosis and treatment of carpal tunnel syndrome. <i>Renal Failure</i> , 1993 , 15, 533-7	2.9	7