

Martin Schobben

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1705748/publications.pdf>

Version: 2024-02-01

17
papers

549
citations

687363

13
h-index

888059

17
g-index

23
all docs

23
docs citations

23
times ranked

624
citing authors

#	ARTICLE	IF	CITATIONS
1	Baghuk Mountain (Central Iran): high-resolution stratigraphy of a continuous Central Tethyan Permian–Triassic boundary section. <i>Fossil Record</i> , 2021, 24, 171-192.	1.4	3
2	The effect of geographic range and climate on extinction risk in the deep-time amphibian fossil record. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2020, 537, 109414.	2.3	2
3	Macroecological patterns of the terrestrial vegetation history during the end-Triassic biotic crisis in the central European Basin: A palynological study of the Bonenburg section (NW-Germany) and its supra-regional implications. <i>Global and Planetary Change</i> , 2020, 194, 103286.	3.5	27
4	A nutrient control on marine anoxia during the end-Permian mass extinction. <i>Nature Geoscience</i> , 2020, 13, 640-646.	12.9	56
5	Catastrophic soil loss associated with end-Triassic deforestation. <i>Earth-Science Reviews</i> , 2020, 210, 103332.	9.1	34
6	Aras Valley (northwest Iran): high-resolution stratigraphy of a continuous central Tethyan Permian–Triassic boundary section. <i>Fossil Record</i> , 2020, 23, 33-69.	1.4	12
7	Interpreting the Carbon Isotope Record of Mass Extinctions. <i>Elements</i> , 2019, 15, 331-337.	0.5	29
8	Increased Stability in Carbon Isotope Records Reflects Emerging Complexity of the Biosphere. <i>Frontiers in Earth Science</i> , 2019, 7, .	1.8	7
9	A comparative study of total organic carbon- $\delta^{13}\text{C}$ signatures in the Triassic–Jurassic transitional beds of the Central European Basin and western Tethys shelf seas. <i>Newsletters on Stratigraphy</i> , 2019, 52, 461-486.	1.2	14
10	Pre-mass extinction decline of latest Permian ammonoids. <i>Geology</i> , 2018, 46, 283-286.	4.4	30
11	Volatile earliest Triassic sulfur cycle: A consequence of persistent low seawater sulfate concentrations and a high sulfur cycle turnover rate?. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2017, 486, 74-85.	2.3	20
12	Latest Permian carbonate carbon isotope variability traces heterogeneous organic carbon accumulation and authigenic carbonate formation. <i>Climate of the Past</i> , 2017, 13, 1635-1659.	3.4	18
13	Eutrophication, microbial-sulfate reduction and mass extinctions. <i>Communicative and Integrative Biology</i> , 2016, 9, e1115162.	1.4	17
14	Discerning primary versus diagenetic signals in carbonate carbon and oxygen isotope records: An example from the Permian–Triassic boundary of Iran. <i>Chemical Geology</i> , 2016, 422, 94-107.	3.3	65
15	The ammonoids from the Late Permian <i>Paratirolites</i> Limestone of Julfa (East Azerbaijan, Iran). <i>Journal of Systematic Palaeontology</i> , 2016, 14, 841-890.	1.5	16
16	Flourishing ocean drives the end-Permian marine mass extinction. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, 10298-10303.	7.1	78
17	Palaeotethys seawater temperature rise and an intensified hydrological cycle following the end-Permian mass extinction. <i>Gondwana Research</i> , 2014, 26, 675-683.	6.0	114