

Kovcs Zsolt

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1711250/kovacs-zsolt-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

53
papers

951
citations

19
h-index

29
g-index

66
ext. papers

1,132
ext. citations

4.2
avg, IF

4.28
L-index

#	Paper	IF	Citations
53	Adenosine Receptors Modulate the Exogenous Ketogenic Supplement-Evoked Alleviating Effect on Lipopolysaccharide-Generated Increase in Absence Epileptic Activity in WAG/Rij Rats. <i>Nutrients</i> , 2021 , 13,	6.7	0
52	Critical Role of Astrocytic Polyamine and GABA Metabolism in Epileptogenesis.. <i>Frontiers in Cellular Neuroscience</i> , 2021 , 15, 787319	6.1	1
51	Beneficial Effects of Exogenous Ketogenic Supplements on Aging Processes and Age-Related Neurodegenerative Diseases. <i>Nutrients</i> , 2021 , 13,	6.7	2
50	Enhancement of Ketone Supplements-Evoked Effect on Absence Epileptic Activity by Co-Administration of Uridine in Wistar Albino Glaxo Rijswijk Rats. <i>Nutrients</i> , 2021 , 13,	6.7	4
49	Inhibition of adenosine A1 receptors abolished the nutritional ketosis-evoked delay in the onset of isoflurane-induced anesthesia in Wistar Albino Glaxo Rijswijk rats. <i>BMC Anesthesiology</i> , 2020 , 20, 30	2.4	3
48	Exogenous Ketone Supplements Improved Motor Performance in Preclinical Rodent Models. <i>Nutrients</i> , 2020 , 12,	6.7	5
47	Age- and Sex-Dependent Modulation of Exogenous Ketone Supplement-Evoked Effects on Blood Glucose and Ketone Body Levels in Wistar Albino Glaxo Rijswijk Rats. <i>Frontiers in Neuroscience</i> , 2020 , 14, 618422	5.1	2
46	Exogenous Ketones Lower Blood Glucose Level in Rested and Exercised Rodent Models. <i>Nutrients</i> , 2019 , 11,	6.7	15
45	Elevated Plus Maze Test Combined with Video Tracking Software to Investigate the Anxiolytic Effect of Exogenous Ketogenic Supplements. <i>Journal of Visualized Experiments</i> , 2019 ,	1.6	6
44	Connexin 43 Differentially Regulates Epileptiform Activity in Models of Convulsive and Non-convulsive Epilepsies. <i>Frontiers in Cellular Neuroscience</i> , 2019 , 13, 173	6.1	13
43	Therapeutic Potential of Exogenous Ketone Supplement Induced Ketosis in the Treatment of Psychiatric Disorders: Review of Current Literature. <i>Frontiers in Psychiatry</i> , 2019 , 10, 363	5	21
42	Exogenous Ketone Supplementation Decreased the Lipopolysaccharide-Induced Increase in Absence Epileptic Activity in Wistar Albino Glaxo Rijswijk Rats. <i>Frontiers in Molecular Neuroscience</i> , 2019 , 12, 45	6.1	11
41	Uric acid and allopurinol aggravate absence epileptic activity in Wistar Albino Glaxo Rijswijk rats. <i>Brain Research</i> , 2018 , 1686, 1-9	3.7	2
40	Suckling induced activation pattern in the brain of rat pups. <i>Nutritional Neuroscience</i> , 2018 , 21, 317-327	3.6	3
39	Anxiolytic Effect of Exogenous Ketone Supplementation Is Abolished by Adenosine A1 Receptor Inhibition in Wistar Albino Glaxo/Rijswijk Rats. <i>Frontiers in Behavioral Neuroscience</i> , 2018 , 12, 29	3.5	16
38	Alterations in hippocampal and cortical densities of functionally different interneurons in rat models of absence epilepsy. <i>Epilepsy Research</i> , 2018 , 145, 40-50	3	11
37	The Role of Lipids and Membranes in the Pathogenesis of Alzheimer's Disease: A Comprehensive View. <i>Current Alzheimer Research</i> , 2018 , 15, 1191-1212	3	34

36	Exogenous ketones lower blood glucose level. <i>FASEB Journal</i> , 2018 , 32, 925-11	0.9	
35	Nutritional ketosis delays the onset of isoflurane induced anesthesia. <i>BMC Anesthesiology</i> , 2018 , 18, 85	2.4	9
34	Absence epileptic activity in Wistar Albino Glaxo Rijswijk rat mothers. <i>Brain Research</i> , 2017 , 1657, 368-376	3.7	2
33	Adenosine A1 Receptor Antagonism Abolished the Anti-seizure Effects of Exogenous Ketone Supplementation in Wistar Albino Glaxo Rijswijk Rats. <i>Frontiers in Molecular Neuroscience</i> , 2017 , 10, 235	6.1	32
32	Exogenous Ketone Supplements Reduce Anxiety-Related Behavior in Sprague-Dawley and Wistar Albino Glaxo/Rijswijk Rats. <i>Frontiers in Molecular Neuroscience</i> , 2016 , 9, 137	6.1	50
31	Guanosine may increase absence epileptic activity by means of A2A adenosine receptors in Wistar Albino Glaxo Rijswijk rats. <i>Brain Research Bulletin</i> , 2016 , 124, 172-81	3.9	11
30	Absence epileptic activity changing effects of non-adenosine nucleoside inosine, guanosine and uridine in Wistar Albino Glaxo Rijswijk rats. <i>Neuroscience</i> , 2015 , 300, 593-608	3.9	25
29	Modulatory effects of inosine, guanosine and uridine on lipopolysaccharide-evoked increase in spike-wave discharge activity in Wistar Albino Glaxo/Rijswijk rats. <i>Brain Research Bulletin</i> , 2015 , 118, 46-57	3.9	8
28	Non-adenosine nucleoside inosine, guanosine and uridine as promising antiepileptic drugs: a summary of current literature. <i>Mini-Reviews in Medicinal Chemistry</i> , 2015 , 14, 1033-42	3.2	19
27	Effects of nucleosides on glia - neuron interactions open up new vistas in the development of more effective antiepileptic drugs. <i>Current Medicinal Chemistry</i> , 2015 , 22, 1500-14	4.3	3
26	Dysfunction of Endoplasmic Reticulum (ER) and Mitochondria (MT) in Alzheimer's Disease: The Role of the ER-MT Cross-Talk. <i>Current Alzheimer Research</i> , 2015 , 12, 655-72	3	45
25	PACAP application improves functional outcome of chronic retinal ischemic injury in rats-evidence from electroretinographic measurements. <i>Journal of Molecular Neuroscience</i> , 2014 , 54, 293-9	3.3	33
24	Brain protein expression changes in WAG/Rij rats, a genetic rat model of absence epilepsy after peripheral lipopolysaccharide treatment. <i>Brain, Behavior, and Immunity</i> , 2014 , 35, 86-95	16.6	18
23	Lipopolysaccharide induced increase in seizure activity in two animal models of absence epilepsy WAG/Rij and GAERS rats and Long Evans rats. <i>Brain Research Bulletin</i> , 2014 , 104, 7-18	3.9	45
22	Receptors of peptides as therapeutic targets in epilepsy research. <i>Current Medicinal Chemistry</i> , 2014 , 21, 764-87	4.3	36
21	The antiepileptic potential of nucleosides. <i>Current Medicinal Chemistry</i> , 2014 , 21, 788-821	4.3	26
20	Anatomical Distribution of Nucleoside System in the Human Brain and Implications for Therapy 2013 , 621-656		1
19	Uridine modulates neuronal activity and inhibits spike-wave discharges of absence epileptic Long Evans and Wistar Albino Glaxo/Rijswijk rats. <i>Brain Research Bulletin</i> , 2013 , 97, 16-23	3.9	15

18	5'-nucleotidases, nucleosides and their distribution in the brain: pathological and therapeutic implications. <i>Current Medicinal Chemistry</i> , 2013 , 20, 4217-40	4.3	19
17	Neonatal tricyclic antidepressant clomipramine treatment reduces the spike-wave discharge activity of the adult WAG/Rij rat. <i>Brain Research Bulletin</i> , 2012 , 89, 102-7	3.9	10
16	Intracerebroventricularly administered lipopolysaccharide enhances spike-wave discharges in freely moving WAG/Rij rats. <i>Brain Research Bulletin</i> , 2011 , 85, 410-6	3.9	48
15	Doxycycline could aggravate the absence-like epileptic seizures of WAG/Rij rats via matrix metalloproteinase inhibition. <i>Neurochemistry International</i> , 2011 , 59, 563-6	4.4	5
14	Area, age and gender dependence of the nucleoside system in the brain: a review of current literature. <i>Current Topics in Medicinal Chemistry</i> , 2011 , 11, 1012-33	3	32
13	The effect of intraperitoneally administered dimethyl sulfoxide on absence-like epileptic activity of freely moving WAG/Rij rats. <i>Journal of Neuroscience Methods</i> , 2011 , 197, 133-6	3	19
12	Uridine function in the central nervous system. <i>Current Topics in Medicinal Chemistry</i> , 2011 , 11, 1058-67	3	45
11	Matrix metalloproteinase-9 activity increased by two different types of epileptic seizures that do not induce neuronal death: a possible role in homeostatic synaptic plasticity. <i>Neurochemistry International</i> , 2010 , 56, 799-809	4.4	46
10	Gender- and age-dependent changes in nucleoside levels in the cerebral cortex and white matter of the human brain. <i>Brain Research Bulletin</i> , 2010 , 81, 579-84	3.9	12
9	Nucleoside map of the human central nervous system. <i>Neurochemical Research</i> , 2010 , 35, 452-64	4.6	24
8	Clomipramine increases the incidence and duration of spike-wave discharges in freely moving WAG/Rij rats. <i>Epilepsy Research</i> , 2010 , 90, 167-70	3	6
7	Genistein induces phosphorylation of cAMP response element-binding protein in neonatal hypothalamus in vivo. <i>Journal of Neuroendocrinology</i> , 2009 , 21, 1024-8	3.8	7
6	Regional distribution and effects of postmortal delay on endocannabinoid content of the human brain. <i>Neuroscience</i> , 2008 , 152, 1032-9	3.9	37
5	Suppression of spike-wave discharge activity and c-fos expression by 2-methyl-4-oxo-3H-quinazoline-3-acetyl piperidine (Q5) in vivo. <i>Neuroscience Letters</i> , 2007 , 423, 73-7	3.3	6
4	Facilitation of spike-wave discharge activity by lipopolysaccharides in Wistar Albino Glaxo/Rijswijk rats. <i>Neuroscience</i> , 2006 , 140, 731-42	3.9	57
3	Concentration of nucleosides and related compounds in cerebral and cerebellar cortical areas and white matter of the human brain. <i>Cellular and Molecular Neurobiology</i> , 2006 , 26, 833-44	4.6	9
2	Post mortem degradation of nucleosides in the brain: comparison of human and rat brains for estimation of in vivo concentration of nucleosides. <i>Journal of Neuroscience Methods</i> , 2005 , 148, 88-93	3	16
1	Uridine is released by depolarization and inhibits unit activity in the rat hippocampus. <i>NeuroReport</i> , 1999 , 10, 3049-53	1.7	19

