

Annette H Van Der Helm-Van Mil

List of Publications by Citations

Source:

<https://exaly.com/author-pdf/1711536/annette-h-van-der-helm-van-mil-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

263
papers

11,856
citations

52
h-index

101
g-index

293
ext. papers

14,026
ext. citations

4.8
avg, IF

6.23
L-index

#	Paper	IF	Citations
263	Genome-wide association study meta-analysis identifies seven new rheumatoid arthritis risk loci. <i>Nature Genetics</i> , 2010 , 42, 508-14	36.3	969
262	Refining the complex rheumatoid arthritis phenotype based on specificity of the HLA-DRB1 shared epitope for antibodies to citrullinated proteins. <i>Arthritis and Rheumatism</i> , 2005 , 52, 3433-8		427
261	Common variants at CD40 and other loci confer risk of rheumatoid arthritis. <i>Nature Genetics</i> , 2008 , 40, 1216-23	36.3	416
260	Autoantibodies recognizing carbamylated proteins are present in sera of patients with rheumatoid arthritis and predict joint damage. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 17372-7	11.5	385
259	Antibodies to citrullinated proteins and differences in clinical progression of rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2005 , 7, R949-58	5.7	325
258	Gene-gene and gene-environment interactions involving HLA-DRB1, PTPN22, and smoking in two subsets of rheumatoid arthritis. <i>American Journal of Human Genetics</i> , 2007 , 80, 867-75	11	317
257	Genetic variants at CD28, PRDM1 and CD2/CD58 are associated with rheumatoid arthritis risk. <i>Nature Genetics</i> , 2009 , 41, 1313-8	36.3	272
256	A prediction rule for disease outcome in patients with recent-onset undifferentiated arthritis: how to guide individual treatment decisions. <i>Arthritis and Rheumatism</i> , 2007 , 56, 433-40		267
255	EULAR recommendations for terminology and research in individuals at risk of rheumatoid arthritis: report from the Study Group for Risk Factors for Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 638-41	2.4	263
254	Long-term impact of delay in assessment of patients with early arthritis. <i>Arthritis and Rheumatism</i> , 2010 , 62, 3537-46		262
253	The HLA-DRB1 shared epitope alleles are primarily a risk factor for anti-cyclic citrullinated peptide antibodies and are not an independent risk factor for development of rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2006 , 54, 1117-21		253
252	2016 update of the EULAR recommendations for the management of early arthritis. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 948-959	2.4	243
251	Smoking is a risk factor for anti-CCP antibodies only in rheumatoid arthritis patients who carry HLA-DRB1 shared epitope alleles. <i>Annals of the Rheumatic Diseases</i> , 2006 , 65, 366-71	2.4	216
250	Predicting arthritis outcomes--what can be learned from the Leiden Early Arthritis Clinic?. <i>Rheumatology</i> , 2011 , 50, 93-100	3.9	207
249	A candidate gene approach identifies the TRAF1/C5 region as a risk factor for rheumatoid arthritis. <i>PLoS Medicine</i> , 2007 , 4, e278	11.6	192
248	What is the evidence for the presence of a therapeutic window of opportunity in rheumatoid arthritis? A systematic literature review. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 861-70	2.4	182
247	Classification of rheumatoid arthritis: comparison of the 1987 American College of Rheumatology criteria and the 2010 American College of Rheumatology/European League Against Rheumatism criteria. <i>Arthritis and Rheumatism</i> , 2011 , 63, 37-42		168

246	Quantitative heritability of anti-citrullinated protein antibody-positive and anti-citrullinated protein antibody-negative rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2009 , 60, 916-23		158
245	Prevalence of and predictive factors for sustained disease-modifying antirheumatic drug-free remission in rheumatoid arthritis: results from two large early arthritis cohorts. <i>Arthritis and Rheumatism</i> , 2009 , 60, 2262-71		154
244	Association of HLA-DR3 with anti-cyclic citrullinated peptide antibody-negative rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2005 , 52, 3058-62		140
243	A high body mass index has a protective effect on the amount of joint destruction in small joints in early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2008 , 67, 769-74	2.4	132
242	Validation of a novel multibiomarker test to assess rheumatoid arthritis disease activity. <i>Arthritis Care and Research</i> , 2012 , 64, 1794-803	4.7	130
241	EULAR definition of arthralgia suspicious for progression to rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 491-496	2.4	125
240	Validation of a prediction rule for disease outcome in patients with recent-onset undifferentiated arthritis: moving toward individualized treatment decision-making. <i>Arthritis and Rheumatism</i> , 2008 , 58, 2241-7		121
239	Update on the epidemiology, risk factors, and disease outcomes of rheumatoid arthritis. <i>Best Practice and Research in Clinical Rheumatology</i> , 2018 , 32, 174-187	5.3	120
238	Value of anti-modified citrullinated vimentin and third-generation anti-cyclic citrullinated peptide compared with second-generation anti-cyclic citrullinated peptide and rheumatoid factor in predicting disease outcome in undifferentiated arthritis and rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2009 , 60, 2232-41		114
237	The HLA-DRB1 shared epitope alleles differ in the interaction with smoking and predisposition to antibodies to cyclic citrullinated peptide. <i>Arthritis and Rheumatism</i> , 2007 , 56, 425-32		112
236	Invasiveness of fibroblast-like synoviocytes is an individual patient characteristic associated with the rate of joint destruction in patients with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2005 , 52, 1999-2002		110
235	Advances in the genetics of rheumatoid arthritis point to subclassification into distinct disease subsets. <i>Arthritis Research and Therapy</i> , 2008 , 10, 205	5.7	101
234	Molecular Portraits of Early Rheumatoid Arthritis Identify Clinical and Treatment Response Phenotypes. <i>Cell Reports</i> , 2019 , 28, 2455-2470.e5	10.6	100
233	Synovial cellular and molecular signatures stratify clinical response to csDMARD therapy and predict radiographic progression in early rheumatoid arthritis patients. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 761-772	2.4	96
232	Clinical factors, anticitrullinated peptide antibodies and MRI-detected subclinical inflammation in relation to progression from clinically suspect arthralgia to arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1824-30	2.4	85
231	EULAR definition of erosive disease in light of the 2010 ACR/EULAR rheumatoid arthritis classification criteria. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 479-81	2.4	85
230	Association of a single-nucleotide polymorphism in CD40 with the rate of joint destruction in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2009 , 60, 2242-7		84
229	Baseline serum adipokine levels predict radiographic progression in early rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2011 , 63, 2567-74		83

228	An independent role of protective HLA class II alleles in rheumatoid arthritis severity and susceptibility. <i>Arthritis and Rheumatism</i> , 2005 , 52, 2637-44		77
227	Understanding the genetic contribution to rheumatoid arthritis. <i>Current Opinion in Rheumatology</i> , 2005 , 17, 299-304	5.3	75
226	Rheumatoid arthritis risk allele PTPRC is also associated with response to anti-tumor necrosis factor alpha therapy. <i>Arthritis and Rheumatism</i> , 2010 , 62, 1849-61		74
225	A genetic study on C5-TRAF1 and progression of joint damage in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2015 , 17, 1	5.7	70
224	Characterising arthralgia in the preclinical phase of rheumatoid arthritis using MRI. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1225-32	2.4	69
223	Association of the PTPN22 C1858T single-nucleotide polymorphism with rheumatoid arthritis phenotypes in an inception cohort. <i>Arthritis and Rheumatism</i> , 2005 , 52, 2948-50		68
222	Strategies to predict rheumatoid arthritis development in at-risk populations. <i>Rheumatology</i> , 2016 , 55, 6-15	3.9	63
221	A large-scale rheumatoid arthritis genetic study identifies association at chromosome 9q33.2. <i>PLoS Genetics</i> , 2008 , 4, e1000107	6	63
220	Preventing progression from arthralgia to arthritis: targeting the right patients. <i>Nature Reviews Rheumatology</i> , 2018 , 14, 32-41	8.1	59
219	The ACPA isotype profile reflects long-term radiographic progression in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1110-6	2.4	58
218	Genetic predisposition of the severity of joint destruction in rheumatoid arthritis: a population-based study. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 707-9	2.4	57
217	The specificity of anti-carbamylated protein antibodies for rheumatoid arthritis in a setting of early arthritis. <i>Arthritis Research and Therapy</i> , 2015 , 17, 339	5.7	55
216	Triple Positivity for Anti-Citrullinated Protein Autoantibodies, Rheumatoid Factor, and Anti-Carbamylated Protein Antibodies Conferring High Specificity for Rheumatoid Arthritis: Implications for Very Early Identification of At-Risk Individuals. <i>Arthritis and Rheumatology</i> , 2018 , 70, 1721-1731	9.5	55
215	An evaluation of molecular and clinical remission in rheumatoid arthritis by assessing radiographic progression. <i>Rheumatology</i> , 2013 , 52, 839-46	3.9	54
214	The ACPA recognition profile and subgrouping of ACPA-positive RA patients. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 268-74	2.4	54
213	The association between anti-carbamylated protein (anti-CarP) antibodies and radiographic progression in early rheumatoid arthritis: a study exploring replication and the added value to ACPA and rheumatoid factor. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 112-118	2.4	53
212	Genetic studies on components of the Wnt signalling pathway and the severity of joint destruction in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 769-75	2.4	53
211	Protective effect of noninherited maternal HLA-DR antigens on rheumatoid arthritis development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 19966-70	11.5	52

210	A genome-wide association study of rheumatoid arthritis without antibodies against citrullinated peptides. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, e15	2.4	49
209	Disease-modifying antirheumatic drug-free sustained remission in rheumatoid arthritis: an increasingly achievable outcome with subsidence of disease symptoms. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 867-73	2.4	48
208	MRI of hand and foot joints of patients with anticitrullinated peptide antibody positive arthralgia without clinical arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 1540-4	2.4	46
207	2017 EULAR recommendations for a core data set to support observational research and clinical care in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2018 , 77, 476-479	2.4	45
206	Improved treatment strategies reduce the increased mortality risk in early RA patients. <i>Rheumatology</i> , 2010 , 49, 2210-6	3.9	45
205	Evaluation of magnetic resonance imaging-detected tenosynovitis in the hand and wrist in early arthritis. <i>Arthritis and Rheumatology</i> , 2015 , 67, 869-76	9.5	44
204	Anticitrullinated protein antibodies and rheumatoid factor are associated with increased mortality but with different causes of death in patients with rheumatoid arthritis: a longitudinal study in three European cohorts. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1924-1932	2.4	44
203	ACPA fine-specificity profiles in early rheumatoid arthritis patients do not correlate with clinical features at baseline or with disease progression. <i>Arthritis Research and Therapy</i> , 2013 , 15, R140	5.7	44
202	Fatigue in rheumatoid arthritis; a persistent problem: a large longitudinal study. <i>RMD Open</i> , 2015 , 1, e000041	5.9	43
201	Diagnostic value of anti-MCV antibodies in differentiating early inflammatory arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 730-2	2.4	43
200	No influence of melatonin on cerebral blood flow in humans. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2003 , 88, 5989-94	5.6	43
199	Contribution of Fcγ3 receptor IIIA gene 158V/F polymorphism and copy number variation to the risk of ACPA-positive rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2009 , 68, 1775-80	2.4	42
198	Concordance between inflammation at physical examination and on MRI in patients with early arthritis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 506-12	2.4	41
197	Rheumatoid factor and anti-citrullinated protein antibody positivity, but not level, are associated with increased mortality in patients with rheumatoid arthritis: results from two large independent cohorts. <i>Arthritis Research and Therapy</i> , 2014 , 16, 483	5.7	41
196	A genetic variant in the region of MMP-9 is associated with serum levels and progression of joint damage in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1163-9	2.4	41
195	Identification of CXCL13 as a marker for rheumatoid arthritis outcome using an in silico model of the rheumatic joint. <i>Arthritis and Rheumatism</i> , 2011 , 63, 1265-73		41
194	Emerging patterns of risk factor make-up enable subclassification of rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2007 , 56, 1728-35		41
193	Genetic variants in IL15 associate with progression of joint destruction in rheumatoid arthritis: a multicohort study. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1651-7	2.4	40

192	Window of opportunity in rheumatoid arthritis - definitions and supporting evidence: from old to new perspectives. <i>RMD Open</i> , 2019 , 5, e000870	5.9	39
191	Distinct ACPA fine specificities, formed under the influence of HLA shared epitope alleles, have no effect on radiographic joint damage in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 1461-4	2.4	39
190	Performance of classification criteria for peripheral spondyloarthritis and psoriatic arthritis in the Leiden Early Arthritis cohort. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1366-9	2.4	38
189	The inflammatory disease-associated variants in IL12B and IL23R are not associated with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2008 , 58, 1877-81		38
188	Undifferentiated arthritis characteristics and outcomes when applying the 2010 and 1987 criteria for rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 238-41	2.4	37
187	Association of variants in IL2RA with progression of joint destruction in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2013 , 65, 1684-93		36
186	Genetic variants in the prediction of rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1694-6	2.4	36
185	Brief Report: IRF4 Newly Identified as a Common Susceptibility Locus for Systemic Sclerosis and Rheumatoid Arthritis in a Cross-Disease Meta-Analysis of Genome-Wide Association Studies. <i>Arthritis and Rheumatology</i> , 2016 , 68, 2338-44	9.5	35
184	Identification of a genetic variant for joint damage progression in autoantibody-positive rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 2038-46	2.4	34
183	Loss of metacarpal bone density predicts RA development in recent-onset arthritis. <i>Rheumatology</i> , 2012 , 51, 1037-41	3.9	34
182	The risk of individual autoantibodies, autoantibody combinations and levels for arthritis development in clinically suspect arthralgia. <i>Rheumatology</i> , 2017 , 56, 2145-2153	3.9	33
181	Low-avidity anticitrullinated protein antibodies (ACPA) are associated with a higher rate of joint destruction in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 270-6	2.4	33
180	The prognostic value of baseline erosions in undifferentiated arthritis. <i>Arthritis Research and Therapy</i> , 2009 , 11, R155	5.7	33
179	Smoking is associated with the concurrent presence of multiple autoantibodies in rheumatoid arthritis rather than with anti-citrullinated protein antibodies per se: a multicenter cohort study. <i>Arthritis Research and Therapy</i> , 2016 , 18, 285	5.7	33
178	Aiming for a simpler early arthritis MRI protocol: can Gd contrast administration be eliminated?. <i>European Radiology</i> , 2015 , 25, 1520-7	8	32
177	Variation in IL7R predisposes to sarcoid inflammation. <i>Genes and Immunity</i> , 2009 , 10, 647-53	4.4	32
176	Arthritis of the large joints - in particular, the knee - at first presentation is predictive for a high level of radiological destruction of the small joints in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2007 , 66, 646-50	2.4	32
175	Brief Report: Clinical Trials Aiming to Prevent Rheumatoid Arthritis Cannot Detect Prevention Without Adequate Risk Stratification: A Trial of Methotrexate Versus Placebo in Undifferentiated Arthritis as an Example. <i>Arthritis and Rheumatology</i> , 2017 , 69, 926-931	9.5	30

174	The Course of Bone Marrow Edema in Early Undifferentiated Arthritis and Rheumatoid Arthritis: A Longitudinal Magnetic Resonance Imaging Study at Bone Level. <i>Arthritis and Rheumatology</i> , 2016 , 68, 1080-8	9.5	30
173	Repair of joint erosions in rheumatoid arthritis: prevalence and patient characteristics in a large inception cohort. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 727-9	2.4	30
172	Immunoglobulin A multiple myeloma presenting with Henoch-Schönlein purpura associated with reduced sialylation of IgA1. <i>British Journal of Haematology</i> , 2003 , 122, 915-7	4.5	30
171	Aiming for a shorter rheumatoid arthritis MRI protocol: can contrast-enhanced MRI replace T2 for the detection of bone marrow oedema?. <i>European Radiology</i> , 2014 , 24, 2614-22	8	29
170	Toward a data-driven evaluation of the 2010 American College of Rheumatology/European League Against Rheumatism criteria for rheumatoid arthritis: is it sensible to look at levels of rheumatoid factor?. <i>Arthritis and Rheumatism</i> , 2011 , 63, 1190-9		29
169	Smoking as a risk factor for the radiological severity of rheumatoid arthritis: a study on six cohorts. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1384-7	2.4	28
168	Sustained drug-free remission in rheumatoid arthritis after DAS-driven or non-DAS-driven therapy: a comparison of two cohort studies. <i>Rheumatology</i> , 2012 , 51, 1120-8	3.9	28
167	Acute rheumatic fever and poststreptococcal reactive arthritis reconsidered. <i>Current Opinion in Rheumatology</i> , 2010 , 22, 437-42	5.3	28
166	Variation in radiologic joint destruction in rheumatoid arthritis differs between monozygotic and dizygotic twins and pairs of unrelated patients. <i>Arthritis and Rheumatism</i> , 2006 , 54, 2028-30		28
165	Relationship of multi-biomarker disease activity score and other risk factors with radiographic progression in an observational study of patients with rheumatoid arthritis. <i>Rheumatology</i> , 2016 , 55, 357-66	3.9	27
164	Comparison of methodologies for analysing the progression of joint destruction in rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2013 , 42, 182-9	1.9	27
163	Pretreatment serum levels of anti-cyclic citrullinated peptide antibodies are associated with the response to methotrexate in recent-onset arthritis. <i>Annals of the Rheumatic Diseases</i> , 2008 , 67, 1194-5	2.4	27
162	Aspects of early arthritis. Definition of disease states in early arthritis: remission versus minimal disease activity. <i>Arthritis Research and Therapy</i> , 2006 , 8, 216	5.7	27
161	MR signal intensity: staying on the bright side in MR image interpretation. <i>RMD Open</i> , 2018 , 4, e000728	5.9	27
160	MRI-detected subclinical joint inflammation is associated with radiographic progression. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 2034-7	2.4	26
159	Improved early identification of arthritis: evaluating the efficacy of Early Arthritis Recognition Clinics. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 1295-301	2.4	26
158	Association of the 6q23 region with the rate of joint destruction in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 567-70	2.4	26
157	Differences in the symptomatic phase preceding ACPA-positive and ACPA-negative RA: a longitudinal study in arthralgia during progression to clinical arthritis. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 1751-1754	2.4	25

156	Validation of the EULAR definition of arthralgia suspicious for progression to rheumatoid arthritis. <i>Rheumatology</i> , 2017 , 56, 2123-2128	3.9	25
155	Genetic variation in GREM1 is a risk factor for fibrosis in pulmonary sarcoidosis. <i>Tissue Antigens</i> , 2011 , 77, 112-7		25
154	Rheumatoid arthritis phenotype at presentation differs depending on the number of autoantibodies present. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 716-720	2.4	24
153	Subclinical inflammation on MRI of hand and foot of anticitrullinated peptide antibody-negative arthralgia patients at risk for rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2014 , 16, R92	5.7	22
152	Risk of rheumatoid arthritis development in patients with unclassified arthritis according to the 2010 ACR/EULAR criteria for rheumatoid arthritis. <i>Rheumatology</i> , 2013 , 52, 1265-70	3.9	22
151	Genetic variation of the Fc gamma receptor 3B gene and association with rheumatoid arthritis. <i>PLoS ONE</i> , 2010 , 5, e13173	3.7	22
150	Hypernatremia from a hunger strike as a cause of osmotic myelinolysis. <i>Neurology</i> , 2005 , 64, 574-5	6.5	22
149	Persistently activated, proliferative memory autoreactive B cells promote inflammation in rheumatoid arthritis. <i>Science Translational Medicine</i> , 2020 , 12,	17.5	22
148	Possibilities for preventive treatment in rheumatoid arthritis? Lessons from experimental animal models of arthritis: a systematic literature review and meta-analysis. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 458-467	2.4	21
147	What is the prevalence of MRI-detected inflammation and erosions in small joints in the general population? A collation and analysis of published data. <i>RMD Open</i> , 2015 , 1, e000005	5.9	21
146	Evaluation of the diagnostic accuracy of hand and foot MRI for early Rheumatoid Arthritis. <i>Rheumatology</i> , 2017 , 56, 1367-1377	3.9	20
145	Clinical expertise and its accuracy in differentiating arthralgia patients at risk for rheumatoid arthritis from other patients presenting with joint symptoms. <i>Rheumatology</i> , 2016 , 55, 1140-1	3.9	20
144	Studies on ageing and the severity of radiographic joint damage in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2015 , 17, 222	5.7	20
143	The 2010 ACR/EULAR criteria for rheumatoid arthritis: do they affect the classification or diagnosis of rheumatoid arthritis?. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1596-8	2.4	20
142	Long-term outcome of rheumatoid arthritis defined according to the 2010-classification criteria. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 428-32	2.4	19
141	Association of genetic variants in the IL4 and IL4R genes with the severity of joint damage in rheumatoid arthritis: a study in seven cohorts. <i>Arthritis and Rheumatism</i> , 2013 , 65, 3051-7		19
140	Studying associations between variants in TRAF1-C5 and TNFAIP3-OLIG3 and the progression of joint destruction in rheumatoid arthritis in multiple cohorts. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1753-5	2.4	19
139	Do musculoskeletal ultrasound and magnetic resonance imaging identify synovitis and tenosynovitis at the same joints and tendons? A comparative study in early inflammatory arthritis and clinically suspect arthralgia. <i>Arthritis Research and Therapy</i> , 2019 , 21, 59	5.7	18

138	Genetic Factors for the Severity of ACPA-negative Rheumatoid Arthritis in 2 Cohorts of Early Disease: A Genome-wide Study. <i>Journal of Rheumatology</i> , 2015 , 42, 1383-91	4.1	18
137	DMARD-free remission as novel treatment target in rheumatoid arthritis: A systematic literature review of achievability and sustainability. <i>RMD Open</i> , 2020 , 6,	5.9	18
136	Evaluating joint destruction in rheumatoid arthritis: is it necessary to radiograph both hands and feet?. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 345-9	2.4	18
135	The 2010 ACR/EULAR criteria are not sufficiently accurate in the early identification of autoantibody-negative rheumatoid arthritis: Results from the Leiden-EAC and ESPOIR cohorts. <i>Seminars in Arthritis and Rheumatism</i> , 2017 , 47, 170-174	5.3	17
134	The isotype and IgG subclass distribution of anti-carbamylated protein antibodies in rheumatoid arthritis patients. <i>Arthritis Research and Therapy</i> , 2017 , 19, 190	5.7	17
133	Are MRI-detected erosions specific for RA? A large explorative cross-sectional study. <i>Annals of the Rheumatic Diseases</i> , 2018 , 77, 861-868	2.4	17
132	Are rheumatoid arthritis patients discernible from other early arthritis patients using 1.5T extremity magnetic resonance imaging? a large cross-sectional study. <i>Journal of Rheumatology</i> , 2014 , 41, 1630-7	4.1	17
131	Allele-dose association of the C5orf30 rs26232 variant with joint damage in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2013 , 65, 2555-61		17
130	How do general practitioners identify inflammatory arthritis? A cohort analysis of Dutch general practitioner electronic medical records. <i>Rheumatology</i> , 2016 , 55, 848-53	3.9	17
129	Autoantibodies to two novel peptides in seronegative and early rheumatoid arthritis. <i>Rheumatology</i> , 2016 , 55, 1431-6	3.9	17
128	Magnetic resonance imaging-detected inflammation is associated with functional disability in early arthritis-results of a cross-sectional study. <i>Rheumatology</i> , 2016 , 55, 2167-2175	3.9	16
127	A genetic variant in osteoprotegerin is associated with progression of joint destruction in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2014 , 16, R108	5.7	16
126	An investigation of the added value of an ACPA multiplex assay in an early rheumatoid arthritis setting. <i>Arthritis Research and Therapy</i> , 2015 , 17, 276	5.7	16
125	Reasons for medical help-seeking behaviour of patients with recent-onset arthralgia. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 1302-7	2.4	16
124	Can anti-cyclic citrullinated peptide antibody-negative RA be subdivided into clinical subphenotypes?. <i>Arthritis Research and Therapy</i> , 2011 , 13, R180	5.7	16
123	TRAF1/C5 polymorphism is not associated with increased mortality in rheumatoid arthritis: two large longitudinal studies. <i>Arthritis Research and Therapy</i> , 2010 , 12, R38	5.7	16
122	Views on clinically suspect arthralgia: a focus group study. <i>Clinical Rheumatology</i> , 2016 , 35, 1347-52	3.9	15
121	Tapering towards DMARD-free remission in established rheumatoid arthritis: 2-year results of the TARA trial. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1174-1181	2.4	15

120	Anaemia to predict radiographic progression in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, e16	2.4	15
119	A genetic variant in granzyme B is associated with progression of joint destruction in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2013 , 65, 582-9		15
118	Defining erosive disease typical of RA in the light of the ACR/EULAR 2010 criteria for rheumatoid arthritis; results of the data driven phase. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 590-5	2.4	15
117	Associations between APOE genotypes and disease susceptibility, joint damage and lipid levels in patients with rheumatoid arthritis. <i>PLoS ONE</i> , 2013 , 8, e60970	3.7	15
116	Enhanced treatment strategies and distinct disease outcomes among autoantibody-positive and -negative rheumatoid arthritis patients over 25 years: A longitudinal cohort study in the Netherlands. <i>PLoS Medicine</i> , 2020 , 17, e1003296	11.6	15
115	Depression and anxiety associate with less remission after 1 year in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, e1	2.4	15
114	Association analysis of copy numbers of FC-gamma receptor genes for rheumatoid arthritis and other immune-mediated phenotypes. <i>European Journal of Human Genetics</i> , 2016 , 24, 263-70	5.3	14
113	ACPA-negative RA consists of subgroups: patients with high likelihood of achieving sustained DMARD-free remission can be identified by serological markers at disease presentation. <i>Arthritis Research and Therapy</i> , 2019 , 21, 121	5.7	14
112	Which patients presenting with arthralgia eventually develop rheumatoid arthritis? The current state of the art. <i>RMD Open</i> , 2017 , 3, e000479	5.9	14
111	IL2RA is associated with persistence of rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2015 , 17, 244	5.7	14
110	Serum pyridinoline levels and prediction of severity of joint destruction in rheumatoid arthritis. <i>Journal of Rheumatology</i> , 2013 , 40, 1303-6	4.1	14
109	Association of valine and leucine at HLA-DRB1 position 11 with radiographic progression in rheumatoid arthritis, independent of the shared epitope alleles but not independent of anti-citrullinated protein antibodies. <i>Arthritis and Rheumatology</i> , 2015 , 67, 877-86	9.5	13
108	The diagnostic accuracy of the squeeze test to identify arthritis: a cross-sectional cohort study. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1886-9	2.4	13
107	MRI-detected osteitis is not associated with the presence or level of ACPA alone, but with the combined presence of ACPA and RF. <i>Arthritis Research and Therapy</i> , 2016 , 18, 179	5.7	13
106	Moderate use of alcohol is associated with lower levels of C reactive protein but not with less severe joint inflammation: a cross-sectional study in early RA and healthy volunteers. <i>RMD Open</i> , 2018 , 4, e000577	5.9	12
105	Inflammatory genes TNF and IL6 display no signs of increased H3K4me3 in circulating monocytes from untreated rheumatoid arthritis patients. <i>Genes and Immunity</i> , 2017 , 18, 191-196	4.4	12
104	Evaluating processes underlying the predictive value of baseline erosions for future radiological damage in early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 883-9	2.4	12
103	Patients with early arthritis consume less alcohol than controls, regardless of the type of arthritis. <i>Rheumatology</i> , 2013 , 52, 1701-7	3.9	12

102	Role of NOD1 polymorphism in susceptibility and clinical progression of rheumatoid arthritis. <i>Rheumatology</i> , 2013 , 52, 806-14	3.9	12
101	How to avoid phenotypic misclassification in using joint destruction as an outcome measure for rheumatoid arthritis?. <i>Rheumatology</i> , 2010 , 49, 1429-35	3.9	12
100	Body mass index and extent of MRI-detected inflammation: opposite effects in rheumatoid arthritis versus other arthritides and asymptomatic persons. <i>Arthritis Research and Therapy</i> , 2016 , 18, 245	5.7	12
99	Older age is associated with more MRI-detected inflammation in hand and foot joints. <i>Rheumatology</i> , 2016 , 55, 2212-2219	3.9	12
98	The prevalence of ACPA is lower in rheumatoid arthritis patients with an older age of onset but the composition of the ACPA response appears identical. <i>Arthritis Research and Therapy</i> , 2017 , 19, 115	5.7	11
97	Longitudinal IP-10 Serum Levels Are Associated with the Course of Disease Activity and Remission in Patients with Rheumatoid Arthritis. <i>Vaccine Journal</i> , 2017 , 24,		11
96	Reappraisal of the diagnostic and prognostic value of morning stiffness in arthralgia and early arthritis: results from the Groningen EARC, Leiden EARC, ESPOIR, Leiden EAC and REACH. <i>Arthritis Research and Therapy</i> , 2015 , 17, 108	5.7	11
95	Are baseline high molecular weight adiponectin levels associated with radiographic progression in rheumatoid arthritis and osteoarthritis?. <i>Journal of Rheumatology</i> , 2014 , 41, 853-7	4.1	11
94	Predicting the severity of joint damage in rheumatoid arthritis; the contribution of genetic factors. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 876-82	2.4	10
93	PTGER4 gene variant rs76523431 is a candidate risk factor for radiological joint damage in rheumatoid arthritis patients: a genetic study of six cohorts. <i>Arthritis Research and Therapy</i> , 2015 , 17, 306	5.7	10
92	Automatic quantification of bone marrow edema on MRI of the wrist in patients with early arthritis: A feasibility study. <i>Magnetic Resonance in Medicine</i> , 2018 , 79, 1127-1134	4.4	9
91	Protective effect of HLA-DRB1*13 alleles during specific phases in the development of ACPA-positive RA. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1891-8	2.4	9
90	Does immunological remission, defined as disappearance of autoantibodies, occur with current treatment strategies? A long-term follow-up study in rheumatoid arthritis patients who achieved sustained DMARD-free status. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 1497-1504	2.4	9
89	Risk estimation in rheumatoid arthritis: from bench to bedside. <i>Nature Reviews Rheumatology</i> , 2014 , 10, 171-80	8.1	9
88	Differential methylation within the major histocompatibility complex region in rheumatoid arthritis: a replication study. <i>Rheumatology</i> , 2014 , 53, 2317-8	3.9	9
87	TREAT Early Arthralgia to Reverse or Limit Impending Exacerbation to Rheumatoid arthritis (TREAT EARLIER): a randomized, double-blind, placebo-controlled clinical trial protocol. <i>Trials</i> , 2020 , 21, 862	2.8	9
86	The anti-carbamylated protein antibody response is of overall low avidity despite extensive isotype switching. <i>Rheumatology</i> , 2018 , 57, 1583-1591	3.9	9
85	Imaging detected tenosynovitis of metacarpophalangeal and wrist joints: an increasingly recognised characteristic of rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 114, 131-138	2.2	9

84	Genetic associations with radiological damage in rheumatoid arthritis: Meta-analysis of seven genome-wide association studies of 2,775 cases. <i>PLoS ONE</i> , 2019 , 14, e0223246	3.7	8
83	Macrophage migration inhibitory factor (MIF) -173 polymorphism is associated with clinical erythema nodosum in LfgrnB syndrome. <i>Cytokine</i> , 2014 , 69, 272-6	4	8
82	Functional limitations in the phase of clinically suspect arthralgia are as serious as in early clinical arthritis; a longitudinal study. <i>RMD Open</i> , 2017 , 3, e000419	5.9	8
81	SPP1 rs9138 variant contributes to the severity of radiological damage in anti-citrullinated protein autoantibody-negative rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1840-3	2.4	8
80	A two-stage mixed-effects model approach for gene-set analyses in candidate gene studies. <i>Statistics in Medicine</i> , 2012 , 31, 1190-202	2.3	8
79	Evaluation of the contribution of cumulative levels of inflammation to the variance in joint destruction in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 307-8	2.4	8
78	Symptoms associated with inflammatory arthritis are common in the primary care population: results from the joint symptoms survey. <i>Rheumatology</i> , 2019 , 58, 2009-2014	3.9	7
77	The earlier, the better or the worse? Towards accurate management of patients with arthralgia at risk for RA. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 312-315	2.4	7
76	Does information on novel identified autoantibodies contribute to predicting the progression from undifferentiated arthritis to rheumatoid arthritis: a study on anti-CarP antibodies as an example. <i>Arthritis Research and Therapy</i> , 2018 , 20, 94	5.7	7
75	Evaluation of the joint distribution at disease presentation of patients with rheumatoid arthritis: a large study across continents. <i>RMD Open</i> , 2017 , 3, e000568	5.9	7
74	A new classification of HLA-DRB1 alleles based on acid-base properties of the amino acids located at positions 13, 70 and 71: impact on ACPA status or structural progression, and meta-analysis on 1235 patients with rheumatoid from two cohorts (ESPOIR and EAC cohort). <i>RMD Open</i> , 2015 , 1, e000099	5.9	7
73	Does the season at symptom onset influence the long-term severity of radiographic joint destruction in rheumatoid arthritis?. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 2055-6	2.4	7
72	A search to the target tissue in which RA-specific inflammation starts: a detailed MRI study to improve identification of RA-specific features in the phase of clinically suspect arthralgia. <i>Arthritis Research and Therapy</i> , 2019 , 21, 249	5.7	7
71	Improvement of symptoms in clinically suspect arthralgia and resolution of subclinical joint inflammation: a longitudinal study in patients that did not progress to clinical arthritis. <i>Arthritis Research and Therapy</i> , 2020 , 22, 11	5.7	6
70	The window of opportunity in ACPA-positive rheumatoid arthritis is not explained by ACPA characteristics. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 1697-8	2.4	6
69	Validity of the rheumatoid arthritis MRI score applied to the forefeet using the OMERACT filter: a systematic literature review. <i>RMD Open</i> , 2018 , 4, e000796	5.9	6
68	EULAR points to consider for conducting clinical trials and observational studies in individuals at risk of rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 1286-1298	2.4	6
67	Difficulties making a fist in clinically suspect arthralgia: an easy applicable phenomenon predictive for RA that is related to flexor tenosynovitis. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 1438-1439	2.4	5

66	Rheumatoid Arthritis and Tenosynovitis at the Metatarsophalangeal Joints: An Anatomic and MRI Study of the Forefoot Tendon Sheaths. <i>Radiology</i> , 2020 , 295, 146-154	20.5	5
65	The use of MRI-detected synovitis to determine the number of involved joints for the 2010 ACR/EULAR classification criteria for Rheumatoid Arthritis - is it of additional benefit?. <i>Annals of the Rheumatic Diseases</i> , 2018 , 77, 1125-1129	2.4	5
64	Evaluation of the association between anticarbamylated protein antibodies and the longitudinal course of functional ability in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, e14	2.4	5
63	Is joint pain in patients with arthralgia suspicious for progression to rheumatoid arthritis explained by subclinical inflammation? A cross-sectional MRI study. <i>Rheumatology</i> , 2019 , 58, 86-93	3.9	5
62	Does a genetic variant in FOXO3A predict a milder course of rheumatoid arthritis?. <i>Arthritis and Rheumatology</i> , 2014 , 66, 1678-81	9.5	5
61	Prediction of the Progression of Undifferentiated Arthritis to Rheumatoid Arthritis Using DNA Methylation Profiling. <i>Arthritis and Rheumatology</i> , 2021 , 73, 2229-2239	9.5	5
60	Automatic quantification of tenosynovitis on MRI of the wrist in patients with early arthritis: a feasibility study. <i>European Radiology</i> , 2019 , 29, 4477-4484	8	5
59	Reliability of Magnetic Resonance Imaging (MRI) Scoring of the Metatarsophalangeal Joints of the Foot according to the Rheumatoid Arthritis MRI Score. <i>Journal of Rheumatology</i> , 2020 , 47, 1165-1173	4.1	4
58	The relation between physical joint examination and MRI-depicted inflammation of metatarsophalangeal joints in early arthritis. <i>Arthritis Research and Therapy</i> , 2020 , 22, 67	5.7	4
57	Identifying MRI-detected inflammatory features specific for rheumatoid arthritis: two-fold feature reduction maintains predictive accuracy in clinically suspect arthralgia patients. <i>Seminars in Arthritis and Rheumatism</i> , 2019 , 48, 579-586	5.3	4
56	Is early remission associated with improved survival or is arthritis persistency associated with increased mortality in early arthritis? Comparisons with the general population. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, e25	2.4	4
55	The PTPN22 susceptibility risk variant is not associated with the rate of joint destruction in anti-citrullinated protein antibody-positive rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1730-1	2.4	4
54	Referring early arthritis patients within 6 weeks versus 12 weeks after symptom onset: an observational cohort study. <i>Lancet Rheumatology, The</i> , 2020 , 2, e332-e338	14.2	4
53	Not only synovitis but also tenosynovitis needs to be considered: why it is time to update textbook images of rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 546-547	2.4	4
52	Determining in which pre-arthritis stage HLA-shared epitope alleles and smoking exert their effect on the development of rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	4
51	Substitution of the quantitative serological component in the 2010 criteria for RA with qualitative presence of three autoantibodies yields similar performance: response to the article by Regueiro et al. <i>Arthritis Research and Therapy</i> , 2020 , 22, 85	5.7	4
50	Do autoantibody-responses mature between presentation with arthralgia suspicious for progression to rheumatoid arthritis and development of clinically apparent inflammatory arthritis? A longitudinal serological study. <i>Annals of the Rheumatic Diseases</i> , 2020 ,	2.4	4
49	Development and validation of a clinical rule for recognition of early inflammatory arthritis. <i>BMJ Open</i> , 2019 , 8, e023552	3	4

48	What is the value of musculoskeletal ultrasound in patients presenting with arthralgia to predict inflammatory arthritis development? A systematic literature review. <i>Arthritis Research and Therapy</i> , 2018 , 20, 228	5.7	4
47	Realising early recognition of arthritis in times of increased telemedicine: the value of patient-reported swollen joints. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 668-669	2.4	4
46	To what extent do autoantibodies help to identify high-risk patients in systemic sclerosis?. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 113, 109-117	2.2	4
45	Evaluation of the predictive accuracy of MRI-detected erosions in hand and foot joints in patients with undifferentiated arthritis. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 144-146	2.4	3
44	The additive value of magnetic resonance imaging for bone edema in predicting rheumatoid arthritis development in early undifferentiated arthritis: comment on the article by Duer-Jensen et al. <i>Arthritis and Rheumatism</i> , 2012 , 64, 321-2; author reply 322-3		3
43	Is the modified disease activity score superior to the disease activity score in early arthritis and rheumatoid arthritis? Comment on the article by Baker et al. <i>Arthritis and Rheumatology</i> , 2014 , 66, 3244-325	3.5	3
42	No impact of serum uric acid on the outcome of recent-onset arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1424-5	2.4	3
41	Gene expression identifies patients who develop inflammatory arthritis in a clinically suspect arthralgia cohort. <i>Arthritis Research and Therapy</i> , 2020 , 22, 266	5.7	3
40	EULAR/eumusc.net standards of care for rheumatoid arthritis: cross-sectional analyses of importance, level of implementation and care gaps experienced by patients and rheumatologists across 35 European countries. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1423-1431	2.4	3
39	SAT0029 THE IMMUNE-PATHOGENIC CHARACTERISTICS OF AUTOREACTIVE B CELLS AGAINST CITRULLINATED ANTIGENS IN RHEUMATOID ARTHRITIS 2019 ,		3
38	Early intensive treatment normalises excess mortality in ACPA-negative RA but not in ACPA-positive RA. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, e124	2.4	3
37	Improving the feasibility of MRI in clinically suspect arthralgia for prediction of rheumatoid arthritis by omitting scanning of the feet. <i>Rheumatology</i> , 2020 , 59, 1247-1252	3.9	3
36	Earlier chronotype in patients with rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2021 , 40, 2185-2192	3.9	3
35	Increasing incidence of autoantibody-negative RA is replicated and is partly explained by an aging population. <i>Annals of the Rheumatic Diseases</i> , 2020 ,	2.4	2
34	Does psychological stress in patients with clinically suspect arthralgia associate with subclinical inflammation and progression to inflammatory arthritis?. <i>Arthritis Research and Therapy</i> , 2018 , 20, 93	5.7	2
33	Early identification of rheumatoid arthritis: the risk of overtreatment in perspective. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, e107	2.4	2
32	How to define baseline erosiveness to predict radiologic progression in rheumatoid arthritis. <i>Journal of Rheumatology</i> , 2013 , 40, 1624-5	4.1	2
31	Biomarkers in the prediction of radiographic progression in early rheumatoid arthritis: comment on the article by Young-Min et al. <i>Arthritis and Rheumatism</i> , 2008 , 58, 1558-9; author reply 1559		2

30	Differing time-orders of inflammation decrease between ACPA subsets in RA patients suggest differences in underlying inflammatory pathways. <i>Rheumatology</i> , 2021 , 60, 2969-2975	3.9	2
29	Two-year cost effectiveness between two gradual tapering strategies in rheumatoid arthritis: cost-utility analysis of the TARA trial. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1550-1556	2.4	2
28	Screening for two or three autoantibodies in persons at risk for RA: implications of current data for clinical practice. <i>Rheumatology</i> , 2019 , 58, 914-915	3.9	2
27	The value of the squeeze test for detection of subclinical synovitis in patients with arthralgia suspicious for progression to RA. <i>Rheumatology</i> , 2020 , 59, 3106-3108	3.9	1
26	Osteoprotegerin as biomarker for persistence of rheumatoid arthritis. <i>Rheumatology</i> , 2016 , 55, 949-50	3.9	1
25	Treat-to-Target Strategies in Rheumatoid Arthritis. <i>JAMA - Journal of the American Medical Association</i> , 2019 , 322, 83	27.4	1
24	Is HLA-B27 increased in patients diagnosed with undifferentiated arthritis? Results from the Leiden early arthritis cohort. <i>Journal of Rheumatology</i> , 2014 , 41, 1948-51	4.1	1
23	Diagnostic value of performing chest x-rays in patients presenting with recent-onset arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 627-8	2.4	1
22	Subclinical synovitis in arthralgia: how often does it result in clinical arthritis? Reflecting on starting points for disease-modifying anti-rheumatic drug treatment. <i>Rheumatology</i> , 2021 , 60, 3872-3878	3.9	1
21	Value of imaging detected joint inflammation in explaining fatigue in RA at diagnosis and during the disease course: a large MRI study. <i>RMD Open</i> , 2021 , 7,	5.9	1
20	Correspondence on Role of joint damage, malalignment and inflammation in articular tenderness in rheumatoid arthritis, psoriatic arthritis and osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	1
19	Tenosynovitis has a high sensitivity for early ACPA-positive and ACPA-negative RA: a large cross-sectional MRI study. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	1
18	Development of clinically apparent synovitis: a longitudinal study at the joint level during progression to inflammatory arthritis. <i>RMD Open</i> , 2018 , 4, e000748	5.9	1
17	Morning stiffness precedes the development of RA and associates with systemic and subclinical joint inflammation in arthralgia patients. <i>Rheumatology</i> , 2021 ,	3.9	1
16	A Polymorphism in C-C Chemokine Receptor 5 (CCR5) Associates with LÉgrenB Syndrome and Alters Receptor Expression as well as Functional Response. <i>Cells</i> , 2021 , 10,	7.9	1
15	Factors that influence biological survival in rheumatoid arthritis: results of a real-world academic cohort from the Netherlands. <i>Clinical Rheumatology</i> , 2021 , 40, 2177-2183	3.9	1
14	Methotrexate prolongs TNF inhibitor survival, but only in autoantibody-positive rheumatoid arthritis: a validation study. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	1
13	The eumusc.net standards of care for rheumatoid arthritis: importance and current implementation according to patients and healthcare providers in the Netherlands. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36, 275-283	2.2	1

12	Response to: "Does ACPA-negative RA consist of subgroups related to sustained DMARD-free remission and serological markers at disease presentation" by Masi and Fleischmann. <i>Arthritis Research and Therapy</i> , 2020 , 22, 55	5.7	○
11	Is shoulder involvement in clinically suspect arthralgia an early feature of rheumatoid arthritis? A longitudinal ultrasound study. <i>Rheumatology</i> , 2020 , 59, 2640-2642	3.9	○
10	Increased frequency of intermetatarsal and submetatarsal bursitis in early rheumatoid arthritis: a large case-controlled MRI study. <i>Arthritis Research and Therapy</i> , 2020 , 22, 277	5.7	○
9	The target of ACPA. <i>Rheumatology</i> , 2016 , 55, 1711-3	3.9	○
8	Impaired fertility in men diagnosed with inflammatory arthritis: results of a large multicentre study (iFAME-Fertility). <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 1545-1552	2.4	○
7	Prospective Studies on the Risk of Rheumatoid Arthritis: The European Risk RA Registry.. <i>Frontiers in Medicine</i> , 2022 , 9, 824501	4.9	○
6	ACPA-negative and ACPA-positive RA-patients achieving disease resolution demonstrate distinct patterns of MRI-detected joint-inflammation.. <i>Rheumatology</i> , 2022 ,	3.9	○
5	A1.29 In rheumatoid arthritis, smoking is not primarily associated with anti-citrullinaged protein antibodies, but with the presence of several autoantibodies. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, A12.1-A12	2.4	
4	Distinct ACPA fine-specificities, formed under the influence of HLA shared epitope alleles, have no effect on radiographic joint damage in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, A5-A5	2.4	
3	Association of tumor necrosis factor alpha polymorphism and radiographic progression in rheumatoid arthritis: comment on the article by Khanna et al. <i>Arthritis and Rheumatism</i> , 2007 , 56, 1032-3; author reply1033-4		
2	The value of inquiring about functional impairments for early identification of inflammatory arthritis: a large cross-sectional derivation and validation study from the Netherlands. <i>BMJ Open</i> , 2020 , 10, e040148	3	
1	Vroegtijdige herkenning van artritis met een scorelijst. <i>Huisarts En Wetenschap</i> , 2020 , 63, 14-18	0.1	