

Annette H Van Der Helm-Van Mil

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1711536/annette-h-van-der-helm-van-mil-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

263
papers

11,856
citations

52
h-index

101
g-index

293
ext. papers

14,026
ext. citations

4.8
avg. IF

6.23
L-index

#	Paper	IF	Citations
263	Prospective Studies on the Risk of Rheumatoid Arthritis: The European Risk RA Registry.. <i>Frontiers in Medicine</i> , 2022 , 9, 824501	4.9	0
262	ACPA-negative and ACPA-positive RA-patients achieving disease resolution demonstrate distinct patterns of MRI-detected joint-inflammation.. <i>Rheumatology</i> , 2022 ,	3.9	0
261	Subclinical synovitis in arthralgia: how often does it result in clinical arthritis? Reflecting on starting points for disease-modifying anti-rheumatic drug treatment. <i>Rheumatology</i> , 2021 , 60, 3872-3878	3.9	1
260	Differing time-orders of inflammation decrease between ACPA subsets in RA patients suggest differences in underlying inflammatory pathways. <i>Rheumatology</i> , 2021 , 60, 2969-2975	3.9	2
259	Prediction of the Progression of Undifferentiated Arthritis to Rheumatoid Arthritis Using DNA Methylation Profiling. <i>Arthritis and Rheumatology</i> , 2021 , 73, 2229-2239	9.5	5
258	Value of imaging detected joint inflammation in explaining fatigue in RA at diagnosis and during the disease course: a large MRI study. <i>RMD Open</i> , 2021 , 7,	5.9	1
257	Correspondence on Role of joint damage, malalignment and inflammation in articular tenderness in rheumatoid arthritis, psoriatic arthritis and osteoarthritis? <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	1
256	Determining in which pre-arthritis stage HLA-shared epitope alleles and smoking exert their effect on the development of rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	4
255	Earlier chronotype in patients with rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2021 , 40, 2185-2192	3.9	3
254	Tenosynovitis has a high sensitivity for early ACPA-positive and ACPA-negative RA: a large cross-sectional MRI study. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	1
253	Impaired fertility in men diagnosed with inflammatory arthritis: results of a large multicentre study (iFAME-Fertility). <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 1545-1552	2.4	0
252	Morning stiffness precedes the development of RA and associates with systemic and subclinical joint inflammation in arthralgia patients. <i>Rheumatology</i> , 2021 ,	3.9	1
251	A Polymorphism in C-C Chemokine Receptor 5 (CCR5) Associates with LfgrnB Syndrome and Alters Receptor Expression as well as Functional Response. <i>Cells</i> , 2021 , 10,	7.9	1
250	EULAR points to consider for conducting clinical trials and observational studies in individuals at risk of rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 1286-1298	2.4	6
249	Factors that influence biological survival in rheumatoid arthritis: results of a real-world academic cohort from the Netherlands. <i>Clinical Rheumatology</i> , 2021 , 40, 2177-2183	3.9	1
248	Realising early recognition of arthritis in times of increased telemedicine: the value of patient-reported swollen joints. <i>Annals of the Rheumatic Diseases</i> , 2021 , 80, 668-669	2.4	4
247	Methotrexate prolongs TNF inhibitor survival, but only in autoantibody-positive rheumatoid arthritis: a validation study. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	1

246	DMARD-free remission as novel treatment target in rheumatoid arthritis: A systematic literature review of achievability and sustainability. <i>RMD Open</i> , 2020 , 6,	5.9	18
245	Reliability of Magnetic Resonance Imaging (MRI) Scoring of the Metatarsophalangeal Joints of the Foot according to the Rheumatoid Arthritis MRI Score. <i>Journal of Rheumatology</i> , 2020 , 47, 1165-1173	4.1	4
244	Increasing incidence of autoantibody-negative RA is replicated and is partly explained by an aging population. <i>Annals of the Rheumatic Diseases</i> , 2020 ,	2.4	2
243	Tapering towards DMARD-free remission in established rheumatoid arthritis: 2-year results of the TARA trial. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1174-1181	2.4	15
242	The value of the squeeze test for detection of subclinical synovitis in patients with arthralgia suspicious for progression to RA. <i>Rheumatology</i> , 2020 , 59, 3106-3108	3.9	1
241	Response to: "Does ACPA-negative RA consist of subgroups related to sustained DMARD-free remission and serological markers at disease presentation" by Masi and Fleischmann. <i>Arthritis Research and Therapy</i> , 2020 , 22, 55	5.7	0
240	Is shoulder involvement in clinically suspect arthralgia an early feature of rheumatoid arthritis? A longitudinal ultrasound study. <i>Rheumatology</i> , 2020 , 59, 2640-2642	3.9	0
239	Rheumatoid Arthritis and Tenosynovitis at the Metatarsophalangeal Joints: An Anatomic and MRI Study of the Forefoot Tendon Sheaths. <i>Radiology</i> , 2020 , 295, 146-154	20.5	5
238	The earlier, the better or the worse? Towards accurate management of patients with arthralgia at risk for RA. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 312-315	2.4	7
237	Improvement of symptoms in clinically suspect arthralgia and resolution of subclinical joint inflammation: a longitudinal study in patients that did not progress to clinical arthritis. <i>Arthritis Research and Therapy</i> , 2020 , 22, 11	5.7	6
236	The relation between physical joint examination and MRI-depicted inflammation of metatarsophalangeal joints in early arthritis. <i>Arthritis Research and Therapy</i> , 2020 , 22, 67	5.7	4
235	The value of inquiring about functional impairments for early identification of inflammatory arthritis: a large cross-sectional derivation and validation study from the Netherlands. <i>BMJ Open</i> , 2020 , 10, e040148	3	
234	Referring early arthritis patients within 6 weeks versus 12 weeks after symptom onset: an observational cohort study. <i>Lancet Rheumatology, The</i> , 2020 , 2, e332-e338	14.2	4
233	Vroegtijdige herkenning van artritis met een scorelijst. <i>Huisarts En Wetenschap</i> , 2020 , 63, 14-18	0.1	
232	Not only synovitis but also tenosynovitis needs to be considered: why it is time to update textbook images of rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 546-547	2.4	4
231	TREAT Early Arthralgia to Reverse or Limit Impending Exacerbation to Rheumatoid arthritis (TREAT EARLIER): a randomized, double-blind, placebo-controlled clinical trial protocol. <i>Trials</i> , 2020 , 21, 862	2.8	9
230	Persistently activated, proliferative memory autoreactive B cells promote inflammation in rheumatoid arthritis. <i>Science Translational Medicine</i> , 2020 , 12,	17.5	22
229	Gene expression identifies patients who develop inflammatory arthritis in a clinically suspect arthralgia cohort. <i>Arthritis Research and Therapy</i> , 2020 , 22, 266	5.7	3

228	Increased frequency of intermetatarsal and submetatarsal bursitis in early rheumatoid arthritis: a large case-controlled MRI study. <i>Arthritis Research and Therapy</i> , 2020 , 22, 277	5.7	0
227	EULAR/eumusc.net standards of care for rheumatoid arthritis: cross-sectional analyses of importance, level of implementation and care gaps experienced by patients and rheumatologists across 35 European countries. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1423-1431	2.4	3
226	Enhanced treatment strategies and distinct disease outcomes among autoantibody-positive and -negative rheumatoid arthritis patients over 25 years: A longitudinal cohort study in the Netherlands. <i>PLoS Medicine</i> , 2020 , 17, e1003296	11.6	15
225	Two-year cost effectiveness between two gradual tapering strategies in rheumatoid arthritis: cost-utility analysis of the TARA trial. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 1550-1556	2.4	2
224	Early intensive treatment normalises excess mortality in ACPA-negative RA but not in ACPA-positive RA. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, e124	2.4	3
223	Improving the feasibility of MRI in clinically suspect arthralgia for prediction of rheumatoid arthritis by omitting scanning of the feet. <i>Rheumatology</i> , 2020 , 59, 1247-1252	3.9	3
222	Substitution of the quantitative serological component in the 2010 criteria for RA with qualitative presence of three autoantibodies yields similar performance: response to the article by Regueiro et al. <i>Arthritis Research and Therapy</i> , 2020 , 22, 85	5.7	4
221	Do autoantibody-responses mature between presentation with arthralgia suspicious for progression to rheumatoid arthritis and development of clinically apparent inflammatory arthritis? A longitudinal serological study. <i>Annals of the Rheumatic Diseases</i> , 2020 ,	2.4	4
220	Molecular Portraits of Early Rheumatoid Arthritis Identify Clinical and Treatment Response Phenotypes. <i>Cell Reports</i> , 2019 , 28, 2455-2470.e5	10.6	100
219	Window of opportunity in rheumatoid arthritis - definitions and supporting evidence: from old to new perspectives. <i>RMD Open</i> , 2019 , 5, e000870	5.9	39
218	Symptoms associated with inflammatory arthritis are common in the primary care population: results from the joint symptoms survey. <i>Rheumatology</i> , 2019 , 58, 2009-2014	3.9	7
217	ACPA-negative RA consists of subgroups: patients with high likelihood of achieving sustained DMARD-free remission can be identified by serological markers at disease presentation. <i>Arthritis Research and Therapy</i> , 2019 , 21, 121	5.7	14
216	Difficulties making a fist in clinically suspect arthralgia: an easy applicable phenomenon predictive for RA that is related to flexor tenosynovitis. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 1438-1439	2.4	5
215	Synovial cellular and molecular signatures stratify clinical response to csDMARD therapy and predict radiographic progression in early rheumatoid arthritis patients. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 761-772	2.4	96
214	Do musculoskeletal ultrasound and magnetic resonance imaging identify synovitis and tenosynovitis at the same joints and tendons? A comparative study in early inflammatory arthritis and clinically suspect arthralgia. <i>Arthritis Research and Therapy</i> , 2019 , 21, 59	5.7	18
213	Evaluation of the predictive accuracy of MRI-detected erosions in hand and foot joints in patients with undifferentiated arthritis. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 144-146	2.4	3
212	Is joint pain in patients with arthralgia suspicious for progression to rheumatoid arthritis explained by subclinical inflammation? A cross-sectional MRI study. <i>Rheumatology</i> , 2019 , 58, 86-93	3.9	5
211	Early identification of rheumatoid arthritis: the risk of overtreatment in perspective. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, e107	2.4	2

210	Identifying MRI-detected inflammatory features specific for rheumatoid arthritis: two-fold feature reduction maintains predictive accuracy in clinically suspect arthralgia patients. <i>Seminars in Arthritis and Rheumatism</i> , 2019 , 48, 579-586	5.3	4
209	Does immunological remission, defined as disappearance of autoantibodies, occur with current treatment strategies? A long-term follow-up study in rheumatoid arthritis patients who achieved sustained DMARD-free status. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 1497-1504	2.4	9
208	Treat-to-Target Strategies in Rheumatoid Arthritis. <i>JAMA - Journal of the American Medical Association</i> , 2019 , 322, 83	27.4	1
207	Genetic associations with radiological damage in rheumatoid arthritis: Meta-analysis of seven genome-wide association studies of 2,775 cases. <i>PLoS ONE</i> , 2019 , 14, e0223246	3.7	8
206	Screening for two or three autoantibodies in persons at risk for RA: implications of current data for clinical practice. <i>Rheumatology</i> , 2019 , 58, 914-915	3.9	2
205	SAT0029 THE IMMUNE-PATHOGENIC CHARACTERISTICS OF AUTOREACTIVE B CELLS AGAINST CITRULLINATED ANTIGENS IN RHEUMATOID ARTHRITIS 2019 ,		3
204	A search to the target tissue in which RA-specific inflammation starts: a detailed MRI study to improve identification of RA-specific features in the phase of clinically suspect arthralgia. <i>Arthritis Research and Therapy</i> , 2019 , 21, 249	5.7	7
203	Automatic quantification of tenosynovitis on MRI of the wrist in patients with early arthritis: a feasibility study. <i>European Radiology</i> , 2019 , 29, 4477-4484	8	5
202	Depression and anxiety associate with less remission after 1 year in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, e1	2.4	15
201	Development and validation of a clinical rule for recognition of early inflammatory arthritis. <i>BMJ Open</i> , 2019 , 8, e023552	3	4
200	Are MRI-detected erosions specific for RA? A large explorative cross-sectional study. <i>Annals of the Rheumatic Diseases</i> , 2018 , 77, 861-868	2.4	17
199	The use of MRI-detected synovitis to determine the number of involved joints for the 2010 ACR/EULAR classification criteria for Rheumatoid Arthritis - is it of additional benefit?. <i>Annals of the Rheumatic Diseases</i> , 2018 , 77, 1125-1129	2.4	5
198	Moderate use of alcohol is associated with lower levels of C reactive protein but not with less severe joint inflammation: a cross-sectional study in early RA and healthy volunteers. <i>RMD Open</i> , 2018 , 4, e000577	5.9	12
197	2017 EULAR recommendations for a core data set to support observational research and clinical care in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2018 , 77, 476-479	2.4	45
196	Automatic quantification of bone marrow edema on MRI of the wrist in patients with early arthritis: A feasibility study. <i>Magnetic Resonance in Medicine</i> , 2018 , 79, 1127-1134	4.4	9
195	Does psychological stress in patients with clinically suspect arthralgia associate with subclinical inflammation and progression to inflammatory arthritis?. <i>Arthritis Research and Therapy</i> , 2018 , 20, 93	5.7	2
194	Does information on novel identified autoantibodies contribute to predicting the progression from undifferentiated arthritis to rheumatoid arthritis: a study on anti-CarP antibodies as an example. <i>Arthritis Research and Therapy</i> , 2018 , 20, 94	5.7	7
193	Preventing progression from arthralgia to arthritis: targeting the right patients. <i>Nature Reviews Rheumatology</i> , 2018 , 14, 32-41	8.1	59

192	Validity of the rheumatoid arthritis MRI score applied to the forefeet using the OMERACT filter: a systematic literature review. <i>RMD Open</i> , 2018 , 4, e000796	5.9	6
191	Update on the epidemiology, risk factors, and disease outcomes of rheumatoid arthritis. <i>Best Practice and Research in Clinical Rheumatology</i> , 2018 , 32, 174-187	5.3	120
190	Development of clinically apparent synovitis: a longitudinal study at the joint level during progression to inflammatory arthritis. <i>RMD Open</i> , 2018 , 4, e000748	5.9	1
189	What is the value of musculoskeletal ultrasound in patients presenting with arthralgia to predict inflammatory arthritis development? A systematic literature review. <i>Arthritis Research and Therapy</i> , 2018 , 20, 228	5.7	4
188	Triple Positivity for Anti-Citrullinated Protein Autoantibodies, Rheumatoid Factor, and Anti-Carbamylated Protein Antibodies Conferring High Specificity for Rheumatoid Arthritis: Implications for Very Early Identification of At-Risk Individuals. <i>Arthritis and Rheumatology</i> , 2018 , 70, 1721-1731	9.5	55
187	The anti-carbamylated protein antibody response is of overall low avidity despite extensive isotype switching. <i>Rheumatology</i> , 2018 , 57, 1583-1591	3.9	9
186	MR signal intensity: staying on the bright side in MR image interpretation. <i>RMD Open</i> , 2018 , 4, e000728	5.9	27
185	The eumusc.net standards of care for rheumatoid arthritis: importance and current implementation according to patients and healthcare providers in the Netherlands. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36, 275-283	2.2	1
184	To what extent do autoantibodies help to identify high-risk patients in systemic sclerosis?. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 113, 109-117	2.2	4
183	Imaging detected tenosynovitis of metacarpophalangeal and wrist joints: an increasingly recognised characteristic of rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 114, 131-138	2.2	9
182	The association between anti-carbamylated protein (anti-CarP) antibodies and radiographic progression in early rheumatoid arthritis: a study exploring replication and the added value to ACPA and rheumatoid factor. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 112-118	2.4	53
181	Rheumatoid arthritis phenotype at presentation differs depending on the number of autoantibodies present. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 716-720	2.4	24
180	Brief Report: Clinical Trials Aiming to Prevent Rheumatoid Arthritis Cannot Detect Prevention Without Adequate Risk Stratification: A Trial of Methotrexate Versus Placebo in Undifferentiated Arthritis as an Example. <i>Arthritis and Rheumatology</i> , 2017 , 69, 926-931	9.5	30
179	2016 update of the EULAR recommendations for the management of early arthritis. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 948-959	2.4	243
178	Possibilities for preventive treatment in rheumatoid arthritis? Lessons from experimental animal models of arthritis: a systematic literature review and meta-analysis. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 458-467	2.4	21
177	EULAR definition of arthralgia suspicious for progression to rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 491-496	2.4	125
176	The 2010 ACR/EULAR criteria are not sufficiently accurate in the early identification of autoantibody-negative rheumatoid arthritis: Results from the Leiden-EAC and ESPOIR cohorts. <i>Seminars in Arthritis and Rheumatism</i> , 2017 , 47, 170-174	5.3	17
175	Evaluation of the diagnostic accuracy of hand and foot MRI for early Rheumatoid Arthritis. <i>Rheumatology</i> , 2017 , 56, 1367-1377	3.9	20

174	The prevalence of ACPA is lower in rheumatoid arthritis patients with an older age of onset but the composition of the ACPA response appears identical. <i>Arthritis Research and Therapy</i> , 2017 , 19, 115	5.7	11
173	Longitudinal IP-10 Serum Levels Are Associated with the Course of Disease Activity and Remission in Patients with Rheumatoid Arthritis. <i>Vaccine Journal</i> , 2017 , 24,		11
172	Differences in the symptomatic phase preceding ACPA-positive and ACPA-negative RA: a longitudinal study in arthralgia during progression to clinical arthritis. <i>Annals of the Rheumatic Diseases</i> , 2017 , 76, 1751-1754	2.4	25
171	The risk of individual autoantibodies, autoantibody combinations and levels for arthritis development in clinically suspect arthralgia. <i>Rheumatology</i> , 2017 , 56, 2145-2153	3.9	33
170	The isotype and IgG subclass distribution of anti-carbamylated protein antibodies in rheumatoid arthritis patients. <i>Arthritis Research and Therapy</i> , 2017 , 19, 190	5.7	17
169	Inflammatory genes TNF α and IL6 display no signs of increased H3K4me3 in circulating monocytes from untreated rheumatoid arthritis patients. <i>Genes and Immunity</i> , 2017 , 18, 191-196	4.4	12
168	Evaluation of the joint distribution at disease presentation of patients with rheumatoid arthritis: a large study across continents. <i>RMD Open</i> , 2017 , 3, e000568	5.9	7
167	Functional limitations in the phase of clinically suspect arthralgia are as serious as in early clinical arthritis; a longitudinal study. <i>RMD Open</i> , 2017 , 3, e000419	5.9	8
166	Which patients presenting with arthralgia eventually develop rheumatoid arthritis? The current state of the art. <i>RMD Open</i> , 2017 , 3, e000479	5.9	14
165	Validation of the EULAR definition of arthralgia suspicious for progression to rheumatoid arthritis. <i>Rheumatology</i> , 2017 , 56, 2123-2128	3.9	25
164	Views on clinically suspect arthralgia: a focus group study. <i>Clinical Rheumatology</i> , 2016 , 35, 1347-52	3.9	15
163	Association analysis of copy numbers of FC-gamma receptor genes for rheumatoid arthritis and other immune-mediated phenotypes. <i>European Journal of Human Genetics</i> , 2016 , 24, 263-70	5.3	14
162	Strategies to predict rheumatoid arthritis development in at-risk populations. <i>Rheumatology</i> , 2016 , 55, 6-15	3.9	63
161	Magnetic resonance imaging-detected inflammation is associated with functional disability in early arthritis-results of a cross-sectional study. <i>Rheumatology</i> , 2016 , 55, 2167-2175	3.9	16
160	Brief Report: IRF4 Newly Identified as a Common Susceptibility Locus for Systemic Sclerosis and Rheumatoid Arthritis in a Cross-Disease Meta-Analysis of Genome-Wide Association Studies. <i>Arthritis and Rheumatology</i> , 2016 , 68, 2338-44	9.5	35
159	Disease-modifying antirheumatic drug-free sustained remission in rheumatoid arthritis: an increasingly achievable outcome with subsidence of disease symptoms. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 867-73	2.4	48
158	Anticitrullinated protein antibodies and rheumatoid factor are associated with increased mortality but with different causes of death in patients with rheumatoid arthritis: a longitudinal study in three European cohorts. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1924-1932	2.4	44
157	Clinical expertise and its accuracy in differentiating arthralgia patients at risk for rheumatoid arthritis from other patients presenting with joint symptoms. <i>Rheumatology</i> , 2016 , 55, 1140-1	3.9	20

156	Relationship of multi-biomarker disease activity score and other risk factors with radiographic progression in an observational study of patients with rheumatoid arthritis. <i>Rheumatology</i> , 2016 , 55, 357-66	3.9	27
155	Osteoprotegerin as biomarker for persistence of rheumatoid arthritis. <i>Rheumatology</i> , 2016 , 55, 949-50	3.9	1
154	Evaluation of the association between anticarbamylated protein antibodies and the longitudinal course of functional ability in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, e14	2.4	5
153	Protective effect of HLA-DRB1*13 alleles during specific phases in the development of ACPA-positive RA. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1891-8	2.4	9
152	Clinical factors, anticitrullinated peptide antibodies and MRI-detected subclinical inflammation in relation to progression from clinically suspect arthralgia to arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016 , 75, 1824-30	2.4	85
151	The Course of Bone Marrow Edema in Early Undifferentiated Arthritis and Rheumatoid Arthritis: A Longitudinal Magnetic Resonance Imaging Study at Bone Level. <i>Arthritis and Rheumatology</i> , 2016 , 68, 1080-8	9.5	30
150	The target of ACPA. <i>Rheumatology</i> , 2016 , 55, 1711-3	3.9	0
149	Smoking is associated with the concurrent presence of multiple autoantibodies in rheumatoid arthritis rather than with anti-citrullinated protein antibodies per se: a multicenter cohort study. <i>Arthritis Research and Therapy</i> , 2016 , 18, 285	5.7	33
148	MRI-detected osteitis is not associated with the presence or level of ACPA alone, but with the combined presence of ACPA and RF. <i>Arthritis Research and Therapy</i> , 2016 , 18, 179	5.7	13
147	Body mass index and extent of MRI-detected inflammation: opposite effects in rheumatoid arthritis versus other arthritides and asymptomatic persons. <i>Arthritis Research and Therapy</i> , 2016 , 18, 245	5.7	12
146	How do general practitioners identify inflammatory arthritis? A cohort analysis of Dutch general practitioner electronic medical records. <i>Rheumatology</i> , 2016 , 55, 848-53	3.9	17
145	Autoantibodies to two novel peptides in seronegative and early rheumatoid arthritis. <i>Rheumatology</i> , 2016 , 55, 1431-6	3.9	17
144	Older age is associated with more MRI-detected inflammation in hand and foot joints. <i>Rheumatology</i> , 2016 , 55, 2212-2219	3.9	12
143	A genetic study on C5-TRAF1 and progression of joint damage in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2015 , 17, 1	5.7	70
142	Aiming for a simpler early arthritis MRI protocol: can Gd contrast administration be eliminated?. <i>European Radiology</i> , 2015 , 25, 1520-7	8	32
141	A genome-wide association study of rheumatoid arthritis without antibodies against citrullinated peptides. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, e15	2.4	49
140	Genetic Factors for the Severity of ACPA-negative Rheumatoid Arthritis in 2 Cohorts of Early Disease: A Genome-wide Study. <i>Journal of Rheumatology</i> , 2015 , 42, 1383-91	4.1	18
139	Association of valine and leucine at HLA-DRB1 position 11 with radiographic progression in rheumatoid arthritis, independent of the shared epitope alleles but not independent of anti-citrullinated protein antibodies. <i>Arthritis and Rheumatology</i> , 2015 , 67, 877-86	9.5	13

138	The diagnostic accuracy of the squeeze test to identify arthritis: a cross-sectional cohort study. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1886-9	2.4	13
137	Evaluation of magnetic resonance imaging-detected tenosynovitis in the hand and wrist in early arthritis. <i>Arthritis and Rheumatology</i> , 2015 , 67, 869-76	9.5	44
136	Fatigue in rheumatoid arthritis; a persistent problem: a large longitudinal study. <i>RMD Open</i> , 2015 , 1, e000041	5.9	43
135	Predicting the severity of joint damage in rheumatoid arthritis; the contribution of genetic factors. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 876-82	2.4	10
134	What is the prevalence of MRI-detected inflammation and erosions in small joints in the general population? A collation and analysis of published data. <i>RMD Open</i> , 2015 , 1, e000005	5.9	21
133	A new classification of HLA-DRB1 alleles based on acid-base properties of the amino acids located at positions 13, 70 and 71: impact on ACPA status or structural progression, and meta-analysis on 1235 patients with rheumatoid from two cohorts (ESPOIR and EAC cohort). <i>RMD Open</i> , 2015 , 1, e000099	5.9	7
132	Studies on ageing and the severity of radiographic joint damage in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2015 , 17, 222	5.7	20
131	The specificity of anti-carbamylated protein antibodies for rheumatoid arthritis in a setting of early arthritis. <i>Arthritis Research and Therapy</i> , 2015 , 17, 339	5.7	55
130	PTGER4 gene variant rs76523431 is a candidate risk factor for radiological joint damage in rheumatoid arthritis patients: a genetic study of six cohorts. <i>Arthritis Research and Therapy</i> , 2015 , 17, 306	5.7	10
129	Reappraisal of the diagnostic and prognostic value of morning stiffness in arthralgia and early arthritis: results from the Groningen EARC, Leiden EARC, ESPOIR, Leiden EAC and REACH. <i>Arthritis Research and Therapy</i> , 2015 , 17, 108	5.7	11
128	An investigation of the added value of an ACPA multiplex assay in an early rheumatoid arthritis setting. <i>Arthritis Research and Therapy</i> , 2015 , 17, 276	5.7	16
127	Concordance between inflammation at physical examination and on MRI in patients with early arthritis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 506-12	2.4	41
126	Evaluating processes underlying the predictive value of baseline erosions for future radiological damage in early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 883-9	2.4	12
125	IL2RA is associated with persistence of rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2015 , 17, 244	5.7	14
124	Characterising arthralgia in the preclinical phase of rheumatoid arthritis using MRI. <i>Annals of the Rheumatic Diseases</i> , 2015 , 74, 1225-32	2.4	69
123	What is the evidence for the presence of a therapeutic window of opportunity in rheumatoid arthritis? A systematic literature review. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 861-70	2.4	182
122	Risk estimation in rheumatoid arthritis: from bench to bedside. <i>Nature Reviews Rheumatology</i> , 2014 , 10, 171-80	8.1	9
121	Low-avidity anticitrullinated protein antibodies (ACPA) are associated with a higher rate of joint destruction in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 270-6	2.4	33

120	Long-term outcome of rheumatoid arthritis defined according to the 2010-classification criteria. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 428-32	2.4	19
119	Smoking as a risk factor for the radiological severity of rheumatoid arthritis: a study on six cohorts. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1384-7	2.4	28
118	Aiming for a shorter rheumatoid arthritis MRI protocol: can contrast-enhanced MRI replace T2 for the detection of bone marrow oedema?. <i>European Radiology</i> , 2014 , 24, 2614-22	8	29
117	Macrophage migration inhibitory factor (MIF) -173 polymorphism is associated with clinical erythema nodosum in LfgrenB syndrome. <i>Cytokine</i> , 2014 , 69, 272-6	4	8
116	Is HLA-B27 increased in patients diagnosed with undifferentiated arthritis? Results from the Leiden early arthritis cohort. <i>Journal of Rheumatology</i> , 2014 , 41, 1948-51	4.1	1
115	Subclinical inflammation on MRI of hand and foot of anticitrullinated peptide antibody-negative arthralgia patients at risk for rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2014 , 16, R92	5.7	22
114	A genetic variant in osteoprotegerin is associated with progression of joint destruction in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2014 , 16, R108	5.7	16
113	Identification of a genetic variant for joint damage progression in autoantibody-positive rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 2038-46	2.4	34
112	A1.29 In rheumatoid arthritis, smoking is not primarily associated with anti-citrullinated protein antibodies, but with the presence of several autoantibodies. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, A12.1-A12	2.4	
111	Differential methylation within the major histocompatibility complex region in rheumatoid arthritis: a replication study. <i>Rheumatology</i> , 2014 , 53, 2317-8	3.9	9
110	Rheumatoid factor and anti-citrullinated protein antibody positivity, but not level, are associated with increased mortality in patients with rheumatoid arthritis: results from two large independent cohorts. <i>Arthritis Research and Therapy</i> , 2014 , 16, 483	5.7	41
109	A genetic variant in the region of MMP-9 is associated with serum levels and progression of joint damage in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1163-9	2.4	41
108	SPP1 rs9138 variant contributes to the severity of radiological damage in anti-citrullinated protein autoantibody-negative rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 1840-3	2.4	8
107	Are baseline high molecular weight adiponectin levels associated with radiographic progression in rheumatoid arthritis and osteoarthritis?. <i>Journal of Rheumatology</i> , 2014 , 41, 853-7	4.1	11
106	MRI-detected subclinical joint inflammation is associated with radiographic progression. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 2034-7	2.4	26
105	Does a genetic variant in FOXO3A predict a milder course of rheumatoid arthritis?. <i>Arthritis and Rheumatology</i> , 2014 , 66, 1678-81	9.5	5
104	Are rheumatoid arthritis patients discernible from other early arthritis patients using 1.5T extremity magnetic resonance imaging? a large cross-sectional study. <i>Journal of Rheumatology</i> , 2014 , 41, 1630-7	4.1	17
103	Is the modified disease activity score superior to the disease activity score in early arthritis and rheumatoid arthritis? Comment on the article by Baker et al. <i>Arthritis and Rheumatology</i> , 2014 , 66, 3244-5	8.5	3

102	Association of variants in IL2RA with progression of joint destruction in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2013 , 65, 1684-93		36
101	Anaemia to predict radiographic progression in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, e16	2.4	15
100	MRI of hand and foot joints of patients with anticitrullinated peptide antibody positive arthralgia without clinical arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 1540-4	2.4	46
99	ACPA fine-specificity profiles in early rheumatoid arthritis patients do not correlate with clinical features at baseline or with disease progression. <i>Arthritis Research and Therapy</i> , 2013 , 15, R140	5.7	44
98	Genetic studies on components of the Wnt signalling pathway and the severity of joint destruction in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 769-75	2.4	53
97	Evaluation of the contribution of cumulative levels of inflammation to the variance in joint destruction in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 307-8	2.4	8
96	Comparison of methodologies for analysing the progression of joint destruction in rheumatoid arthritis. <i>Scandinavian Journal of Rheumatology</i> , 2013 , 42, 182-9	1.9	27
95	A genetic variant in granzyme B is associated with progression of joint destruction in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2013 , 65, 582-9		15
94	Association of genetic variants in the IL4 and IL4R genes with the severity of joint damage in rheumatoid arthritis: a study in seven cohorts. <i>Arthritis and Rheumatism</i> , 2013 , 65, 3051-7		19
93	Patients with early arthritis consume less alcohol than controls, regardless of the type of arthritis. <i>Rheumatology</i> , 2013 , 52, 1701-7	3.9	12
92	An evaluation of molecular and clinical remission in rheumatoid arthritis by assessing radiographic progression. <i>Rheumatology</i> , 2013 , 52, 839-46	3.9	54
91	Allele-dose association of the C5orf30 rs26232 variant with joint damage in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2013 , 65, 2555-61		17
90	Evaluating joint destruction in rheumatoid arthritis: is it necessary to radiograph both hands and feet?. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 345-9	2.4	18
89	Role of NOD1 polymorphism in susceptibility and clinical progression of rheumatoid arthritis. <i>Rheumatology</i> , 2013 , 52, 806-14	3.9	12
88	How to define baseline erosiveness to predict radiologic progression in rheumatoid arthritis. <i>Journal of Rheumatology</i> , 2013 , 40, 1624-5	4.1	2
87	Serum pyridinoline levels and prediction of severity of joint destruction in rheumatoid arthritis. <i>Journal of Rheumatology</i> , 2013 , 40, 1303-6	4.1	14
86	Is early remission associated with improved survival or is arthritis persistency associated with increased mortality in early arthritis? Comparisons with the general population. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, e25	2.4	4
85	Risk of rheumatoid arthritis development in patients with unclassified arthritis according to the 2010 ACR/EULAR criteria for rheumatoid arthritis. <i>Rheumatology</i> , 2013 , 52, 1265-70	3.9	22

84	Defining erosive disease typical of RA in the light of the ACR/EULAR 2010 criteria for rheumatoid arthritis; results of the data driven phase. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 590-5	2.4	15
83	EULAR definition of erosive disease in light of the 2010 ACR/EULAR rheumatoid arthritis classification criteria. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 479-81	2.4	85
82	Improved early identification of arthritis: evaluating the efficacy of Early Arthritis Recognition Clinics. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 1295-301	2.4	26
81	Reasons for medical help-seeking behaviour of patients with recent-onset arthralgia. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 1302-7	2.4	16
80	Associations between APOE genotypes and disease susceptibility, joint damage and lipid levels in patients with rheumatoid arthritis. <i>PLoS ONE</i> , 2013 , 8, e60970	3.7	15
79	The additive value of magnetic resonance imaging for bone edema in predicting rheumatoid arthritis development in early undifferentiated arthritis: comment on the article by Duer-Jensen et al. <i>Arthritis and Rheumatism</i> , 2012 , 64, 321-2; author reply 322-3		3
78	Validation of a novel multibiomarker test to assess rheumatoid arthritis disease activity. <i>Arthritis Care and Research</i> , 2012 , 64, 1794-803	4.7	130
77	Sustained drug-free remission in rheumatoid arthritis after DAS-driven or non-DAS-driven therapy: a comparison of two cohort studies. <i>Rheumatology</i> , 2012 , 51, 1120-8	3.9	28
76	The 2010 ACR/EULAR criteria for rheumatoid arthritis: do they affect the classification or diagnosis of rheumatoid arthritis?. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1596-8	2.4	20
75	A two-stage mixed-effects model approach for gene-set analyses in candidate gene studies. <i>Statistics in Medicine</i> , 2012 , 31, 1190-202	2.3	8
74	Genetic predisposition of the severity of joint destruction in rheumatoid arthritis: a population-based study. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 707-9	2.4	57
73	Undifferentiated arthritis characteristics and outcomes when applying the 2010 and 1987 criteria for rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 238-41	2.4	37
72	Studying associations between variants in TRAF1-C5 and TNFAIP3-OLIG3 and the progression of joint destruction in rheumatoid arthritis in multiple cohorts. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1753-5	2.4	19
71	Diagnostic value of performing chest x-rays in patients presenting with recent-onset arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 627-8	2.4	1
70	No impact of serum uric acid on the outcome of recent-onset arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1424-5	2.4	3
69	Genetic variants in IL15 associate with progression of joint destruction in rheumatoid arthritis: a multicohort study. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1651-7	2.4	40
68	Loss of metacarpal bone density predicts RA development in recent-onset arthritis. <i>Rheumatology</i> , 2012 , 51, 1037-41	3.9	34
67	The ACPA recognition profile and subgrouping of ACPA-positive RA patients. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 268-74	2.4	54

66	EULAR recommendations for terminology and research in individuals at risk of rheumatoid arthritis: report from the Study Group for Risk Factors for Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 638-41	2.4	263
65	Performance of classification criteria for peripheral spondyloarthritis and psoriatic arthritis in the Leiden Early Arthritis cohort. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 1366-9	2.4	38
64	Does the season at symptom onset influence the long-term severity of radiographic joint destruction in rheumatoid arthritis?. <i>Annals of the Rheumatic Diseases</i> , 2012 , 71, 2055-6	2.4	7
63	Can anti-cyclic citrullinated peptide antibody-negative RA be subdivided into clinical subphenotypes?. <i>Arthritis Research and Therapy</i> , 2011 , 13, R180	5.7	16
62	Genetic variation in GREM1 is a risk factor for fibrosis in pulmonary sarcoidosis. <i>Tissue Antigens</i> , 2011 , 77, 112-7		25
61	Classification of rheumatoid arthritis: comparison of the 1987 American College of Rheumatology criteria and the 2010 American College of Rheumatology/European League Against Rheumatism criteria. <i>Arthritis and Rheumatism</i> , 2011 , 63, 37-42		168
60	Toward a data-driven evaluation of the 2010 American College of Rheumatology/European League Against Rheumatism criteria for rheumatoid arthritis: is it sensible to look at levels of rheumatoid factor?. <i>Arthritis and Rheumatism</i> , 2011 , 63, 1190-9		29
59	Identification of CXCL13 as a marker for rheumatoid arthritis outcome using an in silico model of the rheumatic joint. <i>Arthritis and Rheumatism</i> , 2011 , 63, 1265-73		41
58	Baseline serum adipokine levels predict radiographic progression in early rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2011 , 63, 2567-74		83
57	Predicting arthritis outcomes--what can be learned from the Leiden Early Arthritis Clinic?. <i>Rheumatology</i> , 2011 , 50, 93-100	3.9	207
56	The window of opportunity in ACPA-positive rheumatoid arthritis is not explained by ACPA characteristics. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 1697-8	2.4	6
55	Distinct ACPA fine-specificities, formed under the influence of HLA shared epitope alleles, have no effect on radiographic joint damage in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, A5-A5	2.4	
54	Autoantibodies recognizing carbamylated proteins are present in sera of patients with rheumatoid arthritis and predict joint damage. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 17372-7	11.5	385
53	Distinct ACPA fine specificities, formed under the influence of HLA shared epitope alleles, have no effect on radiographic joint damage in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2011 , 70, 1461-4	2.4	39
52	Genome-wide association study meta-analysis identifies seven new rheumatoid arthritis risk loci. <i>Nature Genetics</i> , 2010 , 42, 508-14	36.3	969
51	Genetic variation of the Fc gamma receptor 3B gene and association with rheumatoid arthritis. <i>PLoS ONE</i> , 2010 , 5, e13173	3.7	22
50	The PTPN22 susceptibility risk variant is not associated with the rate of joint destruction in anti-citrullinated protein antibody-positive rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1730-1	2.4	4
49	Repair of joint erosions in rheumatoid arthritis: prevalence and patient characteristics in a large inception cohort. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 727-9	2.4	30

48	How to avoid phenotypic misclassification in using joint destruction as an outcome measure for rheumatoid arthritis?. <i>Rheumatology</i> , 2010 , 49, 1429-35	3.9	12
47	The ACPA isotype profile reflects long-term radiographic progression in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1110-6	2.4	58
46	Genetic variants in the prediction of rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 1694-6	2.4	36
45	Improved treatment strategies reduce the increased mortality risk in early RA patients. <i>Rheumatology</i> , 2010 , 49, 2210-6	3.9	45
44	Diagnostic value of anti-MCV antibodies in differentiating early inflammatory arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 730-2	2.4	43
43	Association of the 6q23 region with the rate of joint destruction in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010 , 69, 567-70	2.4	26
42	TRAF1/C5 polymorphism is not associated with increased mortality in rheumatoid arthritis: two large longitudinal studies. <i>Arthritis Research and Therapy</i> , 2010 , 12, R38	5.7	16
41	Acute rheumatic fever and poststreptococcal reactive arthritis reconsidered. <i>Current Opinion in Rheumatology</i> , 2010 , 22, 437-42	5.3	28
40	Rheumatoid arthritis risk allele PTPRC is also associated with response to anti-tumor necrosis factor alpha therapy. <i>Arthritis and Rheumatism</i> , 2010 , 62, 1849-61		74
39	Long-term impact of delay in assessment of patients with early arthritis. <i>Arthritis and Rheumatism</i> , 2010 , 62, 3537-46		262
38	Quantitative heritability of anti-citrullinated protein antibody-positive and anti-citrullinated protein antibody-negative rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2009 , 60, 916-23		158
37	Prevalence of and predictive factors for sustained disease-modifying antirheumatic drug-free remission in rheumatoid arthritis: results from two large early arthritis cohorts. <i>Arthritis and Rheumatism</i> , 2009 , 60, 2262-71		154
36	Value of anti-modified citrullinated vimentin and third-generation anti-cyclic citrullinated peptide compared with second-generation anti-cyclic citrullinated peptide and rheumatoid factor in predicting disease outcome in undifferentiated arthritis and rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2009 , 60, 2232-41		114
35	Association of a single-nucleotide polymorphism in CD40 with the rate of joint destruction in rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2009 , 60, 2242-7		84
34	Variation in IL7R predisposes to sarcoid inflammation. <i>Genes and Immunity</i> , 2009 , 10, 647-53	4.4	32
33	Genetic variants at CD28, PRDM1 and CD2/CD58 are associated with rheumatoid arthritis risk. <i>Nature Genetics</i> , 2009 , 41, 1313-8	36.3	272
32	Contribution of Fcγ receptor IIIA gene 158V/F polymorphism and copy number variation to the risk of ACPA-positive rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2009 , 68, 1775-80	2.4	42
31	The prognostic value of baseline erosions in undifferentiated arthritis. <i>Arthritis Research and Therapy</i> , 2009 , 11, R155	5.7	33

30	Common variants at CD40 and other loci confer risk of rheumatoid arthritis. <i>Nature Genetics</i> , 2008 , 40, 1216-23	36.3	416
29	Advances in the genetics of rheumatoid arthritis point to subclassification into distinct disease subsets. <i>Arthritis Research and Therapy</i> , 2008 , 10, 205	5.7	101
28	A high body mass index has a protective effect on the amount of joint destruction in small joints in early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2008 , 67, 769-74	2.4	132
27	Pretreatment serum levels of anti-cyclic citrullinated peptide antibodies are associated with the response to methotrexate in recent-onset arthritis. <i>Annals of the Rheumatic Diseases</i> , 2008 , 67, 1194-5	2.4	27
26	Biomarkers in the prediction of radiographic progression in early rheumatoid arthritis: comment on the article by Young-Min et al. <i>Arthritis and Rheumatism</i> , 2008 , 58, 1558-9; author reply 1559		2
25	The inflammatory disease-associated variants in IL12B and IL23R are not associated with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2008 , 58, 1877-81		38
24	Validation of a prediction rule for disease outcome in patients with recent-onset undifferentiated arthritis: moving toward individualized treatment decision-making. <i>Arthritis and Rheumatism</i> , 2008 , 58, 2241-7		121
23	A large-scale rheumatoid arthritis genetic study identifies association at chromosome 9q33.2. <i>PLoS Genetics</i> , 2008 , 4, e1000107	6	63
22	The HLA-DRB1 shared epitope alleles differ in the interaction with smoking and predisposition to antibodies to cyclic citrullinated peptide. <i>Arthritis and Rheumatism</i> , 2007 , 56, 425-32		112
21	A prediction rule for disease outcome in patients with recent-onset undifferentiated arthritis: how to guide individual treatment decisions. <i>Arthritis and Rheumatism</i> , 2007 , 56, 433-40		267
20	Association of tumor necrosis factor alpha polymorphism and radiographic progression in rheumatoid arthritis: comment on the article by Khanna et al. <i>Arthritis and Rheumatism</i> , 2007 , 56, 1032-3; author reply 1033-4		
19	Emerging patterns of risk factor make-up enable subclassification of rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2007 , 56, 1728-35		41
18	A candidate gene approach identifies the TRAF1/C5 region as a risk factor for rheumatoid arthritis. <i>PLoS Medicine</i> , 2007 , 4, e278	11.6	192
17	Arthritis of the large joints - in particular, the knee - at first presentation is predictive for a high level of radiological destruction of the small joints in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2007 , 66, 646-50	2.4	32
16	Protective effect of noninherited maternal HLA-DR antigens on rheumatoid arthritis development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 19966-70	11.5	52
15	Gene-gene and gene-environment interactions involving HLA-DRB1, PTPN22, and smoking in two subsets of rheumatoid arthritis. <i>American Journal of Human Genetics</i> , 2007 , 80, 867-75	11	317
14	The HLA-DRB1 shared epitope alleles are primarily a risk factor for anti-cyclic citrullinated peptide antibodies and are not an independent risk factor for development of rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2006 , 54, 1117-21		253
13	Variation in radiologic joint destruction in rheumatoid arthritis differs between monozygotic and dizygotic twins and pairs of unrelated patients. <i>Arthritis and Rheumatism</i> , 2006 , 54, 2028-30		28

12	Smoking is a risk factor for anti-CCP antibodies only in rheumatoid arthritis patients who carry HLA-DRB1 shared epitope alleles. <i>Annals of the Rheumatic Diseases</i> , 2006 , 65, 366-71	2.4	216
11	Aspects of early arthritis. Definition of disease states in early arthritis: remission versus minimal disease activity. <i>Arthritis Research and Therapy</i> , 2006 , 8, 216	5.7	27
10	Antibodies to citrullinated proteins and differences in clinical progression of rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2005 , 7, R949-58	5.7	325
9	Understanding the genetic contribution to rheumatoid arthritis. <i>Current Opinion in Rheumatology</i> , 2005 , 17, 299-304	5.3	75
8	Invasiveness of fibroblast-like synoviocytes is an individual patient characteristic associated with the rate of joint destruction in patients with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2005 , 52, 1999-2002		110
7	An independent role of protective HLA class II alleles in rheumatoid arthritis severity and susceptibility. <i>Arthritis and Rheumatism</i> , 2005 , 52, 2637-44		77
6	Association of the PTPN22 C1858T single-nucleotide polymorphism with rheumatoid arthritis phenotypes in an inception cohort. <i>Arthritis and Rheumatism</i> , 2005 , 52, 2948-50		68
5	Association of HLA-DR3 with anti-cyclic citrullinated peptide antibody-negative rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2005 , 52, 3058-62		140
4	Refining the complex rheumatoid arthritis phenotype based on specificity of the HLA-DRB1 shared epitope for antibodies to citrullinated proteins. <i>Arthritis and Rheumatism</i> , 2005 , 52, 3433-8		427
3	Hypernatremia from a hunger strike as a cause of osmotic myelinolysis. <i>Neurology</i> , 2005 , 64, 574-5	6.5	22
2	Immunoglobulin A multiple myeloma presenting with Henoch-Schönlein purpura associated with reduced sialylation of IgA1. <i>British Journal of Haematology</i> , 2003 , 122, 915-7	4.5	30
1	No influence of melatonin on cerebral blood flow in humans. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2003 , 88, 5989-94	5.6	43