

Paul Russell

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1712946/paul-russell-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

94
papers

10,773
citations

53
h-index

103
g-index

107
ext. papers

11,589
ext. citations

12.8
avg, IF

6.08
L-index

#	Paper	IF	Citations
94	EXO5-DNA structure and BLM interactions direct DNA resection critical for ATR-dependent replication restart. <i>Molecular Cell</i> , 2021 , 81, 2989-3006.e9	17.6	8
93	Molecular basis of chromatin remodeling by Rhp26, a yeast CSB ortholog. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 6120-6129	11.5	7
92	Brc1 Promotes the Focal Accumulation and SUMO Ligase Activity of Smc5-Smc6 during Replication Stress. <i>Molecular and Cellular Biology</i> , 2019 , 39,	4.8	12
91	RNase H eliminates R-loops that disrupt DNA replication but is nonessential for efficient DSB repair. <i>EMBO Reports</i> , 2018 , 19,	6.5	39
90	Mre11 complex links sister chromatids to promote repair of a collapsed replication fork. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018 , 115, 8793-8798	11.5	16
89	Mre11-Rad50-dependent activity of ATM/Tel1 at DNA breaks and telomeres in the absence of Nbs1. <i>Molecular Biology of the Cell</i> , 2018 , 29, 1389-1399	3.5	13
88	Tdp1 processes chromate-induced single-strand DNA breaks that collapse replication forks. <i>PLoS Genetics</i> , 2018 , 14, e1007595	6	2
87	Lingering single-strand breaks trigger Rad51-independent homology-directed repair of collapsed replication forks in the polynucleotide kinase/phosphatase mutant of fission yeast. <i>PLoS Genetics</i> , 2017 , 13, e1007013	6	2
86	Multi-BRCT Domain Protein Brc1 Links Rhp18/Rad18 and H2A To Maintain Genome Stability during S Phase. <i>Molecular and Cellular Biology</i> , 2017 , 37,	4.8	2
85	Ctp1-dependent clipping and resection of DNA double-strand breaks by Mre11 endonuclease complex are not genetically separable. <i>Nucleic Acids Research</i> , 2016 , 44, 8241-9	20.1	11
84	Growth and the Environment of <i>Schizosaccharomyces pombe</i> . <i>Cold Spring Harbor Protocols</i> , 2016 , 2016, pdb.top079764	1.2	76
83	Global Fitness Profiling Identifies Arsenic and Cadmium Tolerance Mechanisms in Fission Yeast. <i>G3: Genes, Genomes, Genetics</i> , 2016 , 6, 3317-3333	3.2	15
82	Genetic Interaction Landscape Reveals Critical Requirements for <i>Schizosaccharomyces pombe</i> Brc1 in DNA Damage Response Mutants. <i>G3: Genes, Genomes, Genetics</i> , 2015 , 5, 953-62	3.2	7
81	Ku stabilizes replication forks in the absence of Brc1. <i>PLoS ONE</i> , 2015 , 10, e0126598	3.7	7
80	Critical Function of H2A in S-Phase. <i>PLoS Genetics</i> , 2015 , 11, e1005517	6	10
79	Regulation of the Rhp26/ERCC6/CSB chromatin remodeler by a novel conserved leucine latch motif. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 18566-71	11.5	21
78	Tolerance of deregulated G1/S transcription depends on critical G1/S regulon genes to prevent catastrophic genome instability. <i>Cell Reports</i> , 2014 , 9, 2279-89	10.6	11

77	Mdb1, a fission yeast homolog of human MDC1, modulates DNA damage response and mitotic spindle function. <i>PLoS ONE</i> , 2014 , 9, e97028	3.7	14
76	Regulation of Mus81-Eme1 Holliday junction resolvase in response to DNA damage. <i>Nature Structural and Molecular Biology</i> , 2013 , 20, 598-603	17.6	47
75	Brc1 links replication stress response and centromere function. <i>Cell Cycle</i> , 2013 , 12, 1665-71	4.7	8
74	H2A-binding protein Brc1 affects centromere function in fission yeast. <i>Molecular and Cellular Biology</i> , 2013 , 33, 1410-6	4.8	16
73	Replication fork collapse and genome instability in a deoxycytidylate deaminase mutant. <i>Molecular and Cellular Biology</i> , 2012 , 32, 4445-54	4.8	33
72	Phosphorylation-dependent interactions between Crb2 and Chk1 are essential for DNA damage checkpoint. <i>PLoS Genetics</i> , 2012 , 8, e1002817	6	11
71	Cadmium-induced proteome remodeling regulated by Spc1/Sty1 and Zip1 in fission yeast. <i>Toxicological Sciences</i> , 2012 , 129, 200-12	4.4	13
70	Mre11 ATLD17/18 mutation retains Tel1/ATM activity but blocks DNA double-strand break repair. <i>Nucleic Acids Research</i> , 2012 , 40, 11435-49	20.1	17
69	ABC ATPase signature helices in Rad50 link nucleotide state to Mre11 interface for DNA repair. <i>Nature Structural and Molecular Biology</i> , 2011 , 18, 423-31	17.6	129
68	Regulatory networks integrating cell cycle control with DNA damage checkpoints and double-strand break repair. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2011 , 366, 3562-71	5.8	95
67	Mre11 nuclease activity and Ctp1 regulate Chk1 activation by Rad3ATR and Tel1ATM checkpoint kinases at double-strand breaks. <i>Molecular and Cellular Biology</i> , 2011 , 31, 573-83	4.8	33
66	Release of Ku and MRN from DNA ends by Mre11 nuclease activity and Ctp1 is required for homologous recombination repair of double-strand breaks. <i>PLoS Genetics</i> , 2011 , 7, e1002271	6	168
65	gammaH2A binds Brc1 to maintain genome integrity during S-phase. <i>EMBO Journal</i> , 2010 , 29, 1136-48	13	67
64	Critical functions of Rpa3/Ssb3 in S-phase DNA damage responses in fission yeast. <i>PLoS Genetics</i> , 2010 , 6, e1001138	6	15
63	Tonoplast-localized Abc2 transporter mediates phytochelatin accumulation in vacuoles and confers cadmium tolerance. <i>Journal of Biological Chemistry</i> , 2010 , 285, 40416-26	5.4	74
62	BRCT domain interactions with phospho-histone H2A target Crb2 to chromatin at double-strand breaks and maintain the DNA damage checkpoint. <i>Molecular and Cellular Biology</i> , 2010 , 30, 4732-43	4.8	22
61	Rad3 decorates critical chromosomal domains with gammaH2A to protect genome integrity during S-Phase in fission yeast. <i>PLoS Genetics</i> , 2010 , 6, e1001032	6	58
60	Phosphorylation-regulated binding of Ctp1 to Nbs1 is critical for repair of DNA double-strand breaks. <i>Cell Cycle</i> , 2010 , 9, 1516-22	4.7	24

59	Mms1-Mms22 complex protects genome integrity in <i>Schizosaccharomyces pombe</i> . <i>DNA Repair</i> , 2009 , 8, 1390-9	4.3	9
58	Nbs1 flexibly tethers Ctp1 and Mre11-Rad50 to coordinate DNA double-strand break processing and repair. <i>Cell</i> , 2009 , 139, 87-99	56.2	242
57	Fission yeast Scm3 mediates stable assembly of Cnp1/CENP-A into centromeric chromatin. <i>Molecular Cell</i> , 2009 , 33, 287-98	17.6	151
56	Assays used to study the DNA replication checkpoint in fission yeast. <i>Methods in Molecular Biology</i> , 2009 , 521, 493-507	1.4	17
55	Mus81 is essential for sister chromatid recombination at broken replication forks. <i>EMBO Journal</i> , 2008 , 27, 1378-87	13	90
54	Mre11 dimers coordinate DNA end bridging and nuclease processing in double-strand-break repair. <i>Cell</i> , 2008 , 135, 97-109	56.2	358
53	A genome-wide screen of genes involved in cadmium tolerance in <i>Schizosaccharomyces pombe</i> . <i>Toxicological Sciences</i> , 2008 , 106, 124-39	4.4	46
52	Xlf1 is required for DNA repair by nonhomologous end joining in <i>Schizosaccharomyces pombe</i> . <i>Genetics</i> , 2007 , 175, 963-7	4	14
51	Mms22 preserves genomic integrity during DNA replication in <i>Schizosaccharomyces pombe</i> . <i>Genetics</i> , 2007 , 177, 47-61	4	19
50	Ctp1 is a cell-cycle-regulated protein that functions with Mre11 complex to control double-strand break repair by homologous recombination. <i>Molecular Cell</i> , 2007 , 28, 134-46	17.6	262
49	Rad22Rad52-dependent repair of ribosomal DNA repeats cleaved by Slx1-Slx4 endonuclease. <i>Molecular Biology of the Cell</i> , 2006 , 17, 2081-90	3.5	33
48	Cip1 and Cip2 are novel RNA-recognition-motif proteins that counteract Csx1 function during oxidative stress. <i>Molecular Biology of the Cell</i> , 2006 , 17, 1176-83	3.5	19
47	Histone modification-dependent and -independent pathways for recruitment of checkpoint protein Crb2 to double-strand breaks. <i>Genes and Development</i> , 2006 , 20, 1583-96	12.6	113
46	Constraining G1-specific transcription to late G1 phase: the MBF-associated corepressor Nrm1 acts via negative feedback. <i>Molecular Cell</i> , 2006 , 23, 483-96	17.6	106
45	Sws1 is a conserved regulator of homologous recombination in eukaryotic cells. <i>EMBO Journal</i> , 2006 , 25, 2564-74	13	88
44	Cooperative control of Crb2 by ATM family and Cdc2 kinases is essential for the DNA damage checkpoint in fission yeast. <i>Molecular and Cellular Biology</i> , 2005 , 25, 10721-30	4.8	22
43	ATM activation and its recruitment to damaged DNA require binding to the C terminus of Nbs1. <i>Molecular and Cellular Biology</i> , 2005 , 25, 5363-79	4.8	336
42	Distinct signaling pathways respond to arsenite and reactive oxygen species in <i>Schizosaccharomyces pombe</i> . <i>Eukaryotic Cell</i> , 2005 , 4, 1396-402		42

41	Swi1 and Swi3 are components of a replication fork protection complex in fission yeast. <i>Molecular and Cellular Biology</i> , 2004 , 24, 8342-55	4.8	171
40	Histone H2A phosphorylation controls Crb2 recruitment at DNA breaks, maintains checkpoint arrest, and influences DNA repair in fission yeast. <i>Molecular and Cellular Biology</i> , 2004 , 24, 6215-30	4.8	172
39	Slx1-Slx4 are subunits of a structure-specific endonuclease that maintains ribosomal DNA in fission yeast. <i>Molecular Biology of the Cell</i> , 2004 , 15, 71-80	3.5	99
38	RNA-binding protein Csx1 mediates global control of gene expression in response to oxidative stress. <i>EMBO Journal</i> , 2003 , 22, 6256-66	13	60
37	The endogenous Mus81-Eme1 complex resolves Holliday junctions by a nick and counternick mechanism. <i>Molecular Cell</i> , 2003 , 12, 747-59	17.6	149
36	Retention but not recruitment of Crb2 at double-strand breaks requires Rad1 and Rad3 complexes. <i>Molecular and Cellular Biology</i> , 2003 , 23, 6150-8	4.8	82
35	The fission yeast Rad32 (Mre11)-Rad50-Nbs1 complex is required for the S-phase DNA damage checkpoint. <i>Molecular and Cellular Biology</i> , 2003 , 23, 6564-73	4.8	64
34	Swi1 prevents replication fork collapse and controls checkpoint kinase Cds1. <i>Molecular and Cellular Biology</i> , 2003 , 23, 7861-74	4.8	146
33	Replication checkpoint protein Mrc1 is regulated by Rad3 and Tel1 in fission yeast. <i>Molecular and Cellular Biology</i> , 2003 , 23, 8395-403	4.8	48
32	Fission yeast Mus81-Eme1 Holliday junction resolvase is required for meiotic crossing over but not for gene conversion. <i>Genetics</i> , 2003 , 165, 2289-93	4	96
31	CDK phosphorylation of Drc1 regulates DNA replication in fission yeast. <i>Current Biology</i> , 2002 , 12, 599-605	6.5	57
30	Telomere binding of checkpoint sensor and DNA repair proteins contributes to maintenance of functional fission yeast telomeres. <i>Genetics</i> , 2002 , 161, 1437-52	4	99
29	Mrc1 channels the DNA replication arrest signal to checkpoint kinase Cds1. <i>Nature Cell Biology</i> , 2001 , 3, 966-72	23.4	200
28	DNA replication checkpoint. <i>Current Biology</i> , 2001 , 11, R953-6	6.3	102
27	Threonine-11, phosphorylated by Rad3 and atm in vitro, is required for activation of fission yeast checkpoint kinase Cds1. <i>Molecular and Cellular Biology</i> , 2001 , 21, 3398-404	4.8	47
26	Roles of the mitotic inhibitors Wee1 and Mik1 in the G(2) DNA damage and replication checkpoints. <i>Molecular and Cellular Biology</i> , 2001 , 21, 1499-508	4.8	66
25	Mus81-Eme1 are essential components of a Holliday junction resolvase. <i>Cell</i> , 2001 , 107, 537-48	56.2	450
24	Checkpoints: it takes more than time to heal some wounds. <i>Current Biology</i> , 2000 , 10, R908-11	6.3	54

23	Regulation of mitotic inhibitor Mik1 helps to enforce the DNA damage checkpoint. <i>Molecular Biology of the Cell</i> , 2000 , 11, 1-11	3.5	66
22	Mechanism of caffeine-induced checkpoint override in fission yeast. <i>Molecular and Cellular Biology</i> , 2000 , 20, 4288-94	4.8	64
21	Damage tolerance protein Mus81 associates with the FHA1 domain of checkpoint kinase Cds1. <i>Molecular and Cellular Biology</i> , 2000 , 20, 8758-66	4.8	249
20	Slm9, a novel nuclear protein involved in mitotic control in fission yeast. <i>Genetics</i> , 2000 , 155, 623-31	4	12
19	Cdc25 inhibited in vivo and in vitro by checkpoint kinases Cds1 and Chk1. <i>Molecular Biology of the Cell</i> , 1999 , 10, 833-45	3.5	191
18	Nuclear localization of Cdc25 is regulated by DNA damage and a 14-3-3 protein. <i>Nature</i> , 1999 , 397, 172-5	5.4	517
17	Basis for the checkpoint signal specificity that regulates Chk1 and Cds1 protein kinases. <i>Molecular and Cellular Biology</i> , 1999 , 19, 4262-9	4.8	100
16	Checkpoints on the road to mitosis. <i>Trends in Biochemical Sciences</i> , 1998 , 23, 399-402	10.3	113
15	Mitotic DNA damage and replication checkpoints in yeast. <i>Current Opinion in Cell Biology</i> , 1998 , 10, 749-58	5.8	145
14	Replication checkpoint enforced by kinases Cds1 and Chk1. <i>Science</i> , 1998 , 280, 909-12	33.3	293
13	Heat stress activates fission yeast Spc1/Sty1 MAPK by a MEKK-independent mechanism. <i>Molecular Biology of the Cell</i> , 1998 , 9, 1339-49	3.5	100
12	Negative regulation of Cdc18 DNA replication protein by Cdc2. <i>Molecular Biology of the Cell</i> , 1998 , 9, 63-73	3.5	80
11	Phosphorylation and association with the transcription factor Atf1 regulate localization of Spc1/Sty1 stress-activated kinase in fission yeast. <i>Genes and Development</i> , 1998 , 12, 1464-73	12.6	130
10	Tyrosine phosphorylation of cdc2 is required for the replication checkpoint in <i>Schizosaccharomyces pombe</i> . <i>Molecular and Cellular Biology</i> , 1998 , 18, 3782-7	4.8	99
9	The <i>Schizosaccharomyces pombe</i> S-phase checkpoint differentiates between different types of DNA damage. <i>Genetics</i> , 1998 , 149, 1729-37	4	55
8	Cdc25 mitotic inducer targeted by chk1 DNA damage checkpoint kinase. <i>Science</i> , 1997 , 277, 1495-7	33.3	476
7	Interaction of Cdc2 and Cdc18 with a fission yeast ORC2-like protein. <i>Nature</i> , 1996 , 379, 360-3	50.4	130
6	Cell-cycle control linked to extracellular environment by MAP kinase pathway in fission yeast. <i>Nature</i> , 1995 , 378, 739-43	50.4	417

- 5 Nim1 kinase promotes mitosis by inactivating Wee1 tyrosine kinase. *Nature*, **1993**, 363, 738-41 50.4 144
- 4 Activation of MPF in fission yeast. *Novartis Foundation Symposium*, **1992**, 170, 50-8; discussion 58-71
- 3 Fission yeast p107wee1 mitotic inhibitor is a tyrosine/serine kinase. *Nature*, **1991**, 349, 808-11 50.4 338
- 2 Negative regulation of mitosis by wee1+, a gene encoding a protein kinase homolog. *Cell*, **1987**, 49, 559-572 56.2 916
- 1 cdc25+ functions as an inducer in the mitotic control of fission yeast. *Cell*, **1986**, 45, 145-53 56.2 896