

Henriette Acosta

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1715382/publications.pdf>

Version: 2024-02-01

16
papers

319
citations

933447

10
h-index

996975

15
g-index

16
all docs

16
docs citations

16
times ranked

549
citing authors

#	ARTICLE	IF	CITATIONS
1	Oxytocin receptor polymorphism and childhood social experiences shape adult personality, brain structure and neural correlates of mentalizing. <i>NeuroImage</i> , 2016, 134, 671-684.	4.2	58
2	Maternal Pregnancy-Related Anxiety Is Associated With Sexually Dimorphic Alterations in Amygdala Volume in 4-Year-Old Children. <i>Frontiers in Behavioral Neuroscience</i> , 2019, 13, 175.	2.0	46
3	Adult attachment style modulates neural responses in a mentalizing task. <i>Neuroscience</i> , 2015, 303, 462-473.	2.3	38
4	Intergroup empathy: Enhanced neural resonance for ingroup facial emotion in a shared neural production-perception network. <i>NeuroImage</i> , 2019, 194, 182-190.	4.2	26
5	Newborn amygdalar volumes are associated with maternal prenatal psychological distress in a sex-dependent way. <i>NeuroImage: Clinical</i> , 2020, 28, 102380.	2.7	25
6	Partial Support for an Interaction Between a Polygenic Risk Score for Major Depressive Disorder and Prenatal Maternal Depressive Symptoms on Infant Right Amygdalar Volumes. <i>Cerebral Cortex</i> , 2020, 30, 6121-6134.	2.9	21
7	Emotional Availability Modulates Electrophysiological Correlates of Executive Functions in Preschool Children. <i>Frontiers in Human Neuroscience</i> , 2016, 10, 299.	2.0	15
8	Emotion specific neural activation for the production and perception of facial expressions. <i>Cortex</i> , 2020, 127, 17-28.	2.4	15
9	Outgroup emotion processing in the vACC is modulated by childhood trauma and CACNA1C risk variant. <i>Social Cognitive and Affective Neuroscience</i> , 2018, 13, 341-348.	3.0	13
10	A voxel-based morphometry study on adult attachment style and affective loss. <i>Neuroscience</i> , 2018, 392, 219-229.	2.3	12
11	Schizotypy and mentalizing: An fMRI study. <i>Neuropsychologia</i> , 2019, 124, 299-310.	1.6	11
12	Prenatal maternal depressive symptoms are associated with smaller amygdalar volumes of four-year-old children. <i>Psychiatry Research - Neuroimaging</i> , 2020, 304, 111153.	1.8	11
13	Sex-specific association between infant caudate volumes and a polygenic risk score for major depressive disorder. <i>Journal of Neuroscience Research</i> , 2020, 98, 2529-2540.	2.9	10
14	A variation in the infant oxytocin receptor gene modulates infant hippocampal volumes in association with sex and prenatal maternal anxiety. <i>Psychiatry Research - Neuroimaging</i> , 2021, 307, 111207.	1.8	6
15	Larger bilateral amygdalar volumes are associated with affective loss experiences. <i>Journal of Neuroscience Research</i> , 2021, 99, 1763-1779.	2.9	6
16	Allometry in the corpus callosum in neonates: Sexual dimorphism. <i>Human Brain Mapping</i> , 0, , .	3.6	6