

# Miroslav ZoriÄ

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1721216/publications.pdf>

Version: 2024-02-01

21  
papers

98  
citations

1478505

6  
h-index

1474206

9  
g-index

21  
all docs

21  
docs citations

21  
times ranked

126  
citing authors

#	ARTICLE	IF	CITATIONS
1	Qualitative traits in sunflower breeding: UGAâ€šAM1 phenotyping case study. <i>Crop Science</i> , 2020, 60, 303-319.	1.8	6
2	Origin and diversity study of local common bean ( <i>Phaseolus vulgaris</i> L.) germplasm from Serbia: phaseolin and phenotyping approach. <i>Genetic Resources and Crop Evolution</i> , 2020, 67, 2195-2212.	1.6	3
3	Effect of harvest maturity stage and seeding rate on alfalfa yield and quality. <i>Ratarstvo I Povrtarstvo</i> , 2020, 57, 35-42.	0.5	1
4	Effect of organic and conventional farming system and sowing date on yield, seed oil and protein content in rapeseed cultivars. <i>Ratarstvo I Povrtarstvo</i> , 2020, 57, 55-60.	0.5	0
5	Dissection of Year Related Climatic Variables and Their Effect on Winter Rapeseed ( <i>Brassica Napus</i> L.) Development and Yield. <i>Agronomy</i> , 2019, 9, 517.	3.0	15
6	Variability of red clover genotypes on the basis of morphological markers. <i>Genetika</i> , 2018, 50, 895-906.	0.4	4
7	Phenotypic variability of spike characteristics and genetic structure of population in the collection of spring barley. <i>Selekcija I Semearstvo</i> , 2018, 24, 49-58.	0.4	1
8	Biplot analysis of seed priming effects on maize seedling growth traits. <i>Ratarstvo I Povrtarstvo</i> , 2018, 55, 111-117.	0.5	1
9	Classification of confectionary sunflower genotypes based on morphological characters. <i>Journal of Agricultural Science</i> , 2017, 155, 1594-1609.	1.3	13
10	Direct and indirect effects of morphophysiological traits on kernel protein content of confectionary sunflower. <i>Genetika</i> , 2017, 49, 1015-1021.	0.4	0
11	Yields of ns maize hybrids from different breeding periods. <i>Selekcija I Semearstvo</i> , 2015, 21, 93-102.	0.4	1
12	Enetic diversity in red clover ( <i>Trifolium pratense</i> L.) using SSR markers. <i>Genetika</i> , 2014, 46, 949-961.	0.4	9
13	Viability of oilseed rape ( <i>Brassica napus</i> L.) seeds under salt stress. <i>Genetika</i> , 2014, 46, 137-148.	0.4	4
14	Molecular characterization of barley ( <i>Hordeum vulgare</i> L.) accessions of the Serbian GeneBank by SSR fingerprinting. <i>Genetika</i> , 2013, 45, 167-180.	0.4	5
15	Evaluation of seed and oil yield stability in NS rapeseed cultivars ( <i>Brassica napus</i> L.). <i>Ratarstvo I Povrtarstvo</i> , 2011, 48, 67-76.	0.5	5
16	Effect of nitrogen rate on grain yield of bread wheat genotypes. <i>Genetika</i> , 2010, 42, 279-286.	0.4	2
17	Analysis of grain size in bean ( <i>Phaseolus vulgaris</i> L.) by linear and bilinear models. <i>Genetika</i> , 2010, 42, 535-544.	0.4	1
18	Variability of leaf Cadmium content in tetraploid and hexaploid wheat. <i>Genetika</i> , 2009, 41, 1-10.	0.4	8

#	ARTICLE	IF	CITATIONS
19	Tissue culture and agronomic traits relationship in wheat. <i>Plant Cell, Tissue and Organ Culture</i> , 2008, 95, 107-114.	2.3	13
20	Interpretation of interactions in sunflower agronomic trials using multiplicative models and climatic information. <i>Helia</i> , 2008, 31, 51-64.	0.4	2
21	Assessing wheat performance using environmental information. <i>Genetika</i> , 2007, 39, 413-425.	0.4	4