Antonio Sanchez-Amat

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1723935/publications.pdf

Version: 2024-02-01

55 papers 2,058 citations

218677 26 h-index 243625 44 g-index

59 all docs 59 docs citations

59 times ranked 1975 citing authors

#	Article	IF	CITATIONS
1	Direct CRISPR spacer acquisition from RNA by a natural reverse transcriptase–Cas1 fusion protein. Science, 2016, 351, aad4234.	12.6	170
2	Hydrogen Peroxide Linked to Lysine Oxidase Activity Facilitates Biofilm Differentiation and Dispersal in Several Gram-Negative Bacteria. Journal of Bacteriology, 2008, 190, 5493-5501.	2.2	119
3	A Pluripotent Polyphenol Oxidase from the Melanogenic MarineAlteromonas spShares Catalytic Capabilities of Tyrosinases and Laccases. Biochemical and Biophysical Research Communications, 1997, 240, 787-792.	2.1	97
4	Dimethoxyphenol oxidase activity of different microbial blue multicopper proteins. FEMS Microbiology Letters, 2001, 204, 175-181.	1.8	95
5	Molecular cloning and functional characterization of a unique multipotent polyphenol oxidase from Marinomonas mediterranea. BBA - Proteins and Proteomics, 2001, 1547, 104-116.	2.1	95
6	A tyrosinase with an abnormally high tyrosine hydroxylase/dopa oxidase ratio. FEBS Journal, 2006, 273, 257-270.	4.7	85
7	Type III CRISPR-Cas systems can provide redundancy to counteract viral escape from type I systems. ELife, 2017, 6, .	6.0	81
8	Taxonomic study of Marinomonas strains isolated from the seagrass Posidonia oceanica, with descriptions of Marinomonas balearica sp. nov. and Marinomonas pollencensis sp. nov International Journal of Systematic and Evolutionary Microbiology, 2010, 60, 93-98.	1.7	68
9	Alkali and halide-resistant catalysis by the multipotent oxidase from Marinomonas mediterranea. Journal of Biotechnology, 2005, 117, 73-82.	3 . 8	63
10	Note: Studies on the phylogenetic relationships of melanogenic marine bacteria: Proposal of Marinomonas mediterranea sp. nov International Journal of Systematic and Evolutionary Microbiology, 1999, 49, 1241-1246.	1.7	62
11	The Antimicrobial Activity of Marinocine, Synthesized by Marinomonas mediterranea, Is Due to Hydrogen Peroxide Generated by Its Lysine Oxidase Activity. Journal of Bacteriology, 2006, 188, 2493-2501.	2.2	60
12	A novel type of lysine oxidase: l-lysine-Îμ-oxidase. Biochimica Et Biophysica Acta - Proteins and Proteomics, 2006, 1764, 1577-1585.	2.3	55
13	Comparison of different microbial laccases as tools for industrial uses. New Biotechnology, 2016, 33, 387-398.	4.4	55
14	Marinomonas mediterranea MMB-1 Transposon Mutagenesis: Isolation of a Multipotent Polyphenol Oxidase Mutant. Journal of Bacteriology, 2000, 182, 3754-3760.	2.2	53
15	Purification and partial characterization of marinocine, a new broad-spectrum antibacterial protein produced by Marinomonas mediterranea. Biochimica Et Biophysica Acta - General Subjects, 2005, 1721, 193-203.	2.4	51
16	Characterization of the Melanogenic System in Vibrio cholerae, ATCC 14035. Pigment Cell & Melanoma Research, 1995, 8, 147-152.	3 . 6	49
17	Polyphenol Oxidase Activity Expression in Ralstonia solanacearum. Applied and Environmental Microbiology, 2005, 71, 6808-6815.	3.1	47
18	Identification of an operon involved in tyrosinase activity and melanin synthesis in Marinomonas mediterranea. Gene, 2004, 342, 179-187.	2.2	46

#	Article	IF	CITATIONS
19	Cloning and Molecular Characterization of a SDSâ€Activated Tyrosinase from <i>Marinomonas mediterranea</i> . Pigment Cell & Melanoma Research, 2002, 15, 104-111.	3.6	45
20	The macromolecule with antimicrobial activity synthesized by Pseudoalteromonas luteoviolacea strains is an l-amino acid oxidase. Applied Microbiology and Biotechnology, 2008, 79, 925-930.	3.6	45
21	Marinomonas alcarazii sp. nov., M. rhizomae sp. nov., M. foliarum sp. nov., M. posidonica sp. nov. and M. aquiplantarum sp. nov., isolated from the microbiota of the seagrass Posidonia oceanica. International Journal of Systematic and Evolutionary Microbiology, 2011, 61, 2191-2196.	1.7	43
22	Location and Catalytic Characteristics of a Multipotent Bacterial Polyphenol Oxidase. Pigment Cell & Melanoma Research, 1999, 12, 331-339.	3.6	35
23	Involvement of a novel copper chaperone in tyrosinase activity and melanin synthesis in Marinomonas mediterranea. Microbiology (United Kingdom), 2007, 153, 2241-2249.	1.8	35
24	Effects of epibiotic bacteria on leaf growth and epiphytes of the seagrass Posidonia oceanica. Marine Ecology - Progress Series, 2012, 456, 21-27.	1.9	29
25	Distribution in microbial genomes of genes similar to lodA and goxA which encode a novel family of quinoproteins with amino acid oxidase activity. BMC Genomics, 2015, 16, 231.	2.8	29
26	The protein encoded by the Shewanella colwellianamelA gene is a p-hydroxyphenylpyruvate dioxygenase. FEMS Microbiology Letters, 1994, 124, 179-184.	1.8	28
27	Comparative tyrosine degradation in Vibrio cholerae strains. The strain ATCC 14035 as a prokaryotic melanogenic model of homogentisate-releasing cell. Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology, 1998, 119, 557-562.	1.6	27
28	Finding New Enzymes from Bacterial Physiology: A Successful Approach Illustrated by the Detection of Novel Oxidases in Marinomonas mediterranea. Marine Drugs, 2010, 8, 519-541.	4.6	27
29	Both genes in the <i>Marinomonas mediterranea lodAB</i> operon are required for the expression of the antimicrobial protein lysine oxidase. Molecular Microbiology, 2010, 75, 462-473.	2.5	26
30	Identification in <i><scp>M</scp>arinomonas mediterranea</i> of a novel quinoprotein with glycine oxidase activity. MicrobiologyOpen, 2013, 2, 684-694.	3.0	26
31	A Reverse Transcriptase-Cas1 Fusion Protein Contains a Cas6 Domain Required for Both CRISPR RNA Biogenesis and RNA Spacer Acquisition. Molecular Cell, 2018, 72, 700-714.e8.	9.7	25
32	Regulation of polyphenol oxidase activities and melanin synthesis in Marinomonas mediterranea: identification of ppoS, a gene encoding a sensor histidine kinase a aThe GenBank accession number for the sequence reported in this paper is AF398464 Microbiology (United Kingdom), 2002, 148, 2457-2466.	1.8	23
33	Complete genome sequence of the melanogenic marine bacterium Marinomonas mediterranea type strain (MMB-1T) Standards in Genomic Sciences, 2012, 6, 63-73.	1.5	21
34	Distribution in Different Organisms of Amino Acid Oxidases with FAD or a Quinone As Cofactor and Their Role as Antimicrobial Proteins in Marine Bacteria. Marine Drugs, 2015, 13, 7403-7418.	4.6	21
35	Marinomonas mediterranea is a lysogenic bacterium that synthesizes R-bodies. Microbiology (United) Tj ETQq1	1 0,78431 1.8	4 rgBT /Over
36	Characterization of recombinant biosynthetic precursors of the cysteine tryptophylquinone cofactors of l-lysine-epsilon-oxidase and glycine oxidase from Marinomonas mediterranea. Biochimica Et Biophysica Acta - Proteins and Proteomics, 2015, 1854, 1123-1131.	2.3	20

#	Article	IF	Citations
37	LodB is required for the recombinant synthesis of the quinoprotein l-lysine-ε-oxidase from Marinomonas mediterranea. Applied Microbiology and Biotechnology, 2014, 98, 2981-2989.	3.6	19
38	Molecular cloning of a copper-dependent laccase from the dye-decolorizing strain Stenotrophomonas maltophilia AAP56. Journal of Applied Microbiology, 2011, 111, 1394-1405.	3.1	18
39	Structure and Enzymatic Properties of an Unusual Cysteine Tryptophylquinone-Dependent Glycine Oxidase from <i>Pseudoalteromonas luteoviolacea</i>). Biochemistry, 2018, 57, 1155-1165.	2.5	18
40	Roles of active site residues in LodA, a cysteine tryptophylquinone dependent Îμ-lysine oxidase. Archives of Biochemistry and Biophysics, 2015, 579, 26-32.	3.0	17
41	Regulation of the <i>Marinomonas mediterranea</i> Antimicrobial Protein Lysine Oxidase by <scp>l</scp> -Lysine and the Sensor Histidine Kinase PpoS. Applied and Environmental Microbiology, 2010, 76, 6141-6149.	3.1	16
42	Different recombinant forms of polyphenol oxidase A, a laccase from Marinomonas mediterranea. Protein Expression and Purification, 2016, 123, 60-69.	1.3	15
43	Steadyâ€state kinetic mechanism of LodA, a novel cysteine tryptophylquinoneâ€dependent oxidase. FEBS Letters, 2014, 588, 752-756.	2.8	12
44	Complete genome sequence of Marinomonas posidonica type strain (IVIA-Po-181T). Standards in Genomic Sciences, 2012, 7, 31-43.	1.5	10
45	Interaction of GoxA with Its Modifying Enzyme and Its Subunit Assembly Are Dependent on the Extent of Cysteine Tryptophylquinone Biosynthesis. Biochemistry, 2016, 55, 2305-2308.	2.5	10
46	The Pseudoalteromonas luteoviolacea L-amino Acid Oxidase with Antimicrobial Activity Is a Flavoenzyme. Marine Drugs, 2018, 16, 499.	4.6	9
47	Roles of Copper and a Conserved Aspartic Acid in the Autocatalytic Hydroxylation of a Specific Tryptophan Residue during Cysteine Tryptophylquinone Biogenesis. Biochemistry, 2017, 56, 997-1004.	2.5	7
48	A Small RNA Isolation and Sequencing Protocol and Its Application to Assay CRISPR RNA Biogenesis in Bacteria. Bio-protocol, 2018, 8, .	0.4	5
49	Dimethoxyphenol oxidase activity of different microbial blue multicopper proteins. FEMS Microbiology Letters, 2001, 204, 175-181.	1.8	5
50	A histidine kinase and a response regulator provide phage resistance to Marinomonas mediterranea via CRISPR-Cas regulation. Scientific Reports, 2021, 11, 20564.	3.3	5
51	Characterization of PlGoxB, a flavoprotein required for cysteine tryptophylquinone biosynthesis in glycine oxidase from Pseudoalteromonas luteoviolacea. Archives of Biochemistry and Biophysics, 2019, 674, 108110.	3.0	3
52	Bacterial Laccases: Some Recent Advances and Applications. Microbiology Monographs, 2020, , 27-55.	0.6	2
53	The protein encoded by the Shewanella colwelliana melA gene is a p-hydroxyphenylpyruvate dioxygenase. FEMS Microbiology Letters, 1994, 124, 179-184.	1.8	2
54	R-bodies., 2006,, 331-341.		2

#	÷	Article	lF	CITATIONS
5	5	Isolation of Phages Infecting Marinomonas mediterranea by an Enrichment Protocol. Bio-protocol, $2018, 8, .$	0.4	1