

Nina Åsterås

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1726007/publications.pdf>

Version: 2024-02-01

45
papers

1,178
citations

471371

17
h-index

395590

33
g-index

48
all docs

48
docs citations

48
times ranked

1499
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Critically appraised paper: Progressive exercise is not superior to best practice advice, and steroid injection is not superior to no injection, for rotator cuff disorders [synopsis]. <i>Journal of Physiotherapy</i> , 2022, 68, 71. | 0.7 | 0 |
| 2 | Development of radiographic classification criteria for hand osteoarthritis: a methodological report (Phase 2). <i>RMD Open</i> , 2022, 8, e002024. | 1.8 | 5 |
| 3 | Critically appraised paper: Exercise is safe, clinically effective and cost-effective compared to usual care after non-reconstructive breast cancer surgery [synopsis]. <i>Journal of Physiotherapy</i> , 2022, 68, 145-145. | 0.7 | 0 |
| 4 | A Framework to Guide the Development of Health Care Professional Education and Training in Best Evidence Osteoarthritis Care. <i>Clinics in Geriatric Medicine</i> , 2022, 38, 361-384. | 1.0 | 1 |
| 5 | Best Evidence Osteoarthritis Care. <i>Clinics in Geriatric Medicine</i> , 2022, 38, 287-302. | 1.0 | 4 |
| 6 | Interventions for osteoarthritis pain: A systematic review with network meta-analysis of existing Cochrane reviews. <i>Osteoarthritis and Cartilage Open</i> , 2022, 4, 100242. | 0.9 | 11 |
| 7 | 2022 EULAR points to consider for remote care in rheumatic and musculoskeletal diseases. <i>Annals of the Rheumatic Diseases</i> , 2022, 81, 1065-1071. | 0.5 | 54 |
| 8 | Critically appraised paper: Early surgery is not superior to exercise and education with the option of later surgery for meniscal tears in young adults [synopsis]. <i>Journal of Physiotherapy</i> , 2022, , . | 0.7 | 0 |
| 9 | Relationship between cam morphology, hip symptoms, and hip osteoarthritis: the Musculoskeletal pain in Ullersaker Study (MUST) cohort. <i>HIP International</i> , 2021, 31, 789-796. | 0.9 | 7 |
| 10 | Improving osteoarthritis management in primary healthcare: results from a quasi-experimental study. <i>BMC Musculoskeletal Disorders</i> , 2021, 22, 79. | 0.8 | 6 |
| 11 | Endorsement of the domains of knee and hip osteoarthritis (OA) flare: A report from the OMERACT 2020 inaugural virtual consensus vote from the flares in OA working group. <i>Seminars in Arthritis and Rheumatism</i> , 2021, 51, 618-622. | 1.6 | 12 |
| 12 | Critically appraised paper: Group-based pelvic floor muscle training is not inferior to individual training for the treatment of urinary incontinence in older women [synopsis]. <i>Journal of Physiotherapy</i> , 2021, 67, 219. | 0.7 | 0 |
| 13 | Critically appraised paper: Stable supportive shoes improved knee pain more than flat flexible shoes in people with moderate to severe radiographic medial knee osteoarthritis [synopsis]. <i>Journal of Physiotherapy</i> , 2021, 67, 310. | 0.7 | 0 |
| 14 | Low adherence to exercise may have influenced the proportion of OMERACT-OARSI responders in an integrated osteoarthritis care model: secondary analyses from a cluster-randomised stepped-wedge trial. <i>BMC Musculoskeletal Disorders</i> , 2020, 21, 236. | 0.8 | 11 |
| 15 | Implementing a structured model for osteoarthritis care in primary healthcare: A stepped-wedge cluster-randomised trial. <i>PLoS Medicine</i> , 2019, 16, e1002949. | 3.9 | 31 |
| 16 | The Maternal and Paternal Effects on Clinically and Surgically Defined Osteoarthritis. <i>Arthritis and Rheumatology</i> , 2019, 71, 1844-1848. | 2.9 | 9 |
| 17 | Implementing international osteoarthritis guidelines in primary care: uptake and fidelity among health professionals and patients. <i>Osteoarthritis and Cartilage</i> , 2019, 27, 1138-1147. | 0.6 | 14 |
| 18 | Does occupational therapy delay or reduce the proportion of patients that receives thumb carpometacarpal joint surgery? A multicentre randomised controlled trial. <i>RMD Open</i> , 2019, 5, e001046. | 1.8 | 16 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Measurement properties for the revised patient-reported OsteoArthritis Quality Indicator questionnaire. <i>Osteoarthritis and Cartilage</i> , 2018, 26, 1300-1310. | 0.6 | 12 |
| 20 | The Ottawa Panel guidelines on programmes involving therapeutic exercise for the management of hand osteoarthritis. <i>Clinical Rehabilitation</i> , 2018, 32, 026921551878097. | 1.0 | 13 |
| 21 | Distribution of osteoarthritis in a Norwegian population-based cohort: associations to risk factor profiles and health-related quality of life. <i>Rheumatology International</i> , 2017, 37, 1541-1550. | 1.5 | 6 |
| 22 | Exercise for Hand Osteoarthritis: A Cochrane Systematic Review. <i>Journal of Rheumatology</i> , 2017, 44, 1850-1858. | 1.0 | 41 |
| 23 | The importance of dose in land-based supervised exercise for people with hip osteoarthritis. A systematic review and meta-analysis. <i>Osteoarthritis and Cartilage</i> , 2017, 25, 1563-1576. | 0.6 | 53 |
| 24 | Quality of Community-Based Osteoarthritis Care: A Systematic Review and Meta-Analysis. <i>Arthritis Care and Research</i> , 2016, 68, 1443-1452. | 1.5 | 133 |
| 25 | Consensus on Exercise Reporting Template (CERT): Modified Delphi Study. <i>Physical Therapy</i> , 2016, 96, 1514-1524. | 1.1 | 279 |
| 26 | Patient-reported quality indicators for osteoarthritis: a patient and public generated self-report measure for primary care. <i>Research Involvement and Engagement</i> , 2016, 2, 5. | 1.1 | 22 |
| 27 | Implementing international osteoarthritis treatment guidelines in primary health care: study protocol for the SAMBA stepped wedge cluster randomized controlled trial. <i>Implementation Science</i> , 2015, 10, 165. | 2.5 | 14 |
| 28 | Perceived quality of health care services among people with osteoarthritis – results from a nationwide survey. <i>Patient Preference and Adherence</i> , 2015, 9, 1255. | 0.8 | 11 |
| 29 | Diabetes Is Associated With Increased Hand Pain in Erosive Hand Osteoarthritis: Data From a Population-Based Study. <i>Arthritis Care and Research</i> , 2015, 67, 187-195. | 1.5 | 58 |
| 30 | A tailored hand exercise program improves function of the rheumatoid hand [synopsis]. <i>Journal of Physiotherapy</i> , 2015, 61, 96. | 0.7 | 1 |
| 31 | Development of an evidence-based exercise programme for people with hand osteoarthritis. <i>Scandinavian Journal of Occupational Therapy</i> , 2015, 22, 103-116. | 1.1 | 18 |
| 32 | Quality of hip and knee osteoarthritis management in primary health care in a Norwegian county: a cross-sectional survey. <i>BMC Health Services Research</i> , 2014, 14, 598. | 0.9 | 18 |
| 33 | Exercise programme with telephone follow-up for people with hand osteoarthritis " protocol for a randomised controlled trial. <i>BMC Musculoskeletal Disorders</i> , 2014, 15, 82. | 0.8 | 10 |
| 34 | Limited effects of exercises in people with hand osteoarthritis: results from a randomized controlled trial. <i>Osteoarthritis and Cartilage</i> , 2014, 22, 1224-1233. | 0.6 | 43 |
| 35 | Hand, hip and knee osteoarthritis in a Norwegian population-based study - The MUST protocol. <i>BMC Musculoskeletal Disorders</i> , 2013, 14, 201. | 0.8 | 28 |
| 36 | Patient-Reported Quality of Care for Osteoarthritis: Development and Testing of the OsteoArthritis Quality Indicator Questionnaire. <i>Arthritis Care and Research</i> , 2013, 65, 1043-1051. | 1.5 | 42 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Daily use of a cane for two months reduced pain and improved function in patients with knee osteoarthritis. <i>Journal of Physiotherapy</i> , 2012, 58, 128. | 0.7 | 10 |
| 38 | Quantifying Information Content in Survey Data by Entropy. <i>Entropy</i> , 2010, 12, 161-163. | 1.1 | 9 |
| 39 | Structured functional assessments in general practice increased the use of part-time sick leave: A cluster randomised controlled trial. <i>Scandinavian Journal of Public Health</i> , 2010, 38, 192-199. | 1.2 | 10 |
| 40 | Tai Chi reduces pain and improves physical function for people with knee OA. <i>Journal of Physiotherapy</i> , 2010, 56, 57. | 0.7 | 0 |
| 41 | Implementing structured functional assessments in general practice for persons with long-term sick leave: a cluster randomised controlled trial. <i>BMC Family Practice</i> , 2009, 10, 31. | 2.9 | 22 |
| 42 | Effect of an intervention addressing working technique on the biomechanical load of the neck and shoulders among hairdressers. <i>Applied Ergonomics</i> , 2008, 39, 183-190. | 1.7 | 48 |
| 43 | A randomised comparison of a four- and a five-point scale version of the Norwegian Function Assessment Scale. <i>Health and Quality of Life Outcomes</i> , 2008, 6, 14. | 1.0 | 43 |
| 44 | Functional ability in a population: normative survey data and reliability for the ICF based Norwegian Function Assessment Scale. <i>BMC Public Health</i> , 2007, 7, 278. | 1.2 | 33 |
| 45 | Muscle pain, physical activity, self-efficacy and relaxation ability in adolescents. <i>Advances in Physiotherapy</i> , 2006, 8, 33-40. | 0.2 | 11 |