

Daniel F Legler

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1729917/daniel-f-legler-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

85
papers

6,390
citations

34
h-index

79
g-index

88
ext. papers

7,302
ext. citations

7.7
avg, IF

5.49
L-index

#	Paper	IF	Citations
85	Mechanosensitive ACKR4 scavenges CCR7 chemokines to facilitate T cell de-adhesion and passive transport by flow in inflamed afferent lymphatics.. <i>Cell Reports</i> , 2022 , 38, 110334	10.6	1
84	CD44 engagement enhances acute myeloid leukemia cell adhesion to the bone marrow microenvironment by increasing VLA-4 avidity. <i>Haematologica</i> , 2021 , 106, 2102-2113	6.6	7
83	Delineating the interactions between the cannabinoid CB receptor and its regulatory effectors; β Arrestins and G protein-coupled receptor kinases. <i>British Journal of Pharmacology</i> , 2021 ,	8.6	2
82	A Versatile Toolkit for Semi-Automated Production of Fluorescent Chemokines to Study CCR7 Expression and Functions. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	3
81	Keratinocytes control skin immune homeostasis through de novo-synthesized glucocorticoids. <i>Science Advances</i> , 2021 , 7,	14.3	8
80	CCR7 signalosomes are preassembled on tips of lymphocyte microvilli in proximity to LFA-1. <i>Biophysical Journal</i> , 2021 , 120, 4002-4012	2.9	0
79	CAL-1 as Cellular Model System to Study CCR7-Guided Human Dendritic Cell Migration. <i>Frontiers in Immunology</i> , 2021 , 12, 702453	8.4	0
78	Medullary stromal cells synergize their production and capture of CCL21 for T-cell emigration from neonatal mouse thymus. <i>Blood Advances</i> , 2021 , 5, 99-112	7.8	4
77	Elimination of negative feedback in TLR signalling allows rapid and hypersensitive detection of microbial contaminants.. <i>Scientific Reports</i> , 2021 , 11, 24414	4.9	1
76	ACKR4 Recruits GRK3 Prior to β Arrestins but Can Scavenge Chemokines in the Absence of β Arrestins. <i>Frontiers in Immunology</i> , 2020 , 11, 720	8.4	12
75	Membrane Compartmentalization and Scaffold Proteins in Leukocyte Migration. <i>Frontiers in Cell and Developmental Biology</i> , 2020 , 8, 285	5.7	2
74	FAT10 localises in dendritic cell aggresome-like induced structures and contributes to their disassembly. <i>Journal of Cell Science</i> , 2020 , 133,	5.3	2
73	CCL20 is a novel ligand for the scavenging atypical chemokine receptor 4. <i>Journal of Leukocyte Biology</i> , 2020 , 107, 1137-1154	6.5	12
72	β Arrestin1 and β Arrestin2 Are Required to Support the Activity of the CXCL12/HMGB1 Heterocomplex on CXCR4. <i>Frontiers in Immunology</i> , 2020 , 11, 550824	8.4	4
71	B cell zone reticular cell microenvironments shape CXCL13 gradient formation. <i>Nature Communications</i> , 2020 , 11, 3677	17.4	18
70	CCR5 deficiency/CCR5B2: resistant to HIV infection at the cost of curtailed CD4 T cell memory responses. <i>EMBO Journal</i> , 2020 , 39, e105854	13	4
69	CXCL14 Preferentially Synergizes With Homeostatic Chemokine Receptor Systems. <i>Frontiers in Immunology</i> , 2020 , 11, 561404	8.4	2

68	Biased Signaling of CCL21 and CCL19 Does Not Rely on N-Terminal Differences, but Markedly on the Chemokine Core Domains and Extracellular Loop 2 of CCR7. <i>Frontiers in Immunology</i> , 2019 , 10, 2156	8.4	7
67	Engineering of Nanobodies Recognizing the Human Chemokine Receptor CCR7. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	4
66	IL-4 receptor engagement in human neutrophils impairs their migration and extracellular trap formation. <i>Journal of Allergy and Clinical Immunology</i> , 2019 , 144, 267-279.e4	11.5	36
65	Chemokine Receptor CCR7 Triggers an Endomembrane Signaling Complex for Spatial Rac Activation. <i>Cell Reports</i> , 2019 , 29, 995-1009.e6	10.6	15
64	In Vivo Function of the Lipid Raft Protein Flotillin-1 during CD8 T Cell-Mediated Host Surveillance. <i>Journal of Immunology</i> , 2019 , 203, 2377-2387	5.3	6
63	Beyond migration-Chemokines in lymphocyte priming, differentiation, and modulating effector functions. <i>Journal of Leukocyte Biology</i> , 2018 , 104, 301-312	6.5	17
62	Membrane lipid environment: Potential modulation of chemokine receptor function. <i>Cytokine</i> , 2018 , 109, 72-75	4	5
61	A unique signal sequence of the chemokine receptor CCR7 promotes package into COPII vesicles for efficient receptor trafficking. <i>Journal of Leukocyte Biology</i> , 2018 , 104, 375-389	6.5	6
60	ZAP70 expression enhances chemokine-driven chronic lymphocytic leukemia cell migration and arrest by valency regulation of integrins. <i>FASEB Journal</i> , 2018 , 32, 4824-4835	0.9	11
59	New insights in chemokine signaling. <i>F1000Research</i> , 2018 , 7, 95	3.6	43
58	CCR7 Is Recruited to the Immunological Synapse, Acts as Co-stimulatory Molecule and Drives LFA-1 Clustering for Efficient T Cell Adhesion Through ZAP70. <i>Frontiers in Immunology</i> , 2018 , 9, 3115	8.4	15
57	A structure-activity relationship linking non-planar PCBs to functional deficits of neural crest cells: new roles for connexins. <i>Archives of Toxicology</i> , 2018 , 92, 1225-1247	5.8	9
56	Role of Mechanotransduction and Tension in T Cell Function. <i>Frontiers in Immunology</i> , 2018 , 9, 2638	8.4	64
55	Fluorescently Tagged CCL19 and CCL21 to Monitor CCR7 and ACKR4 Functions. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	16
54	CCL19 with CCL21-tail displays enhanced glycosaminoglycan binding with retained chemotactic potency in dendritic cells. <i>Journal of Leukocyte Biology</i> , 2018 , 104, 401-411	6.5	10
53	Modulation of Chemokine Receptor Function by Cholesterol: New Prospects for Pharmacological Intervention. <i>Molecular Pharmacology</i> , 2017 , 91, 331-338	4.3	27
52	Epithelial chemokine CXCL14 synergizes with CXCL12 allosteric modulation of CXCR4. <i>FASEB Journal</i> , 2017 , 31, 3084-3097	0.9	34
51	The STEAP1(262-270) peptide encapsulated into PLGA microspheres elicits strong cytotoxic T cell immunity in HLA-A*0201 transgenic mice--A new approach to immunotherapy against prostate carcinoma. <i>Prostate</i> , 2016 , 76, 456-68	4.2	9

50	Inflammation-Induced CCR7 Oligomers Form Scaffolds to Integrate Distinct Signaling Pathways for Efficient Cell Migration. <i>Immunity</i> , 2016 , 44, 59-72	32.3	65
49	Distinct CCR7 glycosylation pattern shapes receptor signaling and endocytosis to modulate chemotactic responses. <i>Journal of Leukocyte Biology</i> , 2016 , 99, 993-1007	6.5	46
48	Chemokine axes in breast cancer: factors of the tumor microenvironment reshape the CCR7-driven metastatic spread of luminal-A breast tumors. <i>Journal of Leukocyte Biology</i> , 2016 , 99, 1009-25	6.5	21
47	Common and biased signaling pathways of the chemokine receptor CCR7 elicited by its ligands CCL19 and CCL21 in leukocytes. <i>Journal of Leukocyte Biology</i> , 2016 , 99, 869-82	6.5	102
46	Chemokines: Chemistry, Biochemistry and Biological Function. <i>Chimia</i> , 2016 , 70, 856-859	1.3	41
45	On the move: endocytic trafficking in cell migration. <i>Cellular and Molecular Life Sciences</i> , 2015 , 72, 2119-34	3.3	59
44	Analysis of CCR7 mediated T cell transfectant migration using a microfluidic gradient generator. <i>Journal of Immunological Methods</i> , 2015 , 419, 9-17	2.5	4
43	Loss of GM130 in breast cancer cells and its effects on cell migration, invasion and polarity. <i>Cell Cycle</i> , 2015 , 14, 1139-47	4.7	13
42	In vivo TCR Signaling in CD4(+) T Cells Imprints a Cell-Intrinsic, Transient Low-Motility Pattern Independent of Chemokine Receptor Expression Levels, or Microtubular Network, Integrin, and Protein Kinase C Activity. <i>Frontiers in Immunology</i> , 2015 , 6, 297	8.4	5
41	Loss of Gadkin Affects Dendritic Cell Migration In Vitro. <i>PLoS ONE</i> , 2015 , 10, e0143883	3.7	11
40	Regulation of Sec16 levels and dynamics links proliferation and secretion. <i>Journal of Cell Science</i> , 2015 , 128, 670-82	5.3	22
39	CCR7: roles in cancer cell dissemination, migration and metastasis formation. <i>International Journal of Biochemistry and Cell Biology</i> , 2014 , 54, 78-82	5.6	49
38	Interstitial dendritic cell guidance by haptotactic chemokine gradients. <i>Science</i> , 2013 , 339, 328-32	33.3	365
37	Distinct modulation of chemokine expression patterns in human monocyte-derived dendritic cells by prostaglandin E(2). <i>Cellular Immunology</i> , 2012 , 276, 52-8	4.4	16
36	Converse regulation of CCR7-driven human dendritic cell migration by prostaglandin E ₂ and liver X receptor activation. <i>European Journal of Immunology</i> , 2012 , 42, 2949-58	6.1	23
35	Ubiquitylation of the chemokine receptor CCR7 enables efficient receptor recycling and cell migration. <i>Journal of Cell Science</i> , 2012 , 125, 4463-74	5.3	34
34	Cross-talk between TCR and CCR7 signaling sets a temporal threshold for enhanced T lymphocyte migration. <i>Journal of Immunology</i> , 2011 , 187, 5645-52	5.3	29
33	Definition of key variables for the induction of optimal NY-ESO-1-specific T cells in HLA transgene mice. <i>Journal of Immunology</i> , 2010 , 185, 3445-55	5.3	8

32	Prostaglandin E2 at new glance: novel insights in functional diversity offer therapeutic chances. <i>International Journal of Biochemistry and Cell Biology</i> , 2010 , 42, 198-201	5.6	170
31	Immobilized chemokine fields and soluble chemokine gradients cooperatively shape migration patterns of dendritic cells. <i>Immunity</i> , 2010 , 32, 703-13	32.3	232
30	Soluble CD146 is generated by ectodomain shedding of membrane CD146 in a calcium-induced, matrix metalloprotease-dependent process. <i>Microvascular Research</i> , 2009 , 78, 325-31	3.7	32
29	Prostaglandin E(2) enhances T-cell proliferation by inducing the costimulatory molecules OX40L, CD70, and 4-1BBL on dendritic cells. <i>Blood</i> , 2009 , 113, 2451-60	2.2	80
28	Increased mobility of major histocompatibility complex I-peptide complexes decreases the sensitivity of antigen recognition. <i>Journal of Biological Chemistry</i> , 2008 , 283, 24254-63	5.4	18
27	Reduced expression of cyclooxygenase-2 in primary breast cancer. <i>Journal of the National Cancer Institute</i> , 2008 , 100, 1042-3	9.7	5
26	Distinct motifs in the chemokine receptor CCR7 regulate signal transduction, receptor trafficking and chemotaxis. <i>Journal of Cell Science</i> , 2008 , 121, 2759-67	5.3	41
25	V domain of RAGE interacts with AGEs on prostate carcinoma cells. <i>Prostate</i> , 2008 , 68, 748-58	4.2	38
24	A novel cytosolic class I antigen-processing pathway for endoplasmic-reticulum-targeted proteins. <i>EMBO Reports</i> , 2007 , 8, 945-51	6.5	12
23	Prostaglandin E2 is a key factor for monocyte-derived dendritic cell maturation: enhanced T cell stimulatory capacity despite IDO. <i>Journal of Leukocyte Biology</i> , 2007 , 82, 1106-14	6.5	53
22	Preformed reggie/flotillin caps: stable priming platforms for macrodomain assembly in T cells. <i>FASEB Journal</i> , 2006 , 20, 711-3	0.9	45
21	Opposite fate of endocytosed CCR7 and its ligands: recycling versus degradation. <i>Journal of Immunology</i> , 2006 , 177, 2314-23	5.3	97
20	Prostaglandin E2 is generally required for human dendritic cell migration and exerts its effect via EP2 and EP4 receptors. <i>Journal of Immunology</i> , 2006 , 176, 966-73	5.3	159
19	Posttranscriptional regulation of Fas (CD95) ligand killing activity by lipid rafts. <i>Blood</i> , 2006 , 107, 2790-6	2.2	27
18	Differential insertion of GPI-anchored GFPs into lipid rafts of live cells. <i>FASEB Journal</i> , 2005 , 19, 73-5	0.9	102
17	PrPc capping in T cells promotes its association with the lipid raft proteins reggie-1 and reggie-2 and leads to signal transduction. <i>FASEB Journal</i> , 2004 , 18, 1731-3	0.9	118
16	The alpha v beta 3 integrin as a tumor homing ligand for lymphocytes. <i>European Journal of Immunology</i> , 2004 , 34, 1608-16	6.1	25
15	CCL19/CCL21-triggered signal transduction and migration of dendritic cells requires prostaglandin E2. <i>Blood</i> , 2004 , 103, 1595-601	2.2	195

14	The beta1 and beta3 integrins promote T cell receptor-mediated cytotoxic T lymphocyte activation. <i>Journal of Biological Chemistry</i> , 2003 , 278, 26983-91	5.4	55
13	Soluble major histocompatibility complex-peptide octamers with impaired CD8 binding selectively induce Fas-dependent apoptosis. <i>Journal of Biological Chemistry</i> , 2003 , 278, 4500-9	5.4	39
12	Recruitment of TNF receptor 1 to lipid rafts is essential for TNFalpha-mediated NF-kappaB activation. <i>Immunity</i> , 2003 , 18, 655-64	32.3	369
11	Association of the Epstein-Barr virus latent membrane protein 1 with lipid rafts is mediated through its N-terminal region. <i>Cellular and Molecular Life Sciences</i> , 2002 , 59, 171-80	10.3	21
10	CARMA1 is a critical lipid raft-associated regulator of TCR-induced NF-kappa B activation. <i>Nature Immunology</i> , 2002 , 3, 836-43	19.1	301
9	Changing responsiveness to chemokines allows medullary plasmablasts to leave lymph nodes. <i>European Journal of Immunology</i> , 2001 , 31, 609-16	6.1	100
8	CTL activation is induced by cross-linking of TCR/MHC-peptide-CD8/p56lck adducts in rafts. <i>European Journal of Immunology</i> , 2001 , 31, 1561-70	6.1	37
7	Selective inhibition of CTL activation by a dipalmitoyl-phospholipid that prevents the recruitment of signaling molecules to lipid rafts. <i>FASEB Journal</i> , 2001 , 15, 1601-3	0.9	23
6	Activation-dependent modulation of B lymphocyte migration to chemokines. <i>International Immunology</i> , 2000 , 12, 1285-92	4.9	75
5	The chemokine SLC is expressed in T cell areas of lymph nodes and mucosal lymphoid tissues and attracts activated T cells via CCR7. <i>European Journal of Immunology</i> , 1998 , 28, 2025-34	6.1	265
4	B cell-attracting chemokine 1, a human CXC chemokine expressed in lymphoid tissues, selectively attracts B lymphocytes via BLR1/CXCR5. <i>Journal of Experimental Medicine</i> , 1998 , 187, 655-60	16.6	631
3	Identification of CCR8, the receptor for the human CC chemokine I-309. <i>Journal of Biological Chemistry</i> , 1997 , 272, 17251-4	5.4	144
2	Expression of high- and low-affinity receptors for C3a on the human mast cell line, HMC-1. <i>European Journal of Immunology</i> , 1996 , 26, 753-8	6.1	72
1	The CXC chemokine SDF-1 is the ligand for LESTR/fusin and prevents infection by T-cell-line-adapted HIV-1. <i>Nature</i> , 1996 , 382, 833-5	50.4	1478