

Wenhua Zhang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1738735/publications.pdf>

Version: 2024-02-01

39
papers

1,868
citations

331259

21
h-index

301761

39
g-index

41
all docs

41
docs citations

41
times ranked

2314
citing authors

#	ARTICLE	IF	CITATIONS
1	Emerging roles of phosphoinositide-associated membrane trafficking in plant stress responses. <i>Journal of Genetics and Genomics</i> , 2022, 49, 726-734.	1.7	4
2	Molecular cloning and functional characterization of GmAAPTs from soybean (<i>Glycine max</i>). <i>Plant Signaling and Behavior</i> , 2021, 16, 1845048.	1.2	1
3	The rice aldehyde oxidase OsAO3 gene regulates plant growth, grain yield, and drought tolerance by participating in ABA biosynthesis. <i>Biochemical and Biophysical Research Communications</i> , 2021, 548, 189-195.	1.0	25
4	Rice shaker potassium channel <i>OsAKT2</i> positively regulates salt tolerance and grain yield by mediating K ⁺ redistribution. <i>Plant, Cell and Environment</i> , 2021, 44, 2951-2965.	2.8	41
5	Seed specifically over-expressing DGAT2A enhances oil and linoleic acid contents in soybean seeds. <i>Biochemical and Biophysical Research Communications</i> , 2021, 568, 143-150.	1.0	14
6	A bHLH protein, OsBIM1, positively regulates rice leaf angle by promoting brassinosteroid signaling. <i>Biochemical and Biophysical Research Communications</i> , 2021, 578, 129-135.	1.0	9
7	HSP70-3 Interacts with Phospholipase D γ and Participates in Heat Stress Defense. <i>Plant Physiology</i> , 2021, 185, 1148-1165.	2.3	27
8	An endoplasmic reticulum-localized cytochrome <i>b₅</i> regulates high-affinity K ⁺ transport in response to salt stress in rice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	3.3	27
9	Overexpression of soybean GmPLD β enhances seed oil content and modulates fatty acid composition in transgenic Arabidopsis. <i>Plant Science</i> , 2020, 290, 110298.	1.7	14
10	Involvement of Arabidopsis phospholipase D γ in regulation of ROS-mediated microtubule organization and stomatal movement upon heat shock. <i>Journal of Experimental Botany</i> , 2020, 71, 6555-6570.	2.4	29
11	Phosphatidic acid directly binds with rice potassium channel <i>OsAKT2</i> to inhibit its activity. <i>Plant Journal</i> , 2020, 102, 649-665.	2.8	30
12	The ATP-binding cassette transporter <i>OsPDR1</i> regulates plant growth and pathogen resistance by affecting jasmonates biosynthesis in rice. <i>Plant Science</i> , 2020, 298, 110582.	1.7	17
13	Multiple basic amino acid residues contribute to phosphatidic acid-mediated inhibition of rice potassium channel <i>OsAKT2</i> . <i>Plant Signaling and Behavior</i> , 2020, 15, 1789818.	1.2	2
14	Phosphatidic acid promotes the activation and plasma membrane localization of MKK7 and MKK9 in response to salt stress. <i>Plant Science</i> , 2019, 287, 110190.	1.7	37
15	Tissue-specific accumulation of pH-sensing phosphatidic acid determines plant stress tolerance. <i>Nature Plants</i> , 2019, 5, 1012-1021.	4.7	73
16	Genome-wide analysis and functional characterization of Acyl-CoA:diacylglycerol acyltransferase from soybean identify GmDGAT1A and 1B roles in oil synthesis in Arabidopsis seeds. <i>Journal of Plant Physiology</i> , 2019, 242, 153019.	1.6	24
17	Rice <i>qGL3/OsPPKL1</i> Functions with the GSK3/SHAGGY-Like Kinase <i>OsGSK3</i> to Modulate Brassinosteroid Signaling. <i>Plant Cell</i> , 2019, 31, 1077-1093.	3.1	106
18	Phosphatidic Acid Directly Regulates PINOID-Dependent Phosphorylation and Activation of the PIN-FORMED2 Auxin Efflux Transporter in Response to Salt Stress. <i>Plant Cell</i> , 2019, 31, 250-271.	3.1	97

#	ARTICLE	IF	CITATIONS
19	FLOURY SHRUNKEN ENDOSPERM1 Connects Phospholipid Metabolism and Amyloplast Development in Rice. <i>Plant Physiology</i> , 2018, 177, 698-712.	2.3	35
20	A phosphoinositide-specific phospholipase C pathway elicits stress-induced Ca ²⁺ signals and confers salt tolerance to rice. <i>New Phytologist</i> , 2017, 214, 1172-1187.	3.5	85
21	Peroxisomal CuAO and its product H ₂ O ₂ regulate the distribution of auxin and IBA-dependent lateral root development in Arabidopsis. <i>Journal of Experimental Botany</i> , 2017, 68, 4851-4867.	2.4	33
22	Phospholipase D1 negatively regulates plant thermotolerance by destabilizing cortical microtubules in Arabidopsis. <i>Plant, Cell and Environment</i> , 2017, 40, 2220-2235.	2.8	45
23	Phosphatidic acid binds to and regulates guanine nucleotide exchange factor 8 (GEF8) activity in Arabidopsis. <i>Functional Plant Biology</i> , 2017, 44, 1029.	1.1	7
24	Fine mapping of qSKC-1, a major quantitative trait locus for shoot K ⁺ concentration, in rice seedlings grown under salt stress. <i>Breeding Science</i> , 2017, 67, 286-295.	0.9	11
25	Characterization and Fine Mapping of a Rice Leaf-Rolling Mutant Deficient in Commissural Veins. <i>Crop Science</i> , 2017, 57, 2595-2604.	0.8	2
26	Small interfering RNAs from bidirectional transcripts of GhMML3_A12 regulate cotton fiber development. <i>New Phytologist</i> , 2016, 210, 1298-1310.	3.5	124
27	Arabidopsis thaliana constitutively active ROP11 interacts with the NADPH oxidase respiratory burst oxidase homologue F to regulate reactive oxygen species production in root hairs. <i>Functional Plant Biology</i> , 2016, 43, 221.	1.1	14
28	Comparative Study of Early Cold-Regulated Proteins by Two-Dimensional Difference Gel Electrophoresis Reveals a Key Role for Phospholipase D1 in Mediating Cold Acclimation Signaling Pathway in Rice. <i>Molecular and Cellular Proteomics</i> , 2016, 15, 1397-1411.	2.5	50
29	Regulation of developmental and environmental signaling by interaction between microtubules and membranes in plant cells. <i>Protein and Cell</i> , 2016, 7, 81-88.	4.8	16
30	The role of putrescine in the regulation of proteins and fatty acids of thylakoid membranes under salt stress. <i>Scientific Reports</i> , 2015, 5, 14390.	1.6	95
31	The potassium transporter HAK21 functions in the maintenance of ion homeostasis and tolerance to salt stress in rice. <i>Plant, Cell and Environment</i> , 2015, 38, 2766-2779.	2.8	155
32	Identification and Fine Mapping of a Mutation Conferring Salt Sensitivity in Rice (<i>Oryza sativa</i>) Tj ETQq0.0.0.rgBT /Overlock 10	0.8	11
33	The Rice High-Affinity Potassium Transporter1;1 Is Involved in Salt Tolerance and Regulated by an MYB-Type Transcription Factor. <i>Plant Physiology</i> , 2015, 168, 1076-1090.	2.3	206
34	A CC-NBS-LRR type gene GHNTR1 confers resistance to southern root-knot nematode in <i>Nicotiana benthamiana</i> and <i>Nicotiana tabacum</i> . <i>European Journal of Plant Pathology</i> , 2015, 142, 715-729.	0.8	11
35	Physiological characterisation and fine mapping of a salt-tolerant mutant in rice (<i>Oryza sativa</i>). <i>Functional Plant Biology</i> , 2015, 42, 1026.	1.1	22
36	The mitogen-activated protein kinase cascade MKK1-MPK4 mediates salt signaling in rice. <i>Plant Science</i> , 2014, 227, 181-189.	1.7	70

#	ARTICLE	IF	CITATIONS
37	Phosphatidic Acid Regulates Microtubule Organization by Interacting with MAP65-1 in Response to Salt Stress in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2012, 24, 4555-4576.	3.1	219
38	Quantitative dissection of lipid degradation in rice seeds during accelerated aging. <i>Plant Growth Regulation</i> , 2012, 66, 49-58.	1.8	41
39	Phospholipase D in the signaling networks of plant response to abscisic acid and reactive oxygen species. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2005, 1736, 1-9.	1.2	38