

# Genichiro Ishii

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1742837/publications.pdf>

Version: 2024-02-01

117  
papers

5,132  
citations

117453

34  
h-index

98622

67  
g-index

119  
all docs

119  
docs citations

119  
times ranked

6539  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Lactic acid promotes PD-1 expression in regulatory T cells in highly glycolytic tumor microenvironments. <i>Cancer Cell</i> , 2022, 40, 201-218.e9.   | 7.7 | 266       |
| 2  | FDG uptake in PET is associated with the tumor microenvironment in metastatic lymph nodes and prognosis in N2 lung adenocarcinoma. <i>Cancer Science</i> , 2022, , .  | 1.7 | 3         |
| 3  | Cancer-associated fibroblasts and the tumor microenvironment in non-small cell lung cancer. <i>Expert Review of Anticancer Therapy</i> , 2022, 22, 169-182.   | 1.1 | 7         |
| 4  | Pathologic method for extracting good prognosis group in triple-negative breast cancer after neoadjuvant chemotherapy. <i>Cancer Science</i> , 2022, 113, 1507-1518.  | 1.7 | 6         |
| 5  | Component with abundant immune-related cells in combined hepatocellular cholangiocarcinoma identified by cluster analysis. <i>Cancer Science</i> , 2022, , .  | 1.7 | 3         |
| 6  | Histological tumor necrosis in pancreatic cancer after neoadjuvant therapy. <i>Oncology Reports</i> , 2022, 48, .   | 1.2 | 2         |
| 7  | Prognostic impact of the number of peri-tumoral alveolar macrophages in patients with stage I lung adenocarcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2022, 148, 3437-3447.                | 1.2 | 3         |
| 8  | Tumor-Infiltrating T Cells Concurrently Overexpress CD200R with Immune Checkpoints PD-1, CTLA-4, and TIM-3 in Non-Small-Cell Lung Cancer. <i>Pathobiology</i> , 2021, 88, 218-227.                                | 1.9 | 2         |
| 9  | Evaluation of the morphological features and unfavorable prognostic impact of dirty necrosis in renal cell carcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2021, 147, 1089-1100.             | 1.2 | 5         |
| 10 | Prognostic impact of the tumor immune microenvironment in pulmonary pleomorphic carcinoma. <i>Lung Cancer</i> , 2021, 153, 56-65.   | 0.9 | 7         |
| 11 | Relationship between podoplanin-expressing cancer-associated fibroblasts and the immune microenvironment of early lung squamous cell carcinoma. <i>Lung Cancer</i> , 2021, 153, 1-10.                             | 0.9 | 43        |
| 12 | Pathological features and prognostic implications of ground-glass opacity components on computed tomography for clinical stage I lung adenocarcinoma. <i>Surgery Today</i> , 2021, 51, 1188-1202.                 | 0.7 | 9         |
| 13 | Sarcomatoid hepatocellular carcinoma is distinct from ordinary hepatocellular carcinoma: Clinicopathologic, transcriptomic and immunologic analyses. <i>International Journal of Cancer</i> , 2021, 149, 546-560. | 2.3 | 18        |
| 14 | Impact of previous history of choledochojejunostomy on the incidence of organ/space surgical site infection after hepatectomy. <i>Asian Journal of Surgery</i> , 2021, 44, 1520-1520.                             | 0.2 | 0         |
| 15 | Prognostic impact of extranodal extension in patients with pN1-N2 lung adenocarcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2021, 147, 3699-3707.  | 1.2 | 6         |
| 16 | Drug-exposed cancer-associated fibroblasts facilitate gastric cancer cell progression following chemotherapy. <i>Gastric Cancer</i> , 2021, 24, 810-822.  | 2.7 | 8         |
| 17 | The immunological impact of preoperative chemoradiotherapy on the tumor microenvironment of pancreatic cancer. <i>Cancer Science</i> , 2021, 112, 2895-2904.  | 1.7 | 9         |
| 18 | Correlation between the number of viable tumor cells and immune cells in the tumor microenvironment in non-small cell lung cancer after induction therapy. <i>Pathology International</i> , 2021, 71, 512-520.    | 0.6 | 0         |

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 19 | High proportion of tumor necrosis predicts poor survival in surgically resected high-grade neuroendocrine carcinoma of the lung. <i>Lung Cancer</i> , 2021, 157, 1-8.  | 0.9  | 3         |
| 20 | Predictive markers based on transcriptome modules for vinorelbine-based adjuvant chemotherapy for lung adenocarcinoma patients. <i>Lung Cancer</i> , 2021, 158, 115-125.   | 0.9  | 2         |
| 21 | Clinicopathological, gene expression and genetic features of stage I lung adenocarcinoma with necrosis. <i>Lung Cancer</i> , 2021, 159, 74-83.   | 0.9  | 5         |
| 22 | The CLIP1-LTK fusion is an oncogenic driver in non-small cell lung cancer. <i>Nature</i> , 2021, 600, 319-323.   | 13.7 | 37        |
| 23 | Highly immunogenic cancer cells require activation of the WNT pathway for immunological escape. <i>Science Immunology</i> , 2021, 6, eabc6424.   | 5.6  | 64        |
| 24 | Cancer cell niche factors secreted from cancer-associated fibroblast by loss of H3K27me3. <i>Gut</i> , 2020, 69, 243-251.  | 6.1  | 62        |
| 25 | Machine learning-based histological classification that predicts recurrence of peripheral lung squamous cell carcinoma. <i>Lung Cancer</i> , 2020, 147, 252-258.   | 0.9  | 12        |
| 26 | Review of cancer-associated fibroblasts and their microenvironment in post-chemotherapy recurrence. <i>Human Cell</i> , 2020, 33, 938-945.   | 1.2  | 10        |
| 27 | The PD-1 expression balance between effector and regulatory T cells predicts the clinical efficacy of PD-1 blockade therapies. <i>Nature Immunology</i> , 2020, 21, 1346-1358.   | 7.0  | 431       |
| 28 | Low-dose CT lung cancer screening in never-smokers and smokers: results of an eight-year observational study. <i>Translational Lung Cancer Research</i> , 2020, 9, 10-22.  | 1.3  | 30        |
| 29 | Association between the mutational smoking signature and the immune microenvironment in lung adenocarcinoma. <i>Lung Cancer</i> , 2020, 147, 12-20.  | 0.9  | 5         |
| 30 | Optimal method for measuring invasive size that predicts survival in invasive mucinous adenocarcinoma of the lung. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020, 146, 1291-1298.   | 1.2  | 6         |
| 31 | Fibroblasts-dependent invasion of podoplanin-positive cancer stem cells in squamous cell carcinoma. <i>Journal of Cellular Physiology</i> , 2020, 235, 7251-7260.  | 2.0  | 5         |
| 32 | Blockade of EGFR improves responsiveness to PD-1 blockade in EGFR-mutated non-small cell lung cancer. <i>Science Immunology</i> , 2020, 5, .   | 5.6  | 160       |
| 33 | Uptake of collagen type I via macropinocytosis cause mTOR activation and anti-cancer drug resistance. <i>Biochemical and Biophysical Research Communications</i> , 2020, 526, 191-198.   | 1.0  | 19        |
| 34 | Clinical Utility of Histological and Radiological Evaluations of Tumor Necrosis for Predicting Prognosis in Pancreatic Cancer. <i>Pancreas</i> , 2020, 49, 634-641.  | 0.5  | 12        |
| 35 | Feasibility and utility of transbronchial cryobiopsy in precision medicine for lung cancer: Prospective single-arm study. <i>Cancer Science</i> , 2020, 111, 2488-2498.  | 1.7  | 46        |
| 36 | Randomized phase III study of irinotecan/cisplatin (IP) versus etoposide/cisplatin (EP) for completely resected high-grade neuroendocrine carcinoma (HGNEC) of the lung: JCOG1205/1206. <i>Journal of Clinical Oncology</i> , 2020, 38, 9006-9006. | 0.8  | 3         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Impact of SWI/SNF complex mutations in patients with non-small cell lung cancer (NSCLC) treated with immune checkpoint inhibitors: Immuno-oncology biomarker study in LC-SCRUM-Japan (LC-SCRUM-IBIS).. Journal of Clinical Oncology, 2020, 38, 9530-9530.          | 0.8 | 2         |
| 38 | Large pulmonary sclerosing pneumocytoma with massive necrosis and vascular invasion: a case report. Oxford Medical Case Reports, 2019, 2019, .   | 0.2 | 6         |
| 39 | Secretion of high amounts of hepatocyte growth factor is a characteristic feature of cancer-associated fibroblasts with EGFR-TKI resistance-promoting phenotype: A study of 18 cases of cancer-associated fibroblasts. Pathology International, 2019, 69, 472-480. | 0.6 | 15        |
| 40 | Ground-Glass Opacity Is a Strong Prognosticator for Pathologic Stage IA Lung Adenocarcinoma. Annals of Thoracic Surgery, 2019, 108, 249-255.   | 0.7 | 47        |
| 41 | Prognostic Impact of the Number of Metastatic Lymph Nodes on the Eighth Edition of the TNM Classification of NSCLC. Journal of Thoracic Oncology, 2019, 14, 1408-1418.   | 0.5 | 43        |
| 42 | Proportion of goblet cell is associated with malignant potential in invasive mucinous adenocarcinoma of the lung. Pathology International, 2019, 69, 526-535.  | 0.6 | 2         |
| 43 | Non-small cell lung cancer with loss of expression of the SWI/SNF complex is associated with aggressive clinicopathological features, PD-L1-positive status, and high tumor mutation burden. Lung Cancer, 2019, 138, 35-42.  | 0.9 | 53        |
| 44 | Interaction between cancer cells and cancer-associated fibroblasts after cisplatin treatment promotes cancer cell regrowth. Human Cell, 2019, 32, 453-464.   | 1.2 | 7         |
| 45 | Correlation between maximum standardized uptake values on FDG-PET and microenvironmental factors in patients with clinical stage IA radiologic pure-solid lung adenocarcinoma. Lung Cancer, 2019, 136, 57-64.  | 0.9 | 10        |
| 46 | Validity of using immunohistochemistry to predict treatment outcome in patients with non-small cell lung cancer not otherwise specified. Journal of Cancer Research and Clinical Oncology, 2019, 145, 2495-2506.   | 1.2 | 5         |
| 47 | Development of Immortalized Human Tumor Endothelial Cells from Renal Cancer. International Journal of Molecular Sciences, 2019, 20, 4595.  | 1.8 | 8         |
| 48 | Dexamethasone Increases Cisplatin-Loaded Nanocarrier Delivery and Efficacy in Metastatic Breast Cancer by Normalizing the Tumor Microenvironment. ACS Nano, 2019, 13, 6396-6408.   | 7.3 | 97        |
| 49 | Growth patterns of small peripheral squamous cell carcinoma of the lung and their impacts on pathological and biological characteristics of tumor cells. Journal of Cancer Research and Clinical Oncology, 2019, 145, 1773-1783.                                   | 1.2 | 8         |
| 50 | Stapling cartridge lavage cytology in limited resection for pulmonary malignant tumors: assessment of cytological status of the surgical margin. Heliyon, 2019, 5, e01240.   | 1.4 | 3         |
| 51 | Organoid culture containing cancer cells and stromal cells reveals that podoplanin-positive cancer-associated fibroblasts enhance proliferation of lung cancer cells. Lung Cancer, 2019, 134, 100-107.   | 0.9 | 40        |
| 52 | Complications and outcomes in diffuse large B-cell lymphoma with gastric lesions treated with R-CHOP. Cancer Medicine, 2019, 8, 982-989.   | 1.3 | 15        |
| 53 | Clinicopathological characteristics associated with necrosis in pulmonary metastases from colorectal cancer. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin, 2019, 474, 569-575.   | 1.4 | 6         |
| 54 | Area of residual tumor (ART) can predict prognosis after post neoadjuvant therapy resection for pancreatic ductal adenocarcinoma. Scientific Reports, 2019, 9, 17145.  | 1.6 | 15        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | Spatiotemporal characteristics of fibroblasts-dependent cancer cell invasion. <i>Journal of Cancer Research and Clinical Oncology</i> , 2019, 145, 373-381.   | 1.2 | 6         |
| 56 | Radiologic Criteria in Predicting Pathologic Less Invasive Lung Cancer According to TNM 8th Edition. <i>Clinical Lung Cancer</i> , 2019, 20, e163-e170.   | 1.1 | 14        |
| 57 | The efficacy of immune checkpoint inhibitors and PD-L1 status in patients with advanced non-small cell lung cancer harboring oncogenic driver alterations: Immuno-oncology biomarker study in LC-SCRUM-Japan.. <i>Journal of Clinical Oncology</i> , 2019, 37, 9046-9046. | 0.8 | 2         |
| 58 | Assessment of PD-L1 expression and oncogenic gene status in patients with small-cell lung cancer: Immuno-oncology biomarker study in LC-SCRUM-Japan.. <i>Journal of Clinical Oncology</i> , 2019, 37, 8558-8558.  | 0.8 | 0         |
| 59 | Podoplanin-positive cancer-associated fibroblast recruitment within cancer stroma is associated with a higher number of single nucleotide variants in cancer cells in lung adenocarcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2018, 144, 893-900.  | 1.2 | 7         |
| 60 | Immunosuppressive tumor microenvironment of usual interstitial pneumonia-associated squamous cell carcinoma of the lung. <i>Journal of Cancer Research and Clinical Oncology</i> , 2018, 144, 835-844.  | 1.2 | 7         |
| 61 | Metabolic Determinants of Sensitivity to Phosphatidylinositol 3-Kinase Pathway Inhibitor in Small-Cell Lung Carcinoma. <i>Cancer Research</i> , 2018, 78, 2179-2190.  | 0.4 | 33        |
| 62 | A secondary RET mutation in the activation loop conferring resistance to vandetanib. <i>Nature Communications</i> , 2018, 9, 625.   | 5.8 | 75        |
| 63 | Abundant tumor promoting stromal cells in lung adenocarcinoma with hypoxic regions. <i>Lung Cancer</i> , 2018, 115, 56-63.  | 0.9 | 15        |
| 64 | Utility of Site-Specific Biopsy for Diagnosis of Desmoplastic Malignant Mesothelioma. <i>Annals of Thoracic Surgery</i> , 2018, 106, e125-e128.   | 0.7 | 0         |
| 65 | The ratio of cancer cells to stroma within the invasive area is a histologic prognostic parameter of lung adenocarcinoma. <i>Lung Cancer</i> , 2018, 118, 30-35.  | 0.9 | 20        |
| 66 | Characterization of the tumor immune-microenvironment of lung adenocarcinoma associated with usual interstitial pneumonia. <i>Lung Cancer</i> , 2018, 126, 162-169.   | 0.9 | 2         |
| 67 | Link between tumor-promoting fibrous microenvironment and an immunosuppressive microenvironment in stage I lung adenocarcinoma. <i>Lung Cancer</i> , 2018, 126, 64-71.  | 0.9 | 39        |
| 68 | Genetic profiling-based prognostic prediction of patients with advanced small-cell lung cancer in large scale analysis. <i>Lung Cancer</i> , 2018, 126, 182-188.  | 0.9 | 17        |
| 69 | Podoplanin: An emerging cancer biomarker and therapeutic target. <i>Cancer Science</i> , 2018, 109, 1292-1299.  | 1.7 | 134       |
| 70 | Differences of tumor microenvironment between stage I lepidic-positive and lepidic-negative lung adenocarcinomas. <i>Journal of Thoracic and Cardiovascular Surgery</i> , 2018, 156, 1679-1688.e2.  | 0.4 | 21        |
| 71 | Collagen type I induces EGFR <sup>+</sup> TKI <sup>+</sup> resistance in EGFR <sup>+</sup> mutated cancer cells by mTOR activation through Akt <sup>+</sup> independent pathway. <i>Cancer Science</i> , 2018, 109, 2063-2073.  | 1.7 | 39        |
| 72 | Current Status and Issues of PD-L1 Testing in Non-small Cell Lung Cancer. <i>Japanese Journal of Lung Cancer</i> , 2018, 58, 189-195.   | 0.0 | 0         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 73 | Single cell time-lapse analysis reveals that podoplanin enhances cell survival and colony formation capacity of squamous cell carcinoma cells. <i>Scientific Reports</i> , 2017, 7, 39971.  | 1.6 | 18        |
| 74 | The ratio of cancer cells to stroma after induction therapy in the treatment of non-small cell lung cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 215-223.   | 1.2 | 5         |
| 75 | Changes in the tumor microenvironment during lymphatic metastasis of lung squamous cell carcinoma. <i>Cancer Science</i> , 2017, 108, 136-142.  | 1.7 | 17        |
| 76 | Fibroblast-led cancer cell invasion is activated by epithelialâ€mesenchymal transition through platelet-derived growth factor BB secretion of lung adenocarcinoma. <i>Cancer Letters</i> , 2017, 395, 20-30.  | 3.2 | 44        |
| 77 | CD200-positive cancer associated fibroblasts augment the sensitivity of Epidermal Growth Factor Receptor mutation-positive lung adenocarcinomas to EGFR Tyrosine kinase inhibitors. <i>Scientific Reports</i> , 2017, 7, 46662.   | 1.6 | 36        |
| 78 | A novel method to generate single-cell-derived cancer-associated fibroblast clones. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 1409-1419.   | 1.2 | 12        |
| 79 | Crosstalk Between Cancer Associated Fibroblasts and Cancer Cells in the Tumor Microenvironment After Radiotherapy. <i>EBioMedicine</i> , 2017, 17, 7-8.   | 2.7 | 12        |
| 80 | Clinical and Pathological Staging Validation in the Eighth Edition of the TNM Classification for Lung Cancer: Correlation between Solid Size on Thin-Section Computed Tomography and Invasive Size in Pathological Findings in the New T Classification. <i>Journal of Thoracic Oncology</i> , 2017, 12, 1403-1412. | 0.5 | 47        |
| 81 | Clinicopathological significance of caveolin-1 expression by cancer-associated fibroblasts in lung adenocarcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 321-328.  | 1.2 | 20        |
| 82 | Vandetanib in patients with previously treated RET-rearranged advanced non-small-cell lung cancer (LURET): an open-label, multicentre phase 2 trial. <i>Lancet Respiratory Medicine</i> , 2017, 5, 42-50.   | 5.2 | 252       |
| 83 | Genomic Profiling of Large-Cell Neuroendocrine Carcinoma of the Lung. <i>Clinical Cancer Research</i> , 2017, 23, 757-765.  | 3.2 | 144       |
| 84 | Clonal heterogeneity in osteogenic potential of lung cancer-associated fibroblasts: promotional effect of osteogenic progenitor cells on cancer cell migration. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016, 142, 1487-1498.   | 1.2 | 8         |
| 85 | Interstitial growth as an aggressive growth pattern in primary lung cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016, 142, 1591-1598.   | 1.2 | 7         |
| 86 | Editorial: Targeting tumor microenvironment heterogeneity. <i>Advanced Drug Delivery Reviews</i> , 2016, 99, 139.   | 6.6 | 9         |
| 87 | Factors influencing the concordance of histological subtype diagnosis from biopsy and resected specimens of lung adenocarcinoma. <i>Lung Cancer</i> , 2016, 94, 1-6.  | 0.9 | 30        |
| 88 | Aggressive tumor microenvironment of solid predominant lung adenocarcinoma subtype harboring with epidermal growth factor receptor mutations. <i>Lung Cancer</i> , 2016, 91, 7-14.  | 0.9 | 33        |
| 89 | The association of intravascular stromal cells with prognosis in high-grade neuroendocrine carcinoma of the lung. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016, 142, 905-912.   | 1.2 | 3         |
| 90 | Unique intravascular tumor microenvironment predicting recurrence of lung squamous cell carcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016, 142, 593-600.  | 1.2 | 7         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 91  | Drastic morphological and molecular differences between lymph node micrometastatic tumors and macrometastatic tumors of lung adenocarcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016, 142, 37-46.   | 1.2 | 10        |
| 92  | Cancer cell invasion driven by extracellular matrix remodeling is dependent on the properties of cancer-associated fibroblasts. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016, 142, 437-446.  | 1.2 | 33        |
| 93  | Phenotypic and functional heterogeneity of cancer-associated fibroblast within the tumor microenvironment. <i>Advanced Drug Delivery Reviews</i> , 2016, 99, 186-196.  | 6.6 | 340       |
| 94  | Podoplanin-expressing cancer-associated fibroblasts lead and enhance the local invasion of cancer cells in lung adenocarcinoma. <i>International Journal of Cancer</i> , 2015, 137, 784-796.   | 2.3 | 106       |
| 95  | Comparison of the expression levels of molecular markers among the peripheral area and central area of primary tumor and metastatic lymph node tumor in patients with squamous cell carcinoma of the lung. <i>Journal of Cancer Research and Clinical Oncology</i> , 2015, 141, 1417-1425. | 1.2 | 11        |
| 96  | Podoplanin-Positive Cancer-Associated Fibroblasts in the Tumor Microenvironment Induce Primary Resistance to EGFR-TKIs in Lung Adenocarcinoma with EGFR Mutation. <i>Clinical Cancer Research</i> , 2015, 21, 642-651.   | 3.2 | 98        |
| 97  | Therapeutic Priority of the PI3K/AKT/mTOR Pathway in Small Cell Lung Cancers as Revealed by a Comprehensive Genomic Analysis. <i>Journal of Thoracic Oncology</i> , 2014, 9, 1324-1331.  | 0.5 | 150       |
| 98  | Circulating CD14+CD204+ Cells Predict Postoperative Recurrence in Non-“Small-Cell Lung Cancer Patients. <i>Journal of Thoracic Oncology</i> , 2014, 9, 179-188.  | 0.5 | 22        |
| 99  | Distinct clinicopathologic characteristics of lung mucinous adenocarcinoma with KRAS mutation. <i>Human Pathology</i> , 2013, 44, 2636-2642.   | 1.1 | 41        |
| 100 | Podoplanin-Positive Cancer-Associated Fibroblasts Could Have Prognostic Value Independent of Cancer Cell Phenotype in Stage I Lung Squamous Cell Carcinoma. <i>Chest</i> , 2013, 143, 963-970.   | 0.4 | 60        |
| 101 | Forkhead box P3 regulatory T cells coexisting with cancer associated fibroblasts are correlated with a poor outcome in lung adenocarcinoma. <i>Cancer Science</i> , 2013, 104, 409-415.  | 1.7 | 87        |
| 102 | Prognostic Impact of Cancer-Associated Stromal Cells in Patients With Stage I Lung Adenocarcinoma. <i>Chest</i> , 2012, 142, 151-158.  | 0.4 | 106       |
| 103 | Tumor promoting effect of podoplanin-positive fibroblasts is mediated by enhanced RhoA activity. <i>Biochemical and Biophysical Research Communications</i> , 2012, 422, 194-199.  | 1.0 | 45        |
| 104 | Risk Factors for Tumor Recurrence in Patients With Early-Stage (Stage I and II) Non-small Cell Lung Cancer. <i>Chest</i> , 2011, 140, 1494-1502.   | 0.4 | 76        |
| 105 | Clinicopathological characteristics of primary lung adenocarcinoma predominantly composed of goblet cells in surgically resected cases. <i>Pathology International</i> , 2011, 61, 423-429.  | 0.6 | 17        |
| 106 | Podoplanin-Positive Fibroblasts Enhance Lung Adenocarcinoma Tumor Formation: Podoplanin in Fibroblast Functions for Tumor Progression. <i>Cancer Research</i> , 2011, 71, 4769-4779.   | 0.4 | 146       |
| 107 | A Novel Histopathological Evaluation Method Predicting the Outcome of Non-small Cell Lung Cancer Treated by Neoadjuvant Therapy: The Prognostic Importance of the Area of Residual Tumor. <i>Journal of Thoracic Oncology</i> , 2010, 5, 49-55.  | 0.5 | 47        |
| 108 | Stromal Macrophage Expressing CD204 is Associated with Tumor Aggressiveness in Lung Adenocarcinoma. <i>Journal of Thoracic Oncology</i> , 2010, 5, 1507-1515.  | 0.5 | 159       |



| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 109 | Prognostic Impact of Node Involvement Pattern in pN1 Non-small Cell Lung Cancer Patients. <i>Journal of Thoracic Oncology</i> , 2010, 5, 1576-1582.   | 0.5 | 10        |
| 110 | Prognostic impact of intratumoral vascular invasion in non-small cell lung cancer patients. <i>Thorax</i> , 2010, 65, 1092-1098.  | 2.7 | 44        |
| 111 | Fibroblasts associated with cancer cells keep enhanced migration activity after separation from cancer cells: A novel character of tumor educated fibroblasts. <i>International Journal of Oncology</i> , 2010, 37, 317-25. | 1.4 | 25        |
| 112 | Area of residual tumor beyond the muscular layer is a useful predictor of outcome in rectal cancer patients who receive preoperative chemoradiotherapy. <i>Pathology International</i> , 2009, 59, 857-862.                 | 0.6 | 11        |
| 113 | Podoplanin expression by cancer associated fibroblasts predicts poor prognosis of lung adenocarcinoma. <i>International Journal of Cancer</i> , 2008, 123, 1053-1059.   | 2.3 | 199       |
| 114 | Podoplanin, a novel marker of tumor-initiating cells in human squamous cell carcinoma A431. <i>Biochemical and Biophysical Research Communications</i> , 2008, 373, 36-41.  | 1.0 | 136       |
| 115 | Presence of Human Circulating Progenitor Cells for Cancer Stromal Fibroblasts in the Blood of Lung Cancer Patients. <i>Stem Cells</i> , 2007, 25, 1469-1477.  | 1.4 | 36        |
| 116 | Differences in clinicopathological and biological features between central-type and peripheral-type squamous cell carcinoma of the lung. <i>Lung Cancer</i> , 2006, 52, 37-45.  | 0.9 | 31        |
| 117 | Bone-marrow-derived myofibroblasts contribute to the cancer-induced stromal reaction. <i>Biochemical and Biophysical Research Communications</i> , 2003, 309, 232-240.  | 1.0 | 260       |