

Pietro Fusani

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1745970/publications.pdf>

Version: 2024-02-01

14
papers

55
citations

1937685

4
h-index

1720034

7
g-index

14
all docs

14
docs citations

14
times ranked

115
citing authors

#	ARTICLE	IF	CITATIONS
1	Phenolics and a sesquiterpene lactone in the edible shoots of <i>Cicerbita alpina</i> (L.) Wallroth. <i>Journal of Food Composition and Analysis</i> , 2010, 23, 658-663.	3.9	12
2	Cultivation trials on <i>Gentiana lutea</i> L. in Southern and South-eastern Europe. <i>Journal of Applied Research on Medicinal and Aromatic Plants</i> , 2014, 1, 113-122.	1.5	8
3	Seasonal variation in secondary metabolites of edible shoots of Buckâ€™s beard [<i>Aruncus dioicus</i> (Walter) Fernald (Rosaceae)]. <i>Food Chemistry</i> , 2016, 202, 23-30.	8.2	8
4	Harvest in different years of growth influences chemical composition of <i>Echinacea angustifolia</i> roots. <i>Industrial Crops and Products</i> , 2015, 76, 1164-1168.	5.2	7
5	The effect of cold stratification and of gibberellic acid on the seed germination of wild musk yarrow [<i>Achillea erba-rotta</i> subsp. <i>moschata</i> (Wulfen) I. Richardson] populations. <i>Journal of Applied Research on Medicinal and Aromatic Plants</i> , 2017, 7, 108-112.	1.5	6
6	Monoterpenoids from the traditional North Italian vegetable <i>Aruncus dioicus</i> (Walter) Fernald var. <i>vulgaris</i> (Maxim.) H.Hara (Rosaceae). <i>Food Chemistry</i> , 2017, 221, 1851-1859.	8.2	3
7	In vitro evaluation of the effects of methanolic plant extracts on the embryonation rate of <i>Ascaridia galli</i> eggs. <i>Veterinary Research Communications</i> , 2023, 47, 409-419.	1.6	3
8	Volatile oil features of a naturalized population of parsley [<i>Petroselinum crispum</i> (Mill) Nyman] suitable for breeding. <i>Journal of Essential Oil Research</i> , 2017, 29, 240-247.	2.7	2
9	Domestication of Alpine blue-sow-thistle (<i>Cicerbita alpina</i> (L.) Wallr.): six year trial results. <i>Genetic Resources and Crop Evolution</i> , 2012, 59, 465-471.	1.6	1
10	Seed yield and germination characteristics of wild accessions of <i>Arnica montana</i> L. from Trentino (Italy). <i>Journal of Applied Research on Medicinal and Aromatic Plants</i> , 2014, 1, e30-e33.	1.5	1
11	Effect of growth substrates on morpho-quantitative and qualitative characteristics of <i>Echinacea angustifolia</i> var. <i>angustifolia</i> roots. <i>Journal of Herbs, Spices and Medicinal Plants</i> , 2018, 24, 64-73.	1.1	1
12	Seed germination requirements of Buckâ€™s beard [<i>Aruncus dioicus</i> (Walter) Fernald (Rosaceae)]. <i>Plant Biosystems</i> , 2019, 153, 5-11.	1.6	1
13	Evaluation of the Farming Potential of <i>Echinacea Angustifolia</i> DC. Accessions Grown in Italy by Root-Marker Compound Content and Morphological Trait Analyses. <i>Plants</i> , 2020, 9, 873.	3.5	1
14	Volatile Composition Variability of <i>Arnica montana</i> Wild Populations of Trentinoâ€™Alto Adige, Italy, Determined by Headspaceâ€™Solid Phase Microextraction. <i>Chemistry and Biodiversity</i> , 2021, , e2100593.	2.1	1