

Emma Rose

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/174791/publications.pdf>

Version: 2024-02-01

28
papers

2,746
citations

331670

21
h-index

526287

27
g-index

28
all docs

28
docs citations

28
times ranked

5814
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | The moderating role of socioeconomic status on level of responsibility, executive functioning, and cortical thinning during adolescence. <i>Developmental Psychobiology</i> , 2021, 63, 291-304. | 1.6 | 0 |
| 2 | Temporal Difference Error Prediction Signal Dysregulation in Cocaine Dependence. <i>Neuropsychopharmacology</i> , 2014, 39, 1732-1742. | 5.4 | 25 |
| 3 | The miRâ€137 schizophrenia susceptibility variant rs1625579 does not predict variability in brain volume in a sample of schizophrenic patients and healthy individuals. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2014, 165, 467-471. | 1.7 | 17 |
| 4 | Effects of MIR137 on fronto-amygdala functional connectivity. <i>NeuroImage</i> , 2014, 90, 189-195. | 4.2 | 42 |
| 5 | The ENIGMA Consortium: large-scale collaborative analyses of neuroimaging and genetic data. <i>Brain Imaging and Behavior</i> , 2014, 8, 153-182. | 2.1 | 696 |
| 6 | Effects of a novel schizophrenia risk variant rs7914558 at <i>CNNM2</i> on brain structure and attributional style. <i>British Journal of Psychiatry</i> , 2014, 204, 115-121. | 2.8 | 30 |
| 7 | Genome-wide schizophrenia variant at MIR137 does not impact white matter microstructure in healthy participants. <i>Neuroscience Letters</i> , 2014, 574, 6-10. | 2.1 | 15 |
| 8 | Altered medial prefrontal activity during dynamic face processing in schizophrenia spectrum patients. <i>Schizophrenia Research</i> , 2014, 157, 225-230. | 2.0 | 30 |
| 9 | Neuropsychological effects of the <i>CSMD1</i> genome-wide associated schizophrenia risk variant rs10503253. <i>Genes, Brain and Behavior</i> , 2013, 12, 203-209. | 2.2 | 48 |
| 10 | Acute Nicotine Differentially Impacts Anticipatory Valence- and Magnitude-Related Striatal Activity. <i>Biological Psychiatry</i> , 2013, 73, 280-288. | 1.3 | 55 |
| 11 | Brain vs Behavior: An Effect Size Comparison of Neuroimaging and Cognitive Studies of Genetic Risk for Schizophrenia. <i>Schizophrenia Bulletin</i> , 2013, 39, 518-526. | 4.3 | 83 |
| 12 | Neural effects of the <i>CSMD1</i> genome-wide associated schizophrenia risk variant rs10503253. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2013, 162, 530-537. | 1.7 | 30 |
| 13 | The Effect of the Neurogranin Schizophrenia Risk Variant rs12807809 on Brain Structure and Function. <i>Twin Research and Human Genetics</i> , 2012, 15, 296-303. | 0.6 | 26 |
| 14 | ZNF804A and social cognition in patients with schizophrenia and healthy controls. <i>Molecular Psychiatry</i> , 2012, 17, 118-119. | 7.9 | 20 |
| 15 | Identification of common variants associated with human hippocampal and intracranial volumes. <i>Nature Genetics</i> , 2012, 44, 552-561. | 21.4 | 594 |
| 16 | Social cognition in bipolar disorder versus schizophrenia: comparability in mental state decoding deficits. <i>Bipolar Disorders</i> , 2012, 14, 743-748. | 1.9 | 44 |
| 17 | Chronic Exposure to Nicotine Is Associated with Reduced Reward-Related Activity in the Striatum but not the Midbrain. <i>Biological Psychiatry</i> , 2012, 71, 206-213. | 1.3 | 59 |
| 18 | The NOS1 variant rs6490121 is associated with variation in prefrontal function and grey matter density in healthy individuals. <i>NeuroImage</i> , 2012, 60, 614-622. | 4.2 | 26 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | A neuropsychological investigation of the genome wide associated schizophrenia risk variant NRCN rs12807809. Schizophrenia Research, 2011, 125, 304-306. | 2.0 | 23 |
| 20 | ZNF804A risk allele is associated with relatively intact gray matter volume in patients with schizophrenia. NeuroImage, 2011, 54, 2132-2137. | 4.2 | 78 |
| 21 | Brain activation predicts treatment improvement in patients with major depressive disorder. Journal of Psychiatric Research, 2011, 45, 1214-22. | 3.1 | 53 |
| 22 | Nicotine modulation of information processing is not limited to input (attention) but extends to output (intention). Psychopharmacology, 2010, 209, 291-302. | 3.1 | 16 |
| 23 | Abnormal Responses to Monetary Outcomes in Cortex, but not in the Basal Ganglia, in Schizophrenia. Neuropsychopharmacology, 2010, 35, 2427-2439. | 5.4 | 137 |
| 24 | Patients with Schizophrenia have a Reduced Neural Response to Both Unpredictable and Predictable Primary Reinforcers. Neuropsychopharmacology, 2009, 34, 1567-1577. | 5.4 | 166 |
| 25 | Limbic over-activity in depression during preserved performance on the n-back task. NeuroImage, 2006, 29, 203-215. | 4.2 | 111 |
| 26 | The effects of escitalopram on working memory and brain activity in healthy adults during performance of the n-back task. Psychopharmacology, 2006, 185, 339-347. | 3.1 | 47 |
| 27 | Cognitive impairment and fMRI in major depression. Neurotoxicity Research, 2006, 10, 87-92. | 2.7 | 35 |
| 28 | Pattern of impaired working memory during major depression. Journal of Affective Disorders, 2006, 90, 149-161. | 4.1 | 240 |