

Ji Sun Nam

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1751585/publications.pdf>

Version: 2024-02-01

59
papers

1,520
citations

331259

21
h-index

329751

37
g-index

59
all docs

59
docs citations

59
times ranked

2349
citing authors

#	ARTICLE	IF	CITATIONS
1	Elevated TyG Index Predicts Progression of Coronary Artery Calcification. <i>Diabetes Care</i> , 2019, 42, 1569-1573.	4.3	180
2	Association between triglyceride glucose index and arterial stiffness in Korean adults. <i>Cardiovascular Diabetology</i> , 2018, 17, 41.	2.7	169
3	Relationship between the triglyceride glucose index and coronary artery calcification in Korean adults. <i>Cardiovascular Diabetology</i> , 2017, 16, 108.	2.7	140
4	Triglyceride Glucose Index Is Superior to the Homeostasis Model Assessment of Insulin Resistance for Predicting Nonalcoholic Fatty Liver Disease in Korean Adults. <i>Endocrinology and Metabolism</i> , 2019, 34, 179.	1.3	82
5	The effects of aerobic exercise training on serum osteocalcin, adipocytokines and insulin resistance on obese young males. <i>Clinical Endocrinology</i> , 2015, 82, 686-694.	1.2	60
6	Relationship between natural killer cell activity and glucose control in patients with type 2 diabetes and prediabetes. <i>Journal of Diabetes Investigation</i> , 2019, 10, 1223-1228.	1.1	56
7	Usefulness of brachial-ankle pulse wave velocity as a predictive marker of multiple coronary artery occlusive disease in Korean type 2 diabetes patients. <i>Diabetes Research and Clinical Practice</i> , 2009, 85, 30-34.	1.1	45
8	Insulin resistance independently influences arterial stiffness in normoglycemic normotensive postmenopausal women. <i>Menopause</i> , 2010, 17, 779-784.	0.8	44
9	Predictive Clinical Parameters for the Therapeutic Efficacy of Sitagliptin in Korean Type 2 Diabetes Mellitus. <i>Diabetes and Metabolism Journal</i> , 2011, 35, 159.	1.8	43
10	The activation of NF- κ B and AP-1 in peripheral blood mononuclear cells isolated from patients with diabetic nephropathy. <i>Diabetes Research and Clinical Practice</i> , 2008, 81, 25-32.	1.1	41
11	High circulating microRNA-122 expression is a poor prognostic marker in patients with hepatitis B virus-related hepatocellular carcinoma who undergo radiofrequency ablation. <i>Clinical Biochemistry</i> , 2015, 48, 1073-1078.	0.8	40
12	Effect of pioglitazone on serum concentrations of osteoprotegerin in patients with type 2 diabetes mellitus. <i>European Journal of Endocrinology</i> , 2011, 164, 69-74.	1.9	39
13	Some Elements in Thyroid Tissue are Associated with More Advanced Stage of Thyroid Cancer in Korean Women. <i>Biological Trace Element Research</i> , 2016, 171, 54-62.	1.9	39
14	Low levels of circulating microRNA-26a/29a as poor prognostic markers in patients with hepatocellular carcinoma who underwent curative treatment. <i>Clinics and Research in Hepatology and Gastroenterology</i> , 2017, 41, 181-189.	0.7	37
15	Association between nonalcoholic fatty liver disease and coronary artery calcification in postmenopausal women. <i>Menopause</i> , 2015, 22, 1323-1327.	0.8	33
16	Visceral adiposity and leptin are independently associated with C-reactive protein in Korean type 2 diabetic patients. <i>Acta Diabetologica</i> , 2010, 47, 113-118.	1.2	30
17	Insulin resistance is independently associated with peripheral and autonomic neuropathy in Korean type 2 diabetic patients. <i>Acta Diabetologica</i> , 2012, 49, 97-103.	1.2	29
18	Transplantation of insulin-secreting cells differentiated from human adipose tissue-derived stem cells into type 2 diabetes mice. <i>Biochemical and Biophysical Research Communications</i> , 2014, 443, 775-781.	1.0	25

#	ARTICLE	IF	CITATIONS
19	Obesity and Hyperglycemia in Korean Men with Klinefelter Syndrome: The Korean Endocrine Society Registry. <i>Endocrinology and Metabolism</i> , 2016, 31, 598.	1.3	25
20	Plasma miRNA-122 as a predictive marker for treatment response following transarterial chemoembolization in patients with hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2017, 32, 199-207.	1.4	23
21	Fetal bovine serum-free cryopreservation methods for clinical banking of human adipose-derived stem cells. <i>Cryobiology</i> , 2018, 81, 65-73.	0.3	22
22	Cerebral arterial pulsatility and insulin resistance in type 2 diabetic patients. <i>Diabetes Research and Clinical Practice</i> , 2008, 79, 237-242.	1.1	20
23	Anti-diabetic properties of different fractions of Korean red ginseng. <i>Journal of Ethnopharmacology</i> , 2019, 236, 220-230.	2.0	20
24	Association between atherogenic index of plasma and coronary artery calcification progression in Korean adults. <i>Lipids in Health and Disease</i> , 2020, 19, 157.	1.2	19
25	Relationship of low-density lipoprotein (LDL) particle size to thyroid function status in Koreans. <i>Clinical Endocrinology</i> , 2009, 71, 130-136.	1.2	18
26	Hemorheological Approach for Early Detection of Chronic Kidney Disease and Diabetic Nephropathy in Type 2 Diabetes. <i>Diabetes Technology and Therapeutics</i> , 2015, 17, 808-815.	2.4	18
27	Plasma MicroRNA-21, 26a, and 29a-3p as Predictive Markers for Treatment Response Following Transarterial Chemoembolization in Patients with Hepatocellular Carcinoma. <i>Journal of Korean Medical Science</i> , 2018, 33, e6.	1.1	18
28	The effect of Korean Red Ginseng on sarcopenia biomarkers in type 2 diabetes patients. <i>Archives of Gerontology and Geriatrics</i> , 2020, 90, 104108.	1.4	18
29	Association between lipoprotein(a) and nonalcoholic fatty liver disease among Korean adults. <i>Clinica Chimica Acta</i> , 2016, 461, 14-18.	0.5	17
30	Higher serum interleukin-17A levels as a potential biomarker for predicting early disease progression in patients with hepatitis B virus-associated advanced hepatocellular carcinoma treated with sorafenib. <i>Cytokine</i> , 2017, 95, 118-125.	1.4	17
31	Triiodothyronine level predicts visceral obesity and atherosclerosis in euthyroid, overweight and obese subjects: T3 and visceral obesity. <i>Obesity Research and Clinical Practice</i> , 2010, 4, e315-e323.	0.8	14
32	Serum osteoprotegerin is associated with vascular stiffness and the onset of new cardiovascular events in hemodialysis patients. <i>Korean Journal of Internal Medicine</i> , 2013, 28, 668.	0.7	14
33	Use of RBC deformability index as an early marker of diabetic nephropathy. <i>Clinical Hemorheology and Microcirculation</i> , 2019, 72, 75-84.	0.9	13
34	The Plasma Atherogenic Index is an Independent Predictor of Arterial Stiffness in Healthy Koreans. <i>Angiology</i> , 2022, 73, 514-519.	0.8	13
35	Circulating myokine levels in different stages of glucose intolerance. <i>Medicine (United States)</i> , 2020, 99, e19235.	0.4	12
36	Supplementation with Korean Red Ginseng Improves Current Perception Threshold in Korean Type 2 Diabetes Patients: A Randomized, Double-Blind, Placebo-Controlled Trial. <i>Journal of Diabetes Research</i> , 2020, 2020, 1-8.	1.0	11

#	ARTICLE	IF	CITATIONS
37	The increased amount of coffee consumption lowers the incidence of fatty liver disease in Korean men. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020, 30, 1653-1661.	1.1	10
38	Semaphorin 3A is a Novel Adipokine Representing Exercise-Induced Improvements of Metabolism in Metabolically Healthy Obese Young Males. <i>Scientific Reports</i> , 2020, 10, 10005.	1.6	10
39	Adiponectin is independently associated with apolipoprotein B to A-1 ratio in Koreans. <i>Metabolism: Clinical and Experimental</i> , 2010, 59, 677-682.	1.5	9
40	The effects of pioglitazone on cerebrovascular resistance in patients with type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2007, 56, 1081-1086.	1.5	8
41	Serum ferritin levels are associated with arterial stiffness in healthy Korean adults. <i>Vascular Medicine</i> , 2016, 21, 325-330.	0.8	6
42	Calpain-10 and Adiponectin Gene Polymorphisms in Korean Type 2 Diabetes Patients. <i>Endocrinology and Metabolism</i> , 2018, 33, 364.	1.3	6
43	Association between <i>BDNF</i> Polymorphism and Depressive Symptoms in Patients Newly Diagnosed with Type 2 Diabetes Mellitus. <i>Yonsei Medical Journal</i> , 2021, 62, 359.	0.9	6
44	Antioxidant-Rich Dietary Intervention Improves Cardiometabolic Profiles and Arterial Stiffness in Elderly Koreans with Metabolic Syndrome. <i>Yonsei Medical Journal</i> , 2022, 63, 26.	0.9	6
45	Visceral Fat Thickness Predicts Fatty Liver in Koreans with Type 2 Diabetes Mellitus. <i>Journal of Korean Medical Science</i> , 2008, 23, 256.	1.1	5
46	Usefulness of Multidetector Row Computed Tomography for Predicting Cardiac Events in Asymptomatic Chronic Kidney Disease Patients at the Initiation of Renal Replacement Therapy. <i>Scientific World Journal</i> , The, 2013, 2013, 1-6.	0.8	5
47	No effect of vitamin D supplementation on metabolic parameters but on lipids in patients with type 2 diabetes and chronic kidney disease. <i>International Journal for Vitamin and Nutrition Research</i> , 2021, 91, 649-658.	0.6	2
48	Relation between Cerebral Arterial Pulsatility and Insulin Resistance in Type 2 Diabetic Patients. <i>The Journal of Korean Diabetes Association</i> , 2006, 30, 347.	0.1	1
49	Non-Laboratory-Based Simple Screening Model for Nonalcoholic Fatty Liver Disease in Patients with Type 2 Diabetes Developed Using Multi-Center Cohorts. <i>Endocrinology and Metabolism</i> , 2021, 36, 823-834.	1.3	1
50	Oxidative stress, point-of-care test, and metabolic syndrome. <i>Korean Journal of Internal Medicine</i> , 2014, 29, 20.	0.7	1
51	A Case of Acromegaly with Gall Bladder Cancer. <i>Journal of Korean Endocrine Society</i> , 2005, 20, 401.	0.1	0
52	A Case of Turner's Syndrome with Transient Hypopituitarism. <i>Journal of Korean Endocrine Society</i> , 2007, 22, 266.	0.1	0
53	In vivo Corneal Confocal Microscopy and Nerve Growth Factor in Diabetic Microvascular Complications. <i>The Journal of Korean Diabetes Association</i> , 2007, 31, 351.	0.1	0
54	Change in waist circumference and the progression of subclinical atherosclerosis in type 2 diabetes patients. <i>Obesity Research and Clinical Practice</i> , 2011, 5, e202-e209.	0.8	0

#	ARTICLE	IF	CITATIONS
55	The relationship between cholesterol concentration and carotid intima media thickness differs according to gender and menopausal status in Korean type 2 diabetic patients. Clinica Chimica Acta, 2016, 455, 107-112.	0.5	0
56	Reversible Pituitary Dysfunction in a Patient with Cushing's Syndrome due to Adrenal Adenoma. Journal of Korean Endocrine Society, 2006, 21, 146.	0.1	0
57	A Case of Acromegaly Caused by Double Pituitary Adenomas. Journal of Korean Endocrine Society, 2006, 21, 53.	0.1	0
58	SUN-072 Elevated TyG Index Predicts Progression of Coronary Artery Calcification. Journal of the Endocrine Society, 2019, 3, .	0.1	0
59	The Effects of C. lacerata on Insulin Resistance in Type 2 Diabetes Patients. Journal of Diabetes Research, 2022, 2022, 1-10.	1.0	0