Ji Sun Nam

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1751585/publications.pdf

Version: 2024-02-01

	331259	329751
1,520	21	37
citations	h-index	g-index
50		00.40
59	59	2349
docs citations	times ranked	citing authors
	1,520 citations 59 docs citations	1,520 21 citations h-index 59 59

#	Article	IF	CITATIONS
1	The Plasma Atherogenic Index is an Independent Predictor of Arterial Stiffness in Healthy Koreans. Angiology, 2022, 73, 514-519.	0.8	13
2	Antioxidant-Rich Dietary Intervention Improves Cardiometabolic Profiles and Arterial Stiffness in Elderly Koreans with Metabolic Syndrome. Yonsei Medical Journal, 2022, 63, 26.	0.9	6
3	The Effects of C. lacerata on Insulin Resistance in Type 2 Diabetes Patients. Journal of Diabetes Research, 2022, 2022, 1-10.	1.0	O
4	Association between <i>BDNF</i> Polymorphism and Depressive Symptoms in Patients Newly Diagnosed with Type 2 Diabetes Mellitus. Yonsei Medical Journal, 2021, 62, 359.	0.9	6
5	Non-Laboratory-Based Simple Screening Model for Nonalcoholic Fatty Liver Disease in Patients with Type 2 Diabetes Developed Using Multi-Center Cohorts. Endocrinology and Metabolism, 2021, 36, 823-834.	1.3	1
6	No effect of vitamin D supplementation on metabolic parameters but on lipids in patients with type 2 diabetes and chronic kidney disease. International Journal for Vitamin and Nutrition Research, 2021, 91, 649-658.	0.6	2
7	The effect of Korean Red Ginseng on sarcopenia biomarkers in type 2 diabetes patients. Archives of Gerontology and Geriatrics, 2020, 90, 104108.	1.4	18
8	The increased amount of coffee consumption lowers the incidence of fatty liver disease in Korean men. Nutrition, Metabolism and Cardiovascular Diseases, 2020, 30, 1653-1661.	1.1	10
9	Semaphorin 3 C is a Novel Adipokine Representing Exercise-Induced Improvements of Metabolism in Metabolically Healthy Obese Young Males. Scientific Reports, 2020, 10, 10005.	1.6	10
10	Association between atherogenic index of plasma and coronary artery calcification progression in Korean adults. Lipids in Health and Disease, 2020, 19, 157.	1.2	19
11	Circulating myokine levels in different stages of glucose intolerance. Medicine (United States), 2020, 99, e19235.	0.4	12
12	Supplementation with Korean Red Ginseng Improves Current Perception Threshold in Korean Type 2 Diabetes Patients: A Randomized, Double-Blind, Placebo-Controlled Trial. Journal of Diabetes Research, 2020, 2020, 1-8.	1.0	11
13	Triglyceride Glucose Index Is Superior to the Homeostasis Model Assessment of Insulin Resistance for Predicting Nonalcoholic Fatty Liver Disease in Korean Adults. Endocrinology and Metabolism, 2019, 34, 179.	1.3	82
14	Elevated TyG Index Predicts Progression of Coronary Artery Calcification. Diabetes Care, 2019, 42, 1569-1573.	4.3	180
15	Anti-diabetic properties of different fractions of Korean red ginseng. Journal of Ethnopharmacology, 2019, 236, 220-230.	2.0	20
16	Use of RBC deformability index as an early marker of diabetic nephropathy. Clinical Hemorheology and Microcirculation, 2019, 72, 75-84.	0.9	13
17	Relationship between natural killer cell activity and glucose control in patients with typeÂ2 diabetes and prediabetes. Journal of Diabetes Investigation, 2019, 10, 1223-1228.	1.1	56
18	SUN-072 Elevated TyG Index Predicts Progression of Coronary Artery Calcification. Journal of the Endocrine Society, 2019, 3, .	0.1	0

#	Article	IF	Citations
19	Fetal bovine serum-free cryopreservation methods for clinical banking of human adipose-derived stem cells. Cryobiology, 2018, 81, 65-73.	0.3	22
20	Association between triglyceride glucose index and arterial stiffness in Korean adults. Cardiovascular Diabetology, 2018, 17, 41.	2.7	169
21	Calpain-10 and Adiponectin Gene Polymorphisms in Korean Type 2 Diabetes Patients. Endocrinology and Metabolism, 2018, 33, 364.	1.3	6
22	Plasma MicroRNA-21, 26a, and 29a-3p as Predictive Markers for Treatment Response Following Transarterial Chemoembolization in Patients with Hepatocellular Carcinoma. Journal of Korean Medical Science, 2018, 33, e6.	1.1	18
23	Plasma micoRNAâ€122 as a predictive marker for treatment response following transarterial chemoembolization in patients with hepatocellular carcinoma. Journal of Gastroenterology and Hepatology (Australia), 2017, 32, 199-207.	1.4	23
24	Higher serum interleukin-17A levels as a potential biomarker for predicting early disease progression in patients with hepatitis B virus-associated advanced hepatocellular carcinoma treated with sorafenib. Cytokine, 2017, 95, 118-125.	1.4	17
25	Low levels of circulating microRNA-26a/29a as poor prognostic markers in patients with hepatocellular carcinoma who underwent curative treatment. Clinics and Research in Hepatology and Gastroenterology, 2017, 41, 181-189.	0.7	37
26	Relationship between the triglyceride glucose index and coronary artery calcification in Korean adults. Cardiovascular Diabetology, 2017, 16, 108.	2.7	140
27	Obesity and Hyperglycemia in Korean Men with Klinefelter Syndrome: The Korean Endocrine Society Registry. Endocrinology and Metabolism, 2016, 31, 598.	1.3	25
28	Association between lipoprotein(a) and nonalcoholic fatty liver disease among Korean adults. Clinica Chimica Acta, 2016, 461, 14-18.	0.5	17
29	The relationship between cholesterol concentration and carotid intima media thickness differs according to gender and menopausal status in Korean type 2 diabetic patients. Clinica Chimica Acta, 2016, 455, 107-112.	0.5	0
30	Serum ferritin levels are associated with arterial stiffness in healthy Korean adults. Vascular Medicine, 2016, 21, 325-330.	0.8	6
31	Some Elements in Thyroid Tissue are Associated with More Advanced Stage of Thyroid Cancer in Korean Women. Biological Trace Element Research, 2016, 171, 54-62.	1.9	39
32	Association between nonalcoholic fatty liver disease and coronary artery calcification in postmenopausal women. Menopause, 2015, 22, 1323-1327.	0.8	33
33	High circulating microRNA-122 expression is a poor prognostic marker in patients with hepatitis B virus-related hepatocellular carcinoma who undergo radiofrequency ablation. Clinical Biochemistry, 2015, 48, 1073-1078.	0.8	40
34	Hemorheological Approach for Early Detection of Chronic Kidney Disease and Diabetic Nephropathy in Type 2 Diabetes. Diabetes Technology and Therapeutics, 2015, 17, 808-815.	2.4	18
35	The effects of aerobic exercise training on serum osteocalcin, adipocytokines and insulin resistance on obese young males. Clinical Endocrinology, 2015, 82, 686-694.	1.2	60
36	Transplantation of insulin-secreting cells differentiated from human adipose tissue-derived stem cells into type 2 diabetes mice. Biochemical and Biophysical Research Communications, 2014, 443, 775-781.	1.0	25

#	Article	IF	CITATIONS
37	Oxidative stress, point-of-care test, and metabolic syndrome. Korean Journal of Internal Medicine, 2014, 29, 20.	0.7	1
38	Usefulness of Multidetector Row Computed Tomography for Predicting Cardiac Events in Asymptomatic Chronic Kidney Disease Patients at the Initiation of Renal Replacement Therapy. Scientific World Journal, The, 2013, 2013, 1-6.	0.8	5
39	Serum osteoprotegerin is associated with vascular stiffness and the onset of new cardiovascular events in hemodialysis patients. Korean Journal of Internal Medicine, 2013, 28, 668.	0.7	14
40	Insulin resistance is independently associated with peripheral and autonomic neuropathy in Korean type 2 diabetic patients. Acta Diabetologica, 2012, 49, 97-103.	1.2	29
41	Change in waist circumference and the progression of subclinical atherosclerosis in type 2 diabetes patients. Obesity Research and Clinical Practice, 2011, 5, e202-e209.	0.8	0
42	Predictive Clinical Parameters for the Therapeutic Efficacy of Sitagliptin in Korean Type 2 Diabetes Mellitus. Diabetes and Metabolism Journal, 2011, 35, 159.	1.8	43
43	Effect of pioglitazone on serum concentrations of osteoprotegerin in patients with type 2 diabetes mellitus. European Journal of Endocrinology, 2011, 164, 69-74.	1.9	39
44	Visceral adiposity and leptin are independently associated with C-reactive protein in Korean type 2 diabetic patients. Acta Diabetologica, 2010, 47, 113-118.	1.2	30
45	Adiponectin is independently associated with apolipoprotein B to A-1 ratio in Koreans. Metabolism: Clinical and Experimental, 2010, 59, 677-682.	1.5	9
46	Triiodothyronine level predicts visceral obesity and atherosclerosis in euthyroid, overweight and obese subjects: T3 and visceral obesity. Obesity Research and Clinical Practice, 2010, 4, e315-e323.	0.8	14
47	Insulin resistance independently influences arterial stiffness in normoglycemic normotensive postmenopausal women. Menopause, 2010, 17, 779-784.	0.8	44
48	Relationship of lowâ€density lipoprotein (LDL) particle size to thyroid function status in Koreans. Clinical Endocrinology, 2009, 71, 130-136.	1.2	18
49	Usefulness of brachial-ankle pulse wave velocity as a predictive marker of multiple coronary artery occlusive disease in Korean type 2 diabetes patients. Diabetes Research and Clinical Practice, 2009, 85, 30-34.	1.1	45
50	Cerebral arterial pulsatility and insulin resistance in type 2 diabetic patients. Diabetes Research and Clinical Practice, 2008, 79, 237-242.	1.1	20
51	The activation of NF- $\hat{\mathbb{P}}$ B and AP-1 in peripheral blood mononuclear cells isolated from patients with diabetic nephropathy. Diabetes Research and Clinical Practice, 2008, 81, 25-32.	1.1	41
52	Visceral Fat Thickness Predicts Fatty Liver in Koreans with Type 2 Diabetes Mellitus. Journal of Korean Medical Science, 2008, 23, 256.	1.1	5
53	The effects of pioglitazone on cerebrovascular resistance in patients with type 2 diabetes mellitus. Metabolism: Clinical and Experimental, 2007, 56, 1081-1086.	1.5	8
54	A Case of Turner's Syndrome with Transient Hypopituitarism. Journal of Korean Endocrine Society, 2007, 22, 266.	0.1	0

#	Article	IF	CITATIONS
55	In vivo Corneal Confocal Microscopy and Nerve Growth Factor in Diabetic Microvascular Complications. The Journal of Korean Diabetes Association, 2007, 31, 351.	0.1	О
56	Relation between Cerebral Arterial Pulsatility and Insulin Resistance in Type 2 Diabetic Patients. The Journal of Korean Diabetes Association, 2006, 30, 347.	0.1	1
57	Reversible Pituitary Dysfunction in a Patient with Cushing's Syndrome due to Adrenal Adenoma. Journal of Korean Endocrine Society, 2006, 21, 146.	0.1	O
58	A Case of Acromegaly Caused by Double Pituitary Adenomas. Journal of Korean Endocrine Society, 2006, 21, 53.	0.1	0
59	A Case of Acromegaly with Gall Bladder Cancer. Journal of Korean Endocrine Society, 2005, 20, 401.	0.1	0