

# Daniel Armando Morris

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/1751699/daniel-armando-morris-publications-by-citations.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

46  
papers

1,668  
citations

21  
h-index

40  
g-index

51  
ext. papers

2,435  
ext. citations

4.3  
avg, IF

4.22  
L-index

| #  | Paper  | IF   | Citations |
|----|--|------|-----------|
| 46 | How to diagnose heart failure with preserved ejection fraction: the HFA-PEFF diagnostic algorithm: a consensus recommendation from the Heart Failure Association (HFA) of the European Society of Cardiology (ESC). <i>European Heart Journal</i> , <b>2019</b> , 40, 3297-3317                  | 9.5  | 379       |
| 45 | Normal values and clinical relevance of left atrial myocardial function analysed by speckle-tracking echocardiography: multicentre study. <i>European Heart Journal Cardiovascular Imaging</i> , <b>2015</b> , 16, 364-72  | 4.1  | 116       |
| 44 | Sarcopenia in patients with heart failure with preserved ejection fraction: Impact on muscle strength, exercise capacity and quality of life. <i>International Journal of Cardiology</i> , <b>2016</b> , 222, 41-46  | 3.2  | 109       |
| 43 | Potential Usefulness and Clinical Relevance of Adding Left Atrial Strain to Left Atrial Volume Index in the Detection of Left Ventricular Diastolic Dysfunction. <i>JACC: Cardiovascular Imaging</i> , <b>2018</b> , 11, 1405-1415   | 8.4  | 103       |
| 42 | Left atrial systolic and diastolic dysfunction in heart failure with normal left ventricular ejection fraction. <i>Journal of the American Society of Echocardiography</i> , <b>2011</b> , 24, 651-62  | 5.8  | 101       |
| 41 | How to diagnose heart failure with preserved ejection fraction: the HFA-PEFF diagnostic algorithm: a consensus recommendation from the Heart Failure Association (HFA) of the European Society of Cardiology (ESC). <i>European Journal of Heart Failure</i> , <b>2020</b> , 22, 391-412         | 12.3 | 84        |
| 40 | Normal range and usefulness of right ventricular systolic strain to detect subtle right ventricular systolic abnormalities in patients with heart failure: a multicentre study. <i>European Heart Journal Cardiovascular Imaging</i> , <b>2017</b> , 18, 212-223                                 | 4.1  | 70        |
| 39 | Right ventricular myocardial systolic and diastolic dysfunction in heart failure with normal left ventricular ejection fraction. <i>Journal of the American Society of Echocardiography</i> , <b>2011</b> , 24, 886-97   | 5.8  | 65        |
| 38 | Genetic profiling of tumours using both circulating free DNA and circulating tumour cells isolated from the same preserved whole blood sample. <i>Molecular Oncology</i> , <b>2016</b> , 10, 566-74  | 7.9  | 59        |
| 37 | Myocardial systolic and diastolic performance derived by 2-dimensional speckle tracking echocardiography in heart failure with normal left ventricular ejection fraction. <i>Circulation: Heart Failure</i> , <b>2012</b> , 5, 610-20  | 7.6  | 56        |
| 36 | Left ventricular longitudinal systolic function analysed by 2D speckle-tracking echocardiography in heart failure with preserved ejection fraction: a meta-analysis. <i>Open Heart</i> , <b>2017</b> , 4, e000630  | 3    | 42        |
| 35 | Iron deficiency in patients with heart failure with preserved ejection fraction and its association with reduced exercise capacity, muscle strength and quality of life. <i>Clinical Research in Cardiology</i> , <b>2019</b> , 108, 203-211   | 6.1  | 36        |
| 34 | Clinical significance of the assessment of the systolic and diastolic myocardial function of the left atrium in patients with paroxysmal atrial fibrillation and low CHADS(2) index treated with catheter ablation therapy. <i>American Journal of Cardiology</i> , <b>2013</b> , 111, 1002-11   | 3    | 36        |
| 33 | Global cardiac alterations detected by speckle-tracking echocardiography in Fabry disease: left ventricular, right ventricular, and left atrial dysfunction are common and linked to worse symptomatic status. <i>International Journal of Cardiovascular Imaging</i> , <b>2015</b> , 31, 301-13 | 2.5  | 35        |
| 32 | Myocardial systolic and diastolic consequences of left ventricular mechanical dyssynchrony in heart failure with normal left ventricular ejection fraction. <i>European Heart Journal Cardiovascular Imaging</i> , <b>2012</b> , 13, 556-67  | 4.1  | 30        |
| 31 | Multimodality imaging approach in the diagnosis of chronic myocarditis with preserved left ventricular ejection fraction (MCpEF): The role of 2D speckle-tracking echocardiography. <i>International Journal of Cardiology</i> , <b>2017</b> , 243, 374-378                                      | 3.2  | 27        |
| 30 | Exercise training in Diastolic Heart Failure (Ex-DHF): rationale and design of a multicentre, prospective, randomized, controlled, parallel group trial. <i>European Journal of Heart Failure</i> , <b>2017</b> , 19, 1067-1074  | 12.3 | 25        |

|    |   |     |    |
|----|---|-----|----|
| 29 | Right ventricular strain in heart failure: Clinical perspective. <i>Archives of Cardiovascular Diseases</i> , <b>2017</b> , 110, 562-571  | 2.7 | 25 |
| 28 | Normal Global Longitudinal Strain: An Individual Patient Meta-Analysis. <i>JACC: Cardiovascular Imaging</i> , <b>2020</b> , 13, 167-169   | 8.4 | 22 |
| 27 | Clinical Relevance of Left Atrial Strain to Predict Recurrence of Atrial Fibrillation after Catheter Ablation: A Meta-Analysis. <i>Echocardiography</i> , <b>2016</b> , 33, 724-33  | 1.5 | 21 |
| 26 | Left atrial strain predicts recurrence of atrial arrhythmias after catheter ablation of persistent atrial fibrillation. <i>Open Heart</i> , <b>2017</b> , 4, e000572  | 3   | 19 |
| 25 | Diastolic stress test echocardiography in patients with suspected heart failure with preserved ejection fraction: a pilot study. <i>ESC Heart Failure</i> , <b>2019</b> , 6, 146-153  | 3.7 | 19 |
| 24 | Multimodality imaging in patients with heart failure and preserved ejection fraction: an expert consensus document of the European Association of Cardiovascular Imaging. <i>European Heart Journal Cardiovascular Imaging</i> , <b>2021</b> ,                                  | 4.1 | 19 |
| 23 | Left atrial strain as sensitive marker of left ventricular diastolic dysfunction in heart failure. <i>ESC Heart Failure</i> , <b>2020</b> , 7, 1956-1965  | 3.7 | 17 |
| 22 | Left ventricular strain and twisting in heart failure with preserved ejection fraction: an updated review. <i>Heart Failure Reviews</i> , <b>2017</b> , 22, 371-379   | 5   | 16 |
| 21 | Lower limit of normality and clinical relevance of left ventricular early diastolic strain rate for the detection of left ventricular diastolic dysfunction. <i>European Heart Journal Cardiovascular Imaging</i> , <b>2018</b> , 19, 905-915                                   | 4.1 | 13 |
| 20 | Structural and functional cardiac analyses using modern and sensitive myocardial techniques in adult Pompe disease. <i>International Journal of Cardiovascular Imaging</i> , <b>2015</b> , 31, 947-56   | 2.5 | 12 |
| 19 | Early detection of cardiac alterations by left atrial strain in patients with risk for cardiac abnormalities with preserved left ventricular systolic and diastolic function. <i>International Journal of Cardiovascular Imaging</i> , <b>2018</b> , 34, 701-711                | 2.5 | 12 |
| 18 | Cardiac anomaly detection based on time and frequency domain features using tree-based classifiers. <i>Physiological Measurement</i> , <b>2018</b> , 39, 114001   | 2.9 | 12 |
| 17 | Subclinical left atrial dysfunction profiles for prediction of cardiac outcome in the general population. <i>Journal of Hypertension</i> , <b>2020</b> , 38, 2465-2474  | 1.9 | 11 |
| 16 | Phasic Left Atrial Function in Cancer Patients Before Initiation of Anti-Cancer Therapy. <i>Journal of Clinical Medicine</i> , <b>2019</b> , 8,   | 5.1 | 7  |
| 15 | The effect of iron deficiency on cardiac resynchronization therapy: results from the RIDE-CRT Study. <i>ESC Heart Failure</i> , <b>2020</b> , 7, 1072-1084  | 3.7 | 6  |
| 14 | Potential usefulness and clinical relevance of a novel left atrial filling index to estimate left ventricular filling pressures in patients with preserved left ventricular ejection fraction. <i>European Heart Journal Cardiovascular Imaging</i> , <b>2020</b> , 21, 260-269 | 4.1 | 5  |
| 13 | Left atrial function and maximal exercise capacity in heart failure with preserved and mid-range ejection fraction. <i>ESC Heart Failure</i> , <b>2021</b> , 8, 116-128   | 3.7 | 4  |
| 12 | Association between functional capacity and heart rate variability in patients with uncomplicated type 2 diabetes. <i>Blood Pressure</i> , <b>2019</b> , 28, 184-190  | 1.7 | 3  |

|    |  |     |   |
|----|--|-----|---|
| 11 | Left atrial strain predicts exercise capacity in heart failure independently of left ventricular ejection fraction.. <i>ESC Heart Failure</i> , <b>2022</b> ,  | 3.7 | 3 |
| 10 | The non-invasive assessment of myocardial work by pressure-strain analysis: clinical applications. <i>Heart Failure Reviews</i> , <b>2021</b> , 1  | 5   | 3 |
| 9  | The CardioMEMS system in the clinical management of end-stage heart failure patients: three case reports. <i>BMC Cardiovascular Disorders</i> , <b>2018</b> , 18, 155  | 2.3 | 2 |
| 8  | Relation of left atrial function with exercise capacity and muscle endurance in patients with heart failure. <i>ESC Heart Failure</i> , <b>2021</b> ,  | 3.7 | 2 |
| 7  | The Prevalence of Iron Deficiency in Patients with Heart Failure with Preserved Ejection Fraction and its Association with Elevated Pulmonary Pressure, Reduced Exercise Capacity and Quality of Life. <i>Journal of Cardiac Failure</i> , <b>2015</b> , 21, S110-S111 | 3.3 | 1 |
| 6  | Clinical perspectives and evidence of diastolic stress test in heart failure with preserved ejection fraction. <i>Egyptian Heart Journal</i> , <b>2015</b> , 67, 279-288   | 1.3 | 1 |
| 5  | A Patient with Quadricuspid Aortic Valve and Ischemic Stroke. <i>Journal of Heart Valve Disease</i> , <b>2016</b> , 25, 456-458  |     | 1 |
| 4  | Functional tricuspid regurgitation, related right heart remodeling, and available treatment options: good news for patients with heart failure?. <i>Heart Failure Reviews</i> , <b>2021</b> , 1  | 5   | 0 |
| 3  | History of acute coronary syndrome: a common, maybe underestimated, risk factor for heart failure with preserved ejection fraction. <i>Revista Espanola De Cardiologia (English Ed)</i> , <b>2021</b> , 74, 480-481  | 0.7 |   |
| 2  | Antecedentes de sīndrome coronario agudo: un factor de riesgo infraestimado de insuficiencia cardiaca con funciōn conservada. <i>Revista Espanola De Cardiologia</i> , <b>2021</b> , 74, 480-481   | 1.5 |   |
| 1  | Left atrial diverticulum-An unexpected finding in routine transesophageal echocardiography. <i>Echocardiography</i> , <b>2021</b> , 38, 147-148  | 1.5 |   |