

Minghuan Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1754785/publications.pdf>

Version: 2024-02-01

40
papers

556
citations

623734

14
h-index

677142

22
g-index

40
all docs

40
docs citations

40
times ranked

1069
citing authors

#	ARTICLE	IF	CITATIONS
1	Patterns and Treatment Strategies of Osimertinib Resistance in T790M-Positive Non-Small Cell Lung Cancer: A Pooled Analysis. <i>Frontiers in Oncology</i> , 2021, 11, 600844.	2.8	6
2	Combined prognostic value of the SUVmax derived from FDG-PET and the lymphocyte-monocyte ratio in patients with stage IIIB-IV non-small cell lung cancer receiving chemotherapy. <i>BMC Cancer</i> , 2021, 21, 66.	2.6	10
3	Short-term response might influence the treatment-related benefit of adjuvant chemotherapy after concurrent chemoradiotherapy for esophageal squamous cell carcinoma patients. <i>Radiation Oncology</i> , 2021, 16, 195.	2.7	1
4	The PET-Derived Tumor-to-Liver Standard Uptake Ratio (SUVTLR) Is Superior to Tumor SUVmax in Predicting Tumor Response and Survival After Chemoradiotherapy in Patients With Locally Advanced Esophageal Cancer. <i>Frontiers in Oncology</i> , 2020, 10, 1630.	2.8	15
5	Lymphocyte-monocyte ratio as a predictive marker for pathological complete response to neoadjuvant therapy in esophageal squamous cell carcinoma. <i>Translational Cancer Research</i> , 2020, 9, 3842-3853.	1.0	11
6	A review of radiation-induced lymphopenia in patients with esophageal cancer: an immunological perspective for radiotherapy. <i>Therapeutic Advances in Medical Oncology</i> , 2020, 12, 175883592092682.	3.2	28
7	A predictive model for treatment response in patients with locally advanced esophageal squamous cell carcinoma after concurrent chemoradiotherapy: based on SUVmean and NLR. <i>BMC Cancer</i> , 2020, 20, 544.	2.6	17
8	The flow-metabolism ratio might predict treatment response and survival in patients with locally advanced esophageal squamous cell carcinoma. <i>EJNMMI Research</i> , 2020, 10, 57.	2.5	2
9	The impact of the nodal status on the overall survival of non-surgical patients with esophageal squamous cell carcinoma. <i>Radiation Oncology</i> , 2019, 14, 161.	2.7	5
10	<p>Primary tumor location is an important predictor of survival in pulmonary adenocarcinoma</p>. <i>Cancer Management and Research</i> , 2019, Volume 11, 2269-2280.	1.9	12
11	<p>Baseline FDG Uptake And Peripheral Lymphocyte-Monocyte Ratio For Predicting Chemoradiotherapy Response In Patients With Esophageal Squamous Cell Carcinoma</p>. <i>Cancer Management and Research</i> , 2019, Volume 11, 9085-9093.	1.9	2
12	Neutrophil-to-lymphocyte ratio is superior to platelet-to-lymphocyte ratio as a prognostic predictor in advanced non-small-cell lung cancer treated with first-line platinum-based chemotherapy. <i>Future Oncology</i> , 2019, 15, 625-635.	2.4	16
13	High expression level of peptidylprolyl isomerase A is correlated with poor prognosis of liver hepatocellular carcinoma. <i>Oncology Letters</i> , 2019, 18, 4691-4702.	1.8	5
14	Radiotherapy for esophageal carcinoma: dose, response and survival. <i>Cancer Management and Research</i> , 2018, Volume 10, 13-21.	1.9	23
15	Nrf2 and Keap1 abnormalities in esophageal squamous cell carcinoma and association with the effect of chemoradiotherapy. <i>Thoracic Cancer</i> , 2018, 9, 726-735.	1.9	28
16	Prognostic role of neutrophilàlymphocyte ratio on esophageal cancer patients who received definitive chemoradiotherapy. <i>Precision Radiation Oncology</i> , 2018, 2, 32-38.	1.1	0
17	Clinical value of carcinoembryonic antigen for predicting the incidence of brain metastases and survival in small cell lung cancer patients treated with prophylactic cranial irradiation. <i>Cancer Management and Research</i> , 2018, Volume 10, 3199-3205.	1.9	9
18	Magnetic resonance imaging evaluation of treatment efficacy and prognosis for brain metastases in lung cancer patients after radiotherapy: A preliminary study. <i>Thoracic Cancer</i> , 2018, 9, 865-873.	1.9	8

#	ARTICLE	IF	CITATIONS
19	Prognostic value of systemic immune-inflammation index in patients with advanced non-small-cell lung cancer. <i>Future Oncology</i> , 2018, 14, 2643-2650.	2.4	30
20	¹⁸ F-fluorodeoxyglucose positron emission tomography predicts lymph node responses to definitive chemoradiotherapy in esophageal squamous cell carcinoma patients. <i>OncoTargets and Therapy</i> , 2018, Volume 11, 4345-4353.	2.0	4
21	Clinical and radiological characteristics of central pulmonary adenocarcinoma: a comparison with central squamous cell carcinoma and small cell lung cancer and the impact on treatment response. <i>OncoTargets and Therapy</i> , 2018, Volume 11, 2509-2517.	2.0	11
22	Stereotactic ablative radiotherapy in treatment of early-stage non-small cell lung cancer: Unsolved questions and frontiers ahead. <i>Cancer Letters</i> , 2017, 401, 46-52.	7.2	1
23	Prognostic significance of the lymphocyte-to-monocyte ratio and the tumor-infiltrating lymphocyte to tumor-associated macrophage ratio in patients with stage T3N0M0 esophageal squamous cell carcinoma. <i>Cancer Immunology, Immunotherapy</i> , 2017, 66, 343-354.	4.2	42
24	The role of metabolic tumor volume (MTV) measured by [18F] FDG PET/CT in predicting EGFR gene mutation status in non-small cell lung cancer. <i>Oncotarget</i> , 2017, 8, 33736-33744.	1.8	19
25	Consolidative thoracic radiotherapy for extensive stage small cell lung cancer. <i>Oncotarget</i> , 2017, 8, 22251-22261.	1.8	11
26	Identification of risk factors and the pattern of lower cervical lymph node metastasis in esophageal cancer: implications for radiotherapy target delineation. <i>Oncotarget</i> , 2017, 8, 43389-43396.	1.8	4
27	Inhibition of hypoxia-inducible factor-1 α by PX-478 as a potential targeted therapy in ESCC.. <i>Journal of Clinical Oncology</i> , 2017, 35, e14083-e14083.	1.6	0
28	Prognostic significance of the lymphocyte-to-monocyte ratio and the tumor-infiltrating lymphocyte to tumor-associated macrophage ratio in patients with stage T3N0M0 esophageal squamous cell carcinoma.. <i>Journal of Clinical Oncology</i> , 2017, 35, e15602-e15602.	1.6	0
29	Association of CD8+/FOXP3+ ratio and PD-L1 expression with survival in pT3N0M0 stage esophageal squamous cell cancer.. <i>Journal of Clinical Oncology</i> , 2017, 35, e15517-e15517.	1.6	0
30	Details of out-field regional recurrence after involved-field irradiation with concurrent chemotherapy for locally advanced esophageal squamous cell carcinoma. <i>OncoTargets and Therapy</i> , 2016, 9, 3049.	2.0	5
31	Mapping patterns of nodal metastases in esophageal carcinoma: rethinking the clinical target volume for supraclavicular nodal irradiation. <i>Journal of Thoracic Disease</i> , 2016, 8, 3132-3138.	1.4	3
32	Intrapericardial bevacizumab safely and effectively treats malignant pericardial effusion in advanced cancer patients. <i>Oncotarget</i> , 2016, 7, 52436-52441.	1.8	13
33	Postoperative radiation in esophageal squamous cell carcinoma and target volume delineation. <i>OncoTargets and Therapy</i> , 2016, Volume 9, 4187-4196.	2.0	14
34	Postoperative radiation therapy of pT2-3N0M0 esophageal carcinoma—a review. <i>Tumor Biology</i> , 2016, 37, 14443-14450.	1.8	1
35	Prognostic Role of Programmed Death Ligand-1 Expression in Breast Cancer: A Systematic Review and Meta-Analysis. <i>Targeted Oncology</i> , 2016, 11, 753-761.	3.6	73
36	How breast cancer chemotherapy increases the risk of leukemia: Thoughts about a case of diffuse large B-cell lymphoma and leukemia after breast cancer chemotherapy. <i>Cancer Biology and Therapy</i> , 2016, 17, 125-128.	3.4	9

#	ARTICLE	IF	CITATIONS
37	CD8+/FOXP3+ ratio and PD-L1 expression associated with survival in pT3N0M0 stage esophageal squamous cell cancer. <i>Oncotarget</i> , 2016, 7, 71455-71465.	1.8	42
38	FDG-PET Predicts Pain Response and Local Control in Palliative Radiotherapy With or Without Systemic Treatment in Patients With Bone Metastasis From Non-small-cell Lung Cancer. <i>Clinical Lung Cancer</i> , 2015, 16, e111-e119.	2.6	14
39	Involved-field irradiation in definitive chemoradiotherapy for locally advanced esophageal squamous cell carcinoma. <i>Radiation Oncology</i> , 2014, 9, 64.	2.7	38
40	Prognostic value of ¹⁸ F-Deoxy- ¹⁸ F-Fluorothymidine ([¹⁸ F] FLT PET) in patients with recurrent malignant gliomas. <i>Nuclear Medicine and Biology</i> , 2014, 41, 710-715.	0.6	24