

Zhaojun Yang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1776481/publications.pdf>

Version: 2024-02-01

13
papers

312
citations

1307543

7
h-index

1125717

13
g-index

18
all docs

18
docs citations

18
times ranked

533
citing authors

#	ARTICLE	IF	CITATIONS
1	Acarbose compared with metformin as initial therapy in patients with newly diagnosed type 2 diabetes: an open-label, non-inferiority randomised trial. <i>Lancet Diabetes and Endocrinology</i> , 2014, 2, 46-55.	11.4	134
2	BMI and waist circumference are associated with impaired glucose metabolism and type 2 diabetes in normal weight Chinese adults. <i>Journal of Diabetes and Its Complications</i> , 2014, 28, 470-476.	2.3	43
3	Stronger associations of waist circumference and waist-to-height ratio with diabetes than BMI in Chinese adults. <i>Diabetes Research and Clinical Practice</i> , 2019, 147, 9-18.	2.8	34
4	Association between Family History Risk Categories and Prevalence of Diabetes in Chinese Population. <i>PLoS ONE</i> , 2015, 10, e0117044.	2.5	31
5	Sex-Specific Prevalence of Diabetes and Cardiovascular Risk Factors in the Middle-Aged Population of China: A Subgroup Analysis of the 2007-2008 China National Diabetes and Metabolic Disorders Study. <i>PLoS ONE</i> , 2015, 10, e0139039.	2.5	17
6	Prevalences and associated factors of electrocardiographic abnormalities in Chinese adults: a cross-sectional study. <i>BMC Cardiovascular Disorders</i> , 2020, 20, 414.	1.7	14
7	Immune-Related Multiple-Organ Injuries Following ICI Treatment With Tislelizumab in an Advanced Non-Small Cell Lung Cancer Patient: A Case Report. <i>Frontiers in Oncology</i> , 2021, 11, 664809.	2.8	10
8	Gender-differential effects on blood glucose levels between acarbose and metformin in Chinese patients with newly diagnosed type 2 diabetes: a sub-analysis of the MARCH trial. <i>Endocrine Journal</i> , 2021, 68, 69-79.	1.6	10
9	Maternal and paternal histories differentially influence risks for diabetes, insulin secretion and insulin resistance in a Chinese population. <i>Journal of Diabetes Investigation</i> , 2021, 12, 434-445.	2.4	7
10	Assessment of three fasting plasma glucose targets for insulin glargine-based therapy in people with type 2 diabetes mellitus in China: study protocol for a randomized controlled trial. <i>Trials</i> , 2016, 17, 470.	1.6	4
11	Distribution and related factors of cardiometabolic disease stage based on body mass index level in Chinese adults- The National Diabetes and Metabolic Disorders Survey. <i>Diabetes/Metabolism Research and Reviews</i> , 2018, 34, e2963.	4.0	4
12	Acarbose Reduces Low-Grade Albuminuria Compared to Metformin in Chinese Patients with Newly Diagnosed Type 2 Diabetes. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2021, Volume 14, 4451-4458.	2.4	3
13	Factors that influence the efficacy of acarbose and metformin as initial therapy in Chinese patients with newly diagnosed type 2 diabetes: a subanalysis of the MARCH trial. <i>Current Medical Research and Opinion</i> , 2016, 32, 713-719.	1.9	1