Bozena Moskwa

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1777029/publications.pdf

Version: 2024-02-01

516710 677142 44 610 16 citations h-index papers

g-index 44 44 44 587 docs citations times ranked citing authors all docs

22

#	Article	IF	CITATIONS
1	The parasitic fauna of the European bison (Bison bonasus) (Linnaeus, 1758) and their impact on the conservation. Part 1 The summarising list of parasites noted. Acta Parasitologica, 2014, 59, 363-71.	1.1	48
2	Comparative analysis of excretory-secretory antigens of Trichinella spiralis and Trichinella britovi muscle larvae by two-dimensional difference gel electrophoresis and immunoblotting. Proteome Science, 2012, 10, 10.	1.7	34
3	Proteomic analysis of potential immunoreactive proteins from muscle larvae and adult worms of Trichinella spiralis in experimentally infected pigs. Folia Parasitologica, 2015, 62, .	1.3	34
4	Seroprevalence of Toxoplasma gondii and Neospora caninum infection in sheep, goats, and fallow deer farmed on the same area1. Journal of Animal Science, 2018, 96, 2468-2473.	0.5	33
5	The parasitic fauna of the European bison (Bison bonasus) (Linnaeus, 1758) and their impact on the conservation. Part 2 The structure and changes over time. Acta Parasitologica, 2014, 59, 372-9.	1.1	24
6	First detection of Trichinella pseudospiralis infection in raccoon (Procyon lotor) in Central Europe. Veterinary Parasitology, 2018, 254, 114-119.	1.8	23
7	The first detection of Neospora caninum DNA in the colostrum of infected cows. Parasitology Research, 2006, 100, 633-636.	1.6	22
8	First report of Trichinella pseudospiralis in Poland, in red foxes (Vulpes vulpes). Acta Parasitologica, 2013, 58, 149-54.	1.1	22
9	Multilocus genotyping of Giardia duodenalis isolates from red deer (Cervus elaphus) and roe deer (Capreolus capreolus) from Poland. Folia Parasitologica, 2012, 59, 237-240.	1.3	22
10	In vitro isolation and identification of the first Neospora caninum isolate from European bison (Bison) Tj ETQq0 0 0) rgBT /Ov	erlock 10 Tf 20
11	Molecular identification of Trichinella britovi in martens (Martes martes) and badgers (Meles meles); new host records in Poland. Acta Parasitologica, 2012, 57, 402-5.	1.1	20
12	Trichinella britovi muscle larvae and adult worms: stage-specific and common antigens detected by two-dimensional gel electrophoresis-based immunoblotting. Parasites and Vectors, 2018, 11, 584.	2.5	20
13	The first identification of a blood-sucking abomasal nematode Ashworthius sidemi in cattle (Bos) Tj ETQq1 1 0.784	1314 rgBT 1.8	/Overlock 1
14	The first detection of nematodes Ashworthius sidemi in elk Alces alces (L.) in Poland and remarks of ashworthiosis foci limitations. Acta Parasitologica, 2013, 58, 515-8.	1.1	18
15	The usefulness of direct agglutination test, enzyme-linked immunosorbent assay and polymerase chain reaction for the detection of Toxoplasma gondii in wild animals. Veterinary Parasitology, 2016, 228, 85-89.	1.8	18
16	Haemonchus contortus: Characterization of the baculovirus expressed form of aminopeptidase H11. Experimental Parasitology, 2007, 117, 208-213.	1.2	17
17	Survey of Toxoplasma gondii and Neospora caninum in raccoons (Procyon lotor) from the Czech Republic, Germany and Poland. Veterinary Parasitology, 2018, 262, 47-50.	1.8	16
18	The Occurrence of <i>Trichinella</i> spp. in Red Foxes (<i>Vulpes vulpes</i>) in Different Regions of Poland: Current Data. Vector-Borne and Zoonotic Diseases, 2016, 16, 717-721.	1.5	15

#	Article	IF	CITATIONS
19	The occurrence and muscle distribution of Trichinella britovi in raccoon dogs (Nyctereutes) Tj ETQq1 1 0.784314 Parasitology: Parasites and Wildlife, 2019, 9, 149-153.	rgBT /Over 1.5	lock 10 Tf 5 15
20	Wild boars meat as a potential source of human trichinellosis in Poland: current data. Acta Parasitologica, 2015, 60, 530-5.	1.1	14
21	The occurrence of nematodes of the genus Trichinella in wolves (Canis lupus) from the Bieszczady Mountains and Augustowska Forest in Poland. Veterinary Parasitology, 2016, 231, 115-117.	1.8	14
22	Intestinal helminths of raccoon dogs (<i>Nyctereutes procyonoides</i>) and red foxes (<i>Vulpes) Tj ETQq0 0 0 rg Research (Poland), 2016, 60, 273-277.</i>	gBT /Overlo 1.0	ock 10 Tf 50 12
23	The Nematodes Thelazia gulosa Railiet and Henry, 1910 and Thelazia skrjabini Erschov, 1928 as a Cause of Blindness in European Bison (Bison bonasus) in Poland. Acta Parasitologica, 2020, 65, 963-968.	1.1	12
24	Detection of specific antibodies anti-Neospora caninum in the fallow deer (Dama dama). Research in Veterinary Science, 2012, 92, 96-98.	1.9	11
25	Recognition of antigens of three different stages of the Trichinella spiralis by antibodies from pigs infected with T. spiralis. Experimental Parasitology, 2013, 134, 129-137.	1.2	10
26	Immunoproteomic analysis of Trichinella spiralis and Trichinella britovi excretory-secretory muscle larvae proteins recognized by sera from humans infected with Trichinella. PLoS ONE, 2020, 15, e0241918.	2.5	10
27	The first report of Toxoplasma gondii antibodies in free-living European bison (Bison bonasus bonasus) Tj ETQq1	. 0. 784314	· fgBT /Overl
28	Seroprevalence of Trichinella spp. infection in bank voles (Myodes glareolus) – A long term study. International Journal for Parasitology: Parasites and Wildlife, 2019, 9, 144-148.	1.5	7
29	Detection of antibodies to Neospora caninum in moose (Alces alces): the first report in Europe. Folia Parasitologica, 2014, 61, 34-36.	1.3	7
30	Studies on Neospora caninum DNA detection in the oocytes and embryos collected from infected cows. Veterinary Parasitology, 2008, 158, 370-375.	1.8	6
31	The usefulness of DNA derived from third stage larvae in the detection of Ashworthius sidemi infection in European bison, by a simple polymerase chain reaction. Parasites and Vectors, 2014, 7, 215.	2.5	6
32	First Toxoplasma gondii isolate from an aborted foetus of European bison (Bison bonasus bonasus L.). Parasitology Research, 2017, 116, 2457-2461.	1.6	6
33	Trichinella britovi infection and muscle distribution in free-living martens (Martes spp.) from the GÅ,Ä™boki Bród Forest District, Poland. International Journal for Parasitology: Parasites and Wildlife, 2020, 12, 176-180.	1.5	6
34	The Seroprevalence of Toxoplasma gondii in Wild Boars from Three Voivodeships in Poland, MAT Analyses. Acta Parasitologica, 2020, 65, 490-495.	1.1	6
35	Exploiting the potential of 2D DIGE and 2DE immunoblotting for comparative analysis of crude extract of Trichinella britovi and Trichinella spiralis muscle larvae proteomes. Veterinary Parasitology, 2021, 289, 109323.	1.8	6

The first report of Toxoplasma gondii antibodies in free-living European bison (Bison bonasus bonasus) Tj ETQq0 0 Q rgBT /Overlock 10 T

#	Article	IF	CITATIONS
37	Comparison of sensitivity of two primer sets for the detection of Toxoplasma gondii DNA in wildlife. Acta Parasitologica, 2018, 63, 634-639.	1.1	4
38	Sarcocystis cruzi infection in free-living European bison (Bison bonasus bonasus L.) from the BiaÅ,owieÅ⅓a Forest, Poland – A molecular analysis based on the cox1 gene. International Journal for Parasitology: Parasites and Wildlife, 2021, 16, 59-63.	1.5	3
39	The estimation of different ELISA procedures for serodiagnosis of human trichinellosis. Annals of Parasitology, 2006, 52, 231-8.	0.1	3
40	Detection of antibodies to Neospora caninum in moose (Alces alces): the first report in Europe. Folia Parasitologica, 2014, 61, 34-6.	1.3	3
41	Ashworthius sidemi in cattle and wild ruminants in Poland - the current state of play. Annals of Parasitology, 2020, 66, 517-520.	0.1	3
42	Molecular identification of sarcocysts from tissue of fallow deer (Dama dama) farmed in the open pasture system based on ssu rRNA gene. Acta Parasitologica, 2020, 65, 354-360.	1.1	2
43	Use of meat juice from racoons (Procyon lotor) collected from Central Europe for immunological detection of Trichinella spp Veterinary Parasitology, 2021, 297, 109066.	1.8	2
44	Kinetics of anti-Neospora antibodies during the period of two consecutive pregnancies in chronically infected dairy cows. Acta Parasitologica, 2013, 58, 463-7.	1.1	1