

João Eduardo de Araujo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1777408/publications.pdf>

Version: 2024-02-01

36
papers

790
citations

623734

14
h-index

501196

28
g-index

36
all docs

36
docs citations

36
times ranked

949
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Neurochemical mechanisms of the defensive behavior in the dorsal midbrain. <i>Neuroscience and Biobehavioral Reviews</i> , 1999, 23, 863-875. | 6.1 | 263 |
| 2 | Anxiogenic effects of substance P and its 7â€“11 C terminal, but not the 1â€“7 N terminal, injected into the dorsal periaqueductal gray. <i>Peptides</i> , 1999, 20, 1437-1443. | 2.4 | 47 |
| 3 | Nitric oxide synthesis blockade reduced the baroreflex sensitivity in trained rats. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2009, 150, 38-44. | 2.8 | 40 |
| 4 | Aversive effects of the C-fragment of Substance P in the dorsal periaqueductal gray matter. <i>Experimental Brain Research</i> , 1998, 123, 84-89. | 1.5 | 37 |
| 5 | Opposite effects of substance P fragments C (anxiogenic) and N (anxiolytic) injected into dorsal periaqueductal gray. <i>European Journal of Pharmacology</i> , 2001, 432, 43-51. | 3.5 | 32 |
| 6 | Laser photobiomodulation (830 and 660Ånm) in mast cells, VEGF, FGF, and CD34 of the musculocutaneous flap in rats submitted to nicotine. <i>Lasers in Medical Science</i> , 2017, 32, 335-341. | 2.1 | 32 |
| 7 | Modified constraint-induced movement therapy and modified forced-use therapy for stroke patients are both effective to promote balance and gait improvements. <i>Brazilian Journal of Physical Therapy</i> , 2012, 16, 157-165. | 2.5 | 27 |
| 8 | Polycystic ovary syndrome associated with increased adiposity interferes with serum levels of TNF-alpha and IL-6 differently from leptin and adiponectin. <i>Archives of Endocrinology and Metabolism</i> , 2020, 64, 4-10. | 0.6 | 25 |
| 9 | Macroscopic and histological effects of magnetic field exposition in the process of tissue reparation in Wistar rats. <i>Archives of Dermatological Research</i> , 2006, 298, 121-126. | 1.9 | 21 |
| 10 | The immediate effects of local and adjacent acupuncture on the tibialis anterior muscle: a human study. <i>Chinese Medicine</i> , 2008, 3, 17. | 4.0 | 21 |
| 11 | Place aversion induced by microinjections of C-fragment of substance P into the dorsal periaqueductal gray of rats is mediated by tachykinin NK1 receptors. <i>Peptides</i> , 2001, 22, 1447-1452. | 2.4 | 20 |
| 12 | Prevalence and characteristics of chronic pain in Brazil: a national internet-based survey study. <i>Brazilian Journal of Pain</i> , 2018, 1, . | 0.1 | 20 |
| 13 | Neuropathology and behavioral impairments after bilateral global ischemia surgery and exposure to static magnetic field: Evidence in the motor cortex, the hippocampal CA1 region and the neostriatum. <i>International Journal of Radiation Biology</i> , 2013, 89, 595-601. | 1.8 | 17 |
| 14 | Influence of treadmill training on motor performance and organization of exploratory behavior in Meriones unguiculatus with unilateral ischemic stroke: Histological correlates in hippocampal CA1 region and the neostriatum. <i>Neuroscience Letters</i> , 2008, 431, 179-183. | 2.1 | 16 |
| 15 | Chronic cholinergic stimulation promotes changes in cardiovascular autonomic control in spontaneously hypertensive rats. <i>Autonomic Neuroscience: Basic and Clinical</i> , 2015, 193, 97-103. | 2.8 | 16 |
| 16 | Neuropathology and behavioral impairments after three types of global ischemia surgery in Meriones unguiculatus: Evidence in motor cortex, hippocampal CA1 region and the neostriatum. <i>Journal of the Neurological Sciences</i> , 2012, 312, 73-78. | 0.6 | 15 |
| 17 | Anxiogenic effects of activation of NK-1 receptors of the dorsal periaqueductal gray as assessed by the elevated plus-maze, ultrasound vocalizations and tail-flick tests. <i>Neuropeptides</i> , 2007, 41, 365-374. | 2.2 | 13 |
| 18 | Unilateral and Immediate Stimulation of Acupuncture Points Xiaohai (SI8) and Jianwaishu (SI14) of the Small Intestine Meridian Increases Electromyographic Activity and Strength in the Ipsilateral and Contralateral Upper Trapezius Muscle. <i>JAMS Journal of Acupuncture and Meridian Studies</i> , 2016, 9, 250-256. | 0.7 | 13 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Electromyographic and Strength Analyses of Activation Patterns of the Wrist Flexor Muscles after Acupuncture. JAMS Journal of Acupuncture and Meridian Studies, 2014, 7, 231-237. | 0.7 | 12 |
| 20 | Behavioural profile of Wistar rats with unilateral striatal lesion by quinolinic acid (animal model of Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Brain Research, 2015, 233, 1455-1462. | 1.5 | 11 |
| 21 | Analgesic Effects Elicited by Neuroactive Mediators Injected into the ST 36 Acupuncture Point on Inflammatory and Neuropathic Pain in Mice. JAMS Journal of Acupuncture and Meridian Studies, 2018, 11, 280-289. | 0.7 | 11 |
| 22 | Physical training associated with Enalapril but not to Losartan, results in better cardiovascular autonomic effects. Autonomic Neuroscience: Basic and Clinical, 2017, 203, 33-40. | 2.8 | 10 |
| 23 | Neuropathology and behavioral impairments in Wistar rats with a 6-OHDA lesion in the substantia nigra compacta and exposure to a static magnetic field. Electromagnetic Biology and Medicine, 2013, 32, 527-535. | 1.4 | 9 |
| 24 | Behavioral effects and neural changes induced by continuous and not continuous treadmill training, post bilateral cerebral ischemia in gerbils. Behavioural Brain Research, 2015, 291, 20-25. | 2.2 | 9 |
| 25 | IrradiaÃ§Ã£o contralateral de forÃ§a para a ativaÃ§Ã£o do mÃºsculo tibial anterior em portadores da doenÃ§a de Charcot-Marie-Tooth: efeitos de um programa de intervenÃ§Ã£o por FNP. Brazilian Journal of Physical Therapy, 2009, 13, 438-443. | 2.5 | 8 |
| 26 | Amlodipine and enalapril promote distinct effects on cardiovascular autonomic control in spontaneously hypertensive rats. Journal of Hypertension, 2016, 34, 2383-2392. | 0.5 | 8 |
| 27 | Hyperbaric oxygen therapy reduces astrogliosis and helps to recovery brain damage in hydrocephalic young rats. Child's Nervous System, 2018, 34, 1125-1134. | 1.1 | 7 |
| 28 | Single or Multiple Electroacupuncture Sessions in Nonspecific Low Back Pain: Are We Low-Responders to Electroacupuncture?. JAMS Journal of Acupuncture and Meridian Studies, 2018, 11, 54-61. | 0.7 | 7 |
| 29 | Polycystic Ovary Syndrome Presents Higher Sympathetic Cardiac Autonomic Modulation that is not altered by Strength Training. International Journal of Exercise Science, 2016, 9, 554-566. | 0.5 | 7 |
| 30 | Ipsilateral proprioceptive neuromuscular facilitation patterns improve overflow and reduce foot drop in patients with demyelinating polyneuropathy. Journal of Exercise Rehabilitation, 2018, 14, 503-508. | 1.0 | 5 |
| 31 | Aerobic Exercise After Left-Sided Stroke Improves Gait Speed and Endurance. American Journal of Physical Medicine and Rehabilitation, 2021, 100, 576-583. | 1.4 | 5 |
| 32 | Study of Heart Rate Variability and Stress Markers in Basketball Players Submitted to Selective Loads Periodization System. American Journal of Sports Science, 2015, 3, 46. | 0.2 | 4 |
| 33 | Continuous and not continuous 2-week treadmill training enhances the performance in the passive avoidance test in ischemic gerbils. Neuroscience Letters, 2018, 665, 170-175. | 2.1 | 2 |
| 34 | Lower limb auriculotherapy points improves balance in young healthy subjects-assessed by computerized baropodometry. Complementary Therapies in Clinical Practice, 2018, 32, 46-50. | 1.7 | 0 |
| 35 | Effects of Physical Training on Cardiac Modulation in Normal Weight, Overweight, and Obese Individuals: A Comparative Study. Journal of Nutrition & Food Sciences, 2015, 05, . | 1.0 | 0 |
| 36 | AnÃ¡lise da interaÃ§Ã£o funcional do campo magnÃ©tico contÃnuo pÃ³s-injeÃ§Ã£o de drogas ansiolÃticas e ansiogÃnicas, ativadoras e depressoras do comportamento motor, uma abordagem experimental em roedores. , 0, , . | | 0 |