

Hao Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1777771/publications.pdf>

Version: 2024-02-01

10
papers

506
citations

1040056

9
h-index

1372567

10
g-index

11
all docs

11
docs citations

11
times ranked

841
citing authors

#	ARTICLE	IF	CITATIONS
1	Entopeduncular Nucleus Projections to the Lateral Habenula Contribute to Cocaine Avoidance. <i>Journal of Neuroscience</i> , 2021, 41, 298-306.	3.6	13
2	Synaptic Adaptations at the Rostromedial Tegmental Nucleus Underlie Individual Differences in Cocaine Avoidance Behavior. <i>Journal of Neuroscience</i> , 2021, 41, 4620-4630.	3.6	7
3	Three Rostromedial Tegmental Afferents Drive Triply Dissociable Aspects of Punishment Learning and Aversive Valence Encoding. <i>Neuron</i> , 2019, 104, 987-999.e4.	8.1	48
4	Local inhibition of uptake2 transporters augments stress-induced increases in serotonin in the rat central amygdala. <i>Neuroscience Letters</i> , 2019, 701, 119-124.	2.1	11
5	Valence-encoding in the lateral habenula arises from the entopeduncular region. <i>ELife</i> , 2019, 8, .	6.0	34
6	Generality and opponency of rostromedial tegmental (RMTg) roles in valence processing. <i>ELife</i> , 2019, 8, .	6.0	25
7	Cocaine Self-Administration and Extinction Leads to Reduced Glial Fibrillary Acidic Protein Expression and Morphometric Features of Astrocytes in the Nucleus Accumbens Core. <i>Biological Psychiatry</i> , 2016, 80, 207-215.	1.3	133
8	Cq-DREADD Selectively Initiates Glial Glutamate Release and Inhibits Cue-induced Cocaine Seeking. <i>Biological Psychiatry</i> , 2015, 78, 441-451.	1.3	156
9	Serotonergic responses to stress are enhanced in the central amygdala and inhibited in the ventral hippocampus during amphetamine withdrawal. <i>European Journal of Neuroscience</i> , 2014, 40, 3684-3692.	2.6	17
10	Anxiolytic function of the orexin 2/hypocretin A receptor in the basolateral amygdala. <i>Psychoneuroendocrinology</i> , 2014, 40, 17-26.	2.7	59