

Alejandro JimÃ©nez-Sosa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1788270/publications.pdf>

Version: 2024-02-01

172
papers

4,075
citations

136885

32
h-index

149623

56
g-index

178
all docs

178
docs citations

178
times ranked

4899
citing authors

#	ARTICLE	IF	CITATIONS
1	Central venous catheter-related infection in a prospective and observational study of 2,595 catheters. <i>Critical Care</i> , 2005, 9, R631.	2.5	257
2	Influence of an Endotracheal Tube with Polyurethane Cuff and Subglottic Secretion Drainage on Pneumonia. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2007, 176, 1079-1083.	2.5	242
3	Estimated GFR: time for a critical appraisal. <i>Nature Reviews Nephrology</i> , 2019, 15, 177-190.	4.1	187
4	Meropenem by Continuous Versus Intermittent Infusion in Ventilator-Associated Pneumonia due to Gram-Negative Bacilli. <i>Annals of Pharmacotherapy</i> , 2006, 40, 219-223.	0.9	168
5	Clinical cure of ventilator-associated pneumonia treated with piperacillin/tazobactam administered by continuous or intermittent infusion. <i>International Journal of Antimicrobial Agents</i> , 2009, 33, 464-468.	1.1	120
6	Microorganisms responsible for intravascular catheter-related bloodstream infection according to the catheter site*. <i>Critical Care Medicine</i> , 2007, 35, 2424-2427.	0.4	119
7	Matrix metalloproteinase-9, -10, and tissue inhibitor of matrix metalloproteinases-1 blood levels as biomarkers of severity and mortality in sepsis. <i>Critical Care</i> , 2009, 13, R158.	2.5	105
8	Diagnostic accuracy of immunochemical versus guaiac faecal occult blood tests for colorectal cancer screening. <i>Journal of Gastroenterology</i> , 2010, 45, 703-712.	2.3	105
9	Relationship between serum magnesium and parathyroid hormone levels in hemodialysis patients. <i>American Journal of Kidney Diseases</i> , 1999, 34, 43-48.	2.1	93
10	Endoscopic submucosal dissection training with pig models in a Western country. <i>World Journal of Gastroenterology</i> , 2010, 16, 2895.	1.4	88
11	Comparison of clinical cure rates in adults with ventilator-associated pneumonia treated with intravenous ceftazidime administered by continuous or intermittent infusion: A retrospective, nonrandomized, open-label, historical chart review. <i>Clinical Therapeutics</i> , 2007, 29, 2433-2439.	1.1	73
12	Tracheal suction by closed system without daily change versus open system. <i>Intensive Care Medicine</i> , 2006, 32, 538-544.	3.9	68
13	Red Blood Cell Distribution Width during the First Week Is Associated with Severity and Mortality in Septic Patients. <i>PLoS ONE</i> , 2014, 9, e105436.	1.1	68
14	Effects of Colon Capsule Endoscopy on Medical Decision Making in Patients With Incomplete Colonoscopies. <i>Clinical Gastroenterology and Hepatology</i> , 2013, 11, 534-540.e1.	2.4	64
15	Arterial catheter-related infection of 2,949 catheters. <i>Critical Care</i> , 2006, 10, R83.	2.5	60
16	Association of Sepsis-Related Mortality with Early Increase of TIMP-1/MMP-9 Ratio. <i>PLoS ONE</i> , 2014, 9, e94318.	1.1	60
17	Catheter-related infection in critically ill patients. <i>Intensive Care Medicine</i> , 2004, 30, 1681-4.	3.9	58
18	Accidental catheter removal in critically ill patients: a prospective and observational study. <i>Critical Care</i> , 2004, 8, R229.	2.5	55

#	ARTICLE	IF	CITATIONS
19	Ventilator-associated pneumonia using a heated humidifier or a heat and moisture exchanger: a randomized controlled trial [ISRCTN88724583]. <i>Critical Care</i> , 2006, 10, R116.	2.5	55
20	Association between serum soluble CD40 ligand levels and mortality in patients with severe sepsis. <i>Critical Care</i> , 2011, 15, R97.	2.5	53
21	Comparison of Two Intensive Bowel Cleansing Regimens in Patients With Previous Poor Bowel Preparation: A Randomized Controlled Study. <i>American Journal of Gastroenterology</i> , 2017, 112, 951-958.	0.2	51
22	Prognostic Value of Malondialdehyde Serum Levels in Severe Sepsis: A Multicenter Study. <i>PLoS ONE</i> , 2013, 8, e53741.	1.1	49
23	The impact of frailty in older patients with non-ischæmic cardiomyopathy after implantation of cardiac resynchronization therapy defibrillator. <i>Europace</i> , 2015, 17, 598-602.	0.7	47
24	The Use of Rifampicinâ€Miconazoleâ€Impregnated Catheters Reduces the Incidence of Femoral and Jugular Catheterâ€Related Bacteremia. <i>Clinical Infectious Diseases</i> , 2008, 47, 1171-1175.	2.9	46
25	Serum substance P levels are associated with severity and mortality in patients with severe traumatic brain injury. <i>Critical Care</i> , 2015, 19, 192.	2.5	41
26	Factors influencing pregnancy outcome in women with type 2 versus type 1 diabetes mellitus. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2008, 87, 43-49.	1.3	40
27	Impact of a 1-day versus 3-day low-residue diet on bowel cleansing quality before colonoscopy: a randomized controlled trial. <i>Endoscopy</i> , 2019, 51, 628-636.	1.0	40
28	Interatrial block and atrial remodeling assessed using speckle tracking echocardiography. <i>BMC Cardiovascular Disorders</i> , 2018, 18, 38.	0.7	36
29	Association between red blood cell distribution width and mortality of COVID-19 patients. <i>Anaesthesia, Critical Care & Pain Medicine</i> , 2021, 40, 100777.	0.6	36
30	Validation of Fujinon intelligent chromoendoscopy with high definition endoscopes in colonoscopy. <i>World Journal of Gastroenterology</i> , 2009, 15, 5266.	1.4	34
31	Association between Serum Tissue Inhibitor of Matrix Metalloproteinase-1 Levels and Mortality in Patients with Severe Brain Trauma Injury. <i>PLoS ONE</i> , 2014, 9, e94370.	1.1	34
32	Serum caspase-3 levels and mortality are associated in patients with severe traumatic brain injury. <i>BMC Neurology</i> , 2015, 15, 228.	0.8	34
33	Up-regulation of gelatinases in the colorectal adenoma-carcinoma sequence. <i>European Journal of Cancer</i> , 2006, 42, 3246-3252.	1.3	33
34	Serum Malondialdehyde Levels in Patients with Malignant Middle Cerebral Artery Infarction Are Associated with Mortality. <i>PLoS ONE</i> , 2015, 10, e0125893.	1.1	33
35	The 372 T/C genetic polymorphism of TIMP-1 is associated with serum levels of TIMP-1 and survival in patients with severe sepsis. <i>Critical Care</i> , 2013, 17, R94.	2.5	31
36	Sustained high plasma plasminogen activator inhibitor-1 levels are associated with severity and mortality in septic patients. <i>Thrombosis Research</i> , 2014, 134, 182-186.	0.8	31

#	ARTICLE	IF	CITATIONS
37	Platelet cytochrome c oxidase activity and quantity in septic patients*. Critical Care Medicine, 2011, 39, 1289-1294.	0.4	30
38	Association between total antioxidant capacity and mortality in ischemic stroke patients. Annals of Intensive Care, 2016, 6, 39.	2.2	30
39	Impaired HDL cholesterol efflux capacity in systemic lupus erythematosus patients is related to subclinical carotid atherosclerosis. Rheumatology, 2020, 59, 2847-2856.	0.9	30
40	Iohexol plasma clearance simplified by dried blood spot testing. Nephrology Dialysis Transplantation, 2018, 33, 1597-1603.	0.4	29
41	Advanced interatrial block is a surrogate for left atrial strain reduction which predicts atrial fibrillation and stroke. Annals of Noninvasive Electrocardiology, 2019, 24, e12632.	0.5	29
42	Serum Levels of Caspase-Cleaved Cytokeratin-18 in Patients with Severe Traumatic Brain Injury Are Associated with Mortality: A Pilot Study. PLoS ONE, 2015, 10, e0121739.	1.1	29
43	Serum Levels of Caspase-Cleaved Cytokeratin-18 and Mortality Are Associated in Severe Septic Patients: Pilot Study. PLoS ONE, 2014, 9, e109618.	1.1	28
44	Association between serum total antioxidant capacity and mortality in severe septic patients. Journal of Critical Care, 2015, 30, 217.e7-217.e12.	1.0	28
45	Risk for high-grade dysplasia or invasive carcinoma in colorectal flat adenomas in a Spanish population. Gastroenterología Y Hepatología, 2006, 29, 602-609.	0.2	26
46	Association between Interleukin-6 Promoter Polymorphism (-174 G/C), Serum Interleukin-6 Levels and Mortality in Severe Septic Patients. International Journal of Molecular Sciences, 2016, 17, 1861.	1.8	26
47	Decrease of oxidative phosphorylation system function in severe septic patients. Journal of Critical Care, 2015, 30, 935-939.	1.0	25
48	Impact of physician specialty on the cost of nonvariceal upper GI bleeding care. American Journal of Gastroenterology, 2002, 97, 1535-1542.	0.2	24
49	Chlorhexidine-silver sulfadiazine-impregnated venous catheters save costs. American Journal of Infection Control, 2014, 42, 321-324.	1.1	24
50	Serum soluble CD40 Ligand levels are associated with severity and mortality of brain trauma injury patients. Thrombosis Research, 2014, 134, 832-836.	0.8	23
51	Association between Pre-Transplant Serum Malondialdehyde Levels and Survival One Year after Liver Transplantation for Hepatocellular Carcinoma. International Journal of Molecular Sciences, 2016, 17, 500.	1.8	23
52	Low-Dose 3 mg Levobupivacaine Plus 10 μ g Fentanyl Selective Spinal Anesthesia for Gynecological Outpatient Laparoscopy. Anesthesia and Analgesia, 2009, 109, 1456-1461.	1.1	22
53	Survival and mitochondrial function in septic patients according to mitochondrial DNA haplogroup. Critical Care, 2012, 16, R10.	2.5	22
54	Chronic kidney disease staging with cystatin C or creatinine-based formulas: flipping the coin. Nephrology Dialysis Transplantation, 2019, 34, 287-294.	0.4	22

#	ARTICLE	IF	CITATIONS
55	Total antioxidant capacity is associated with mortality of patients with severe traumatic brain injury. <i>BMC Neurology</i> , 2015, 15, 115.	0.8	20
56	Association between serum substance P levels and mortality in patients with severe sepsis. <i>Journal of Critical Care</i> , 2015, 30, 924-928.	1.0	20
57	New founding mutation in MSH2 associated with hereditary nonpolyposis colorectal cancer syndrome on the Island of Tenerife. <i>Cancer Letters</i> , 2006, 244, 268-273.	3.2	19
58	Internal jugular venous catheter-related bacteremia according to central and posterior accesses. <i>Intensive Care Medicine</i> , 2007, 33, 1071-1075.	3.9	18
59	Serum melatonin levels are associated with mortality in severe septic patients. <i>Journal of Critical Care</i> , 2015, 30, 860.e1-860.e6.	1.0	18
60	Chlorhexidine-silver sulfadiazine or rifampicin-miconazole impregnated venous catheters decrease the risk of catheter-related bloodstream infection similarly. <i>American Journal of Infection Control</i> , 2016, 44, 50-53.	1.1	18
61	High prevalence of overweight in a stable spanish hemodialysis population: A cross sectional study. , 2003, 13, 52-59.		17
62	Severe Septic Patients with Mitochondrial DNA Haplogroup JT Show Higher Survival Rates: A Prospective, Multicenter, Observational Study. <i>PLoS ONE</i> , 2013, 8, e73320.	1.1	17
63	Serum melatonin levels in survivor and non-survivor patients with traumatic brain injury. <i>BMC Neurology</i> , 2017, 17, 138.	0.8	17
64	Serum Malondialdehyde Levels and Mortality in Patients with Spontaneous Intracerebral Hemorrhage. <i>World Neurosurgery</i> , 2018, 113, e542-e547.	0.7	17
65	Lower associated costs using rifampicin-miconazole impregnated catheters compared with standard catheters. <i>American Journal of Infection Control</i> , 2011, 39, 895-897.	1.1	16
66	Measurement of glomerular filtration rate: Internal and external validations of the iohexol plasma clearance technique by HPLC. <i>Clinica Chimica Acta</i> , 2014, 430, 84-85.	0.5	16
67	Prognostic Value of Serum Caspase-Cleaved Cytokeratin-18 Levels before Liver Transplantation for One-Year Survival of Patients with Hepatocellular Carcinoma. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1524.	1.8	16
68	Association Between Bendopnea and Key Parameters of Cardiopulmonary Exercise Testing in Patients With Advanced Heart Failure. <i>Journal of Cardiac Failure</i> , 2016, 22, 163-165.	0.7	16
69	One-day versus two-day cleansing for colon capsule endoscopy: a prospective randomized pilot study. <i>Gastroenterología Y Hepatología</i> , 2014, 37, 101-106.	0.2	15
70	Serum total antioxidant capacity during the first week of sepsis and mortality. <i>Journal of Critical Care</i> , 2018, 47, 139-144.	1.0	15
71	The Error of Estimated GFR in Type 2 Diabetes Mellitus. <i>Journal of Clinical Medicine</i> , 2019, 8, 1543.	1.0	15
72	DNA and RNA Oxidative Damage and Mortality of Patients With COVID-19. <i>American Journal of the Medical Sciences</i> , 2021, 361, 585-590.	0.4	15

#	ARTICLE	IF	CITATIONS
73	Subglottic secretion drainage and continuous control of cuff pressure used together save health care costs. <i>American Journal of Infection Control</i> , 2014, 42, 1101-1105.	1.1	14
74	The 4G/4G Genotype of PAI-1 Polymorphism Is Associated with Higher Plasma PAI-1 Concentrations and Mortality in Patients with Severe Sepsis. <i>PLoS ONE</i> , 2015, 10, e0129565.	1.1	14
75	Efficiency of chlorhexidine-silver sulfadiazine-impregnated venous catheters at subclavian sites. <i>American Journal of Infection Control</i> , 2015, 43, 711-714.	1.1	14
76	Serum caspase 3 levels are associated with early mortality in severe septic patients. <i>Journal of Critical Care</i> , 2016, 34, 103-106.	1.0	14
77	Serum melatonin levels are associated with mortality in patients with malignant middle cerebral artery infarction. <i>Journal of International Medical Research</i> , 2018, 46, 3268-3277.	0.4	14
78	Impact of errors of creatinine and cystatin C equations in the selection of living kidney donors. <i>CKJ: Clinical Kidney Journal</i> , 2019, 12, 748-755.	1.4	14
79	Early Mortality of Brain Infarction Patients and Red Blood Cell Distribution Width. <i>Brain Sciences</i> , 2020, 10, 196.	1.1	14
80	Low-Dose, Low-Concentration Levobupivacaine Plus Fentanyl Selective Spinal Anesthesia for Knee Arthroscopy. <i>Anesthesia and Analgesia</i> , 2011, 112, 477-480.	1.1	13
81	Levosimendan Improves Hemodynamic Status in Critically Ill Patients with Severe Aortic Stenosis and Left Ventricular Dysfunction: An Interventional Study. <i>Cardiovascular Therapeutics</i> , 2015, 33, 193-199.	1.1	13
82	Persistently High Serum Substance P Levels and Early Mortality in Patients with Severe Traumatic Brain Injury. <i>World Neurosurgery</i> , 2019, 132, e613-e617.	0.7	13
83	Colorectal cancer screening in a Spanish population. <i>Medicina Clínica</i> , 2009, 133, 736-740.	0.3	12
84	Association between Serum Soluble CD154 Levels and Mortality in Patients with Malignant Middle Cerebral Artery Infarction. <i>International Journal of Molecular Sciences</i> , 2015, 16, 12147-12158.	1.8	12
85	Serum tissue inhibitor of matrix metalloproteinase-1 levels are associated with mortality in patients with malignant middle cerebral artery infarction. <i>BMC Neurology</i> , 2015, 15, 111.	0.8	11
86	Plasma matrix metalloproteinase 9 as an early surrogate biomarker of advanced colorectal neoplasia. <i>Gastroenterología y Hepatología</i> , 2016, 39, 433-441.	0.2	11
87	Non-survivor septic patients have persistently higher serum sCD40L levels than survivors. <i>Journal of Critical Care</i> , 2017, 41, 177-182.	1.0	11
88	The Serum Melatonin Levels and Mortality of Patients with Spontaneous Intracerebral Hemorrhage. <i>Brain Sciences</i> , 2019, 9, 263.	1.1	11
89	Association Between DNA and RNA Oxidative Damage and Mortality of Patients with Traumatic Brain Injury. <i>Neurocritical Care</i> , 2020, 32, 790-795.	1.2	11
90	Red blood cell distribution width and mortality of spontaneous intracerebral hemorrhage patients. <i>Clinical Neurology and Neurosurgery</i> , 2020, 195, 106066.	0.6	11

#	ARTICLE	IF	CITATIONS
91	Extracellular water/total body water ratio as predictor of mortality in hemodialysis patients. <i>Renal Failure</i> , 2021, 43, 821-829.	0.8	11
92	Cost/benefit analysis of chlorhexidine-silver sulfadiazine-impregnated venous catheters for femoral access. <i>American Journal of Infection Control</i> , 2014, 42, 1130-1132.	1.1	10
93	A Simple Method to Measure Renal Function in Swine by the Plasma Clearance of Iohexol. <i>International Journal of Molecular Sciences</i> , 2018, 19, 232.	1.8	10
94	High serum levels of caspase-cleaved cytokeratin-18 are associated with malignant middle cerebral artery infarction patient mortality. <i>BMC Neurology</i> , 2018, 18, 32.	0.8	10
95	Association between DNA and RNA oxidative damage and mortality in septic patients. <i>Journal of Critical Care</i> , 2019, 54, 94-98.	1.0	10
96	Relationship Between Insulin Sensitivity and β -Cell Secretion in Nondiabetic Subjects with Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2019, 46, 229-236.	1.0	10
97	Proprotein convertase subtilisin/kexin type 9 in the dyslipidaemia of patients with axial spondyloarthritis is related to disease activity. <i>Rheumatology</i> , 2021, 60, 2296-2306.	0.9	10
98	Set of Quality Indicators of Pediatric Intensive Care in Spain: Delphi Method Selection. <i>Pediatric Quality & Safety</i> , 2017, 2, e009.	0.4	9
99	Serum melatonin levels during the first seven days of severe sepsis diagnosis are associated with sepsis severity and mortality. <i>Enfermedades Infecciosas Y Microbiología Clínica</i> , 2018, 36, 544-549.	0.3	9
100	Predictive ability of the EVARUCI scale and COMHON index for pressure injury risk in critically ill patients: A diagnostic accuracy study. <i>Australian Critical Care</i> , 2018, 31, 355-361.	0.6	9
101	High Serum DNA and RNA Oxidative Damage in Non-surviving Patients with Spontaneous Intracerebral Hemorrhage. <i>Neurocritical Care</i> , 2020, 33, 90-96.	1.2	9
102	Septic patients with mitochondrial DNA haplogroup JT have higher respiratory complex IV activity and survival rate. <i>Journal of Critical Care</i> , 2016, 33, 95-99.	1.0	8
103	Blood concentrations of proapoptotic sFas and antiapoptotic Bcl2 and COVID-19 patient mortality. <i>Expert Review of Molecular Diagnostics</i> , 2021, 21, 837-844.	1.5	8
104	Sustained Low Serum Substance P Levels in Non-Surviving Septic Patients. <i>International Journal of Molecular Sciences</i> , 2017, 18, 1531.	1.8	7
105	High serum soluble CD40L levels previously to liver transplantation in patients with hepatocellular carcinoma are associated with mortality at one year. <i>Journal of Critical Care</i> , 2018, 43, 316-320.	1.0	7
106	Traumatic Brain Injury Patients Mortality and Serum Total Antioxidant Capacity. <i>Brain Sciences</i> , 2020, 10, 110.	1.1	7
107	Association between neutrophil-to-lymphocyte ratio in the first seven days of sepsis and mortality. <i>Enfermedades Infecciosas Y Microbiología Clínica</i> , 2020, , .	0.3	7
108	Lower arterial catheter-related infection in brachial than in femoral access. <i>American Journal of Infection Control</i> , 2010, 38, e40-e42.	1.1	6

#	ARTICLE	IF	CITATIONS
109	Association between serum levels of caspase-cleaved cytokeratin-18 and early mortality in patients with severe spontaneous intracerebral hemorrhage. <i>BMC Neuroscience</i> , 2018, 19, 23.	0.8	6
110	High serum levels of tissue inhibitor of matrix metalloproteinase-1 during the first week of a malignant middle cerebral artery infarction in non-surviving patients. <i>BMC Neurology</i> , 2019, 19, 167.	0.8	6
111	Maintained high sustained serum malondialdehyde levels after severe brain trauma injury in non-survivor patients. <i>BMC Research Notes</i> , 2019, 12, 789.	0.6	6
112	Red blood cell distribution width as mortality biomarker in patients with traumatic brain injury. <i>Acta Neurologica Belgica</i> , 2021, 121, 715-720.	0.5	6
113	High Serum Caspase-Cleaved Cytokeratin-18 Levels and Mortality of Traumatic Brain Injury Patients. <i>Brain Sciences</i> , 2019, 9, 269.	1.1	5
114	Reply to "Strengths and limitations of estimated and measured GFR". <i>Nature Reviews Nephrology</i> , 2019, 15, 785-786.	4.1	5
115	Persistently high circulating tissue inhibitor of matrix metalloproteinase-1 levels in non-survivor brain trauma injury patients. <i>Journal of Critical Care</i> , 2019, 51, 117-121.	1.0	5
116	High Serum Tissue Inhibitor of Matrix Metalloproteinase-1 Levels and Mortality in Patients with Spontaneous Intracerebral Hemorrhage. <i>World Neurosurgery</i> , 2020, 134, e476-e480.	0.7	5
117	Serum substance P levels and early mortality of spontaneous intracerebral haemorrhage patients. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2020, 29, 104893.	0.7	5
118	Risk of liver fibrosis progression in patients with suboptimal diagnosis of hepatitis C virus infection. <i>European Journal of Gastroenterology and Hepatology</i> , 2020, 32, 528-534.	0.8	5
119	Chlorhexidine-silver sulfadiazine-impregnated venous catheters are efficient even at subclavian sites without tracheostomy. <i>American Journal of Infection Control</i> , 2016, 44, 1526-1529.	1.1	4
120	Facial canal dehiscence in patients with cholesteatoma: concordance between intraoperative inspection, computed tomography and neurophysiological findings. <i>European Archives of Oto-Rhino-Laryngology</i> , 2019, 276, 1915-1920.	0.8	4
121	High serum substance P levels and mortality after malignant middle cerebral artery infarction. <i>Journal of Critical Care</i> , 2020, 57, 1-4.	1.0	4
122	High Serum Levels of Caspase-3 and Early Mortality in Patients with Severe Spontaneous Intracerebral Hemorrhage. <i>Neurocritical Care</i> , 2021, 34, 175-181.	1.2	4
123	Association between serum sFasL concentrations and sepsis mortality. <i>Infectious Diseases</i> , 2021, 53, 38-43.	1.4	4
124	Hepatitis C virus media coverage favorably impacts on antibody testing in the non-interferon era. <i>Journal of Public Health</i> , 2021, 43, 385-391.	1.0	4
125	Low blood caspase-8 levels in survivor patients of traumatic brain injury. <i>Neurological Sciences</i> , 2021, 42, 5065-5070.	0.9	4
126	Lower incidence of catheter-related bloodstream infection in cubital than in femoral artery access. <i>Scandinavian Journal of Infectious Diseases</i> , 2011, 43, 814-817.	1.5	3

#	ARTICLE	IF	CITATIONS
127	Lower mitochondrial dysfunction in survivor septic patients with mitochondrial DNA haplogroup JT. <i>Enfermedades Infecciosas Y Microbiología Clínica</i> , 2018, 36, 539-543.	0.3	3
128	Serum total antioxidant capacity prior to liver transplantation for hepatocellular carcinoma is associated with 1-year liver transplantation survival. <i>Journal of International Medical Research</i> , 2018, 46, 2641-2649.	0.4	3
129	Low Serum Melatonin Levels Prior to Liver Transplantation in Patients with Hepatocellular Carcinoma are Associated with Lower Survival after Liver Transplantation. <i>International Journal of Molecular Sciences</i> , 2019, 20, 1696.	1.8	3
130	A Simplified Iohexol-Based Method to Measure Renal Function in Sheep Models of Renal Disease. <i>Biology</i> , 2020, 9, 259.	1.3	3
131	High serum levels of TAC and early mortality in patients with spontaneous intracerebral haemorrhage. <i>Neurological Sciences</i> , 2021, 42, 1491-1497.	0.9	3
132	Patients with high serum substance P levels previously to liver transplantation for hepatocellular carcinoma have higher risk of one-year liver transplantation mortality. <i>Oncotarget</i> , 2018, 9, 21552-21559.	0.8	3
133	Encuesta de adherencia al tratamiento en enfermedad inflamatoria intestinal. Estudio ENADEII. <i>Gastroenterología Y Hepatología</i> , 2020, 43, 285-292.	0.2	3
134	Serum melatonin levels in predicting mortality in patients with severe traumatic brain injury. <i>Anaesthesia, Critical Care & Pain Medicine</i> , 2021, 40, 100966.	0.6	3
135	Moving average as a method of assessing risk of pressure injury using the COMHON index (Conscious) Tj ETQq1 1 0.784314 rgBT /Over <i>Critical Care</i> , 2022, 35, 696-700.	0.6	3
136	Consumption of "gofio", a roasted cereal flour from the Canary Islands, is associated with exercise capacity and risk of coronary heart disease in the elderly. <i>International Journal of Cardiology</i> , 2015, 187, 41-43.	0.8	2
137	Impact of bendopnea on postoperative outcomes in patients with severe aortic stenosis undergoing aortic valve replacement. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2018, 27, 808-812.	0.5	2
138	Non-survivor patients with malignant middle cerebral artery infarction showed persistently high serum malondialdehyde levels. <i>BMC Neurology</i> , 2019, 19, 238.	0.8	2
139	High serum caspase-3 levels in hepatocellular carcinoma prior to liver transplantation and high mortality risk during the first year after liver transplantation. <i>Expert Review of Molecular Diagnostics</i> , 2019, 19, 635-640.	1.5	2
140	Nonsurviving Patients with Severe Traumatic Brain Injury Have Maintained High Serum sCD40L Levels. <i>World Neurosurgery</i> , 2019, 126, e1537-e1541.	0.7	2
141	Higher Serum Melatonin Levels during the First Week of Malignant Middle Cerebral Artery Infarction in Non-Surviving Patients. <i>Brain Sciences</i> , 2019, 9, 346.	1.1	2
142	High Serum Soluble Fas Ligand Levels in Non-survivor Traumatic Brain Injury Patients. <i>Neurocritical Care</i> , 2021, 35, 249-254.	1.2	2
143	An Enhanced High-Volume Preparation for Colonoscopy Is Not Better Than a Conventional Low-Volume One in Patients at Risk of Poor Bowel Cleansing: A Randomized Controlled Trial. <i>Frontiers in Medicine</i> , 2021, 8, 654847.	1.2	2
144	Increasing the low residue diet to 3 days does not improve the bowel cleansing in hard to prepare patients: Post hoc analysis of a randomized controlled trial. <i>Gastroenterología Y Hepatología</i> , 2021, 44, 183-190.	0.2	2

#	ARTICLE	IF	CITATIONS
145	Usefulness of prioritization systems during the resumption of gastrointestinal endoscopy activity during the COVID-19 pandemic. <i>Endoscopy</i> , 2021, 53, 662-663.	1.0	2
146	Association of serum soluble Fas concentrations and mortality of septic patients. <i>Enfermedades Infecciosas Y Microbiología Clínica</i> , 2020, , .	0.3	2
147	Immunogenicity and Acceptance of Influenza A (H1N1) Vaccine in a Cohort of Chronic Hepatitis C Patients Receiving Pegylated-Interferon Treatment. <i>PLoS ONE</i> , 2012, 7, e48610.	1.1	2
148	Mortality of spontaneous intracerebral haemorrhage patients and high serum caspase-8 concentrations. <i>Anaesthesia, Critical Care & Pain Medicine</i> , 2021, 40, 100969.	0.6	2
149	Mortality prediction of patients with spontaneous intracerebral hemorrhage by serum soluble Fas ligand concentrations. <i>Expert Review of Molecular Diagnostics</i> , 2022, 22, 233-238.	1.5	2
150	Association between neutrophil-to-lymphocyte ratio in the first seven days of sepsis and mortality. <i>Enfermedades Infecciosas Y Microbiología Clínica (English Ed)</i> , 2022, 40, 235-240.	0.2	2
151	In Reply to "Predictors of Outcome in Spontaneous Intracerebral Hemorrhage" Role of Oxidative Stress Biomarkers. <i>World Neurosurgery</i> , 2018, 120, 602.	0.7	1
152	High Serum sCD40L Levels During the First Week of Malignant Middle Cerebral Artery Infarction and Mortality. <i>World Neurosurgery</i> , 2019, 132, e630-e636.	0.7	1
153	Serum Caspase-3 Levels and Early Mortality of Patients with Malignant Middle Cerebral Artery Infarction. <i>Neurocritical Care</i> , 2019, 31, 486-493.	1.2	1
154	Delirium in Patients With Heart Failure: Influence of Attendance Times in an Emergency Department. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2019, 72, 82-84.	0.4	1
155	Proyecto educativo de investigación clínica en el programa de formación sanitaria especializada del Hospital Universitario de Canarias: evaluación del impacto a 10 años. <i>Educacion Medica</i> , 2020, 21, 299-305.	0.3	1
156	Patients with suboptimal hepatitis B virus diagnostic characterization are at risk of liver fibrosis progression. <i>European Journal of Gastroenterology and Hepatology</i> , 2020, 32, 426-432.	0.8	1
157	Non-Survivor Ischemic Stroke Patients Maintain High Serum Caspase-Cleaved Cytokeratin-18 Levels. <i>Brain Sciences</i> , 2020, 10, 132.	1.1	1
158	Serum sFasL concentrations and mortality prediction in patients with sepsis. <i>Infectious Diseases</i> , 2021, 53, 643-646.	1.4	1
159	Serum Levels of B-cell Lymphoma-2 Anti-Apoptotic Protein and Malignant Middle Cerebral Artery Infarction Mortality. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2021, 30, 105717.	0.7	1
160	Skin insertion site culture for the prediction of primary bloodstream infection. <i>Irish Journal of Medical Science</i> , 2022, 191, 1269-1275.	0.8	1
161	Mortality prediction by serum melatonin levels of patients with spontaneous intracerebral hemorrhage. <i>Neurological Sciences</i> , 2021, , 1.	0.9	1
162	Circulating Bcl-2 concentrations and septic patient mortality. <i>Enfermedades Infecciosas Y Microbiología Clínica (English Ed)</i> , 2021, 39, 330-334.	0.2	1

#	ARTICLE	IF	CITATIONS
163	Association of serum soluble Fas concentrations and mortality of septic patients. <i>Enfermedades Infecciosas Y Microbiología Clínica (English Ed)</i> , 2021, 39, 493-497.	0.2	1
164	Predictive factors for inadequate bowel cleansing in colon capsule endoscopy. <i>Gastroenterología Y Hepatología</i> , 2022, 45, 605-613.	0.2	1
165	Response to Letter "High serum soluble CD40L levels previously to liver transplantation in patients with hepatocellular carcinoma are associated with mortality at one year: Methodological issues": <i>Journal of Critical Care</i> , 2018, 43, 371.	1.0	0
166	Serum B cell lymphoma-2 concentrations and mortality of patients with spontaneous intracerebral hemorrhage. <i>Neurological Sciences</i> , 2021, 42, 3631-3636.	0.9	0
167	Serum tissue inhibitor of MMP-1 levels at any moment of the first week of spontaneous intracerebral hemorrhage may predict early mortality. <i>Biomarkers in Medicine</i> , 2021, 15, 1245-1251.	0.6	0
168	Mortality prediction of ischemic stroke patients without thrombectomy by blood total antioxidant capacity. <i>Journal of Integrative Neuroscience</i> , 2020, 19, 501.	0.8	0
169	Mortality prediction of septic patients by blood caspase-8 levels. <i>Diagnostic Microbiology and Infectious Disease</i> , 2022, 102, 115639.	0.8	0
170	High mortality rate of septic patients with high blood granzyme B concentrations. <i>Diagnostic Microbiology and Infectious Disease</i> , 2022, 103, 115694.	0.8	0
171	Blood Bcl-2 levels to predict the mortality of septic patients. <i>Biomarkers in Medicine</i> , 2022, 16, 427-433.	0.6	0
172	Association between serum concentrations of anti-apoptotic B-cell lymphoma-2 protein and traumatic brain injury mortality. <i>Expert Review of Molecular Diagnostics</i> , 2022, 22, 125-129.	1.5	0