

Alejandro JimÃ©nez-Sosa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1788270/publications.pdf>

Version: 2024-02-01

172
papers

4,075
citations

136885

32
h-index

149623

56
g-index

178
all docs

178
docs citations

178
times ranked

4899
citing authors

#	ARTICLE	IF	CITATIONS
1	Skin insertion site culture for the prediction of primary bloodstream infection. <i>Irish Journal of Medical Science</i> , 2022, 191, 1269-1275.	0.8	1
2	Mortality prediction of septic patients by blood caspase-8 levels. <i>Diagnostic Microbiology and Infectious Disease</i> , 2022, 102, 115639.	0.8	0
3	Predictive factors for inadequate bowel cleansing in colon capsule endoscopy. <i>Gastroenterology & Hepatology</i> , 2022, 45, 605-613.	0.2	1
4	High mortality rate of septic patients with high blood granzyme B concentrations. <i>Diagnostic Microbiology and Infectious Disease</i> , 2022, 103, 115694.	0.8	0
5	Blood Bcl-2 levels to predict the mortality of septic patients. <i>Biomarkers in Medicine</i> , 2022, 16, 427-433.	0.6	0
6	Moving average as a method of assessing risk of pressure injury using the COMHON index (Conscious) Tj ETQq0 0 0 rgBT /Overlock 10 T Critical Care, 2022, 35, 696-700.	0.6	3
7	Mortality prediction of patients with spontaneous intracerebral hemorrhage by serum soluble Fas ligand concentrations. <i>Expert Review of Molecular Diagnostics</i> , 2022, 22, 233-238.	1.5	2
8	Association between serum concentrations of anti-apoptotic B-cell lymphoma-2 protein and traumatic brain injury mortality. <i>Expert Review of Molecular Diagnostics</i> , 2022, 22, 125-129.	1.5	0
9	Association between neutrophil-to-lymphocyte ratio in the first seven days of sepsis and mortality. <i>Enfermedades Infecciosas Y Microbiología Clínica (English Ed)</i> , 2022, 40, 235-240.	0.2	2
10	High Serum Levels of Caspase-3 and Early Mortality in Patients with Severe Spontaneous Intracerebral Hemorrhage. <i>Neurocritical Care</i> , 2021, 34, 175-181.	1.2	4
11	Red blood cell distribution width as mortality biomarker in patients with traumatic brain injury. <i>Acta Neurologica Belgica</i> , 2021, 121, 715-720.	0.5	6
12	Association between red blood cell distribution width and mortality of COVID-19 patients. <i>Anaesthesia, Critical Care & Pain Medicine</i> , 2021, 40, 100777.	0.6	36
13	Proprotein convertase subtilisin/kexin type 9 in the dyslipidaemia of patients with axial spondyloarthritis is related to disease activity. <i>Rheumatology</i> , 2021, 60, 2296-2306.	0.9	10
14	Association between serum sFasL concentrations and sepsis mortality. <i>Infectious Diseases</i> , 2021, 53, 38-43.	1.4	4
15	High serum levels of TAC and early mortality in patients with spontaneous intracerebral haemorrhage. <i>Neurological Sciences</i> , 2021, 42, 1491-1497.	0.9	3
16	Hepatitis C virus media coverage favorably impacts on antibody testing in the non-interferon era. <i>Journal of Public Health</i> , 2021, 43, 385-391.	1.0	4
17	Serum B cell lymphoma-2 concentrations and mortality of patients with spontaneous intracerebral hemorrhage. <i>Neurological Sciences</i> , 2021, 42, 3631-3636.	0.9	0
18	High Serum Soluble Fas Ligand Levels in Non-survivor Traumatic Brain Injury Patients. <i>Neurocritical Care</i> , 2021, 35, 249-254.	1.2	2

#	ARTICLE	IF	CITATIONS
19	Extracellular water/total body water ratio as predictor of mortality in hemodialysis patients. <i>Renal Failure</i> , 2021, 43, 821-829.	0.8	11
20	Serum sFasL concentrations and mortality prediction in patients with sepsis. <i>Infectious Diseases</i> , 2021, 53, 643-646.	1.4	1
21	An Enhanced High-Volume Preparation for Colonoscopy Is Not Better Than a Conventional Low-Volume One in Patients at Risk of Poor Bowel Cleansing: A Randomized Controlled Trial. <i>Frontiers in Medicine</i> , 2021, 8, 654847.	1.2	2
22	Increasing the low residue diet to 3 days does not improve the bowel cleansing in hard to prepare patients: Post hoc analysis of a randomized controlled trial. <i>Gastroenterología Y Hepatología</i> , 2021, 44, 183-190.	0.2	2
23	Low blood caspase-8 levels in survivor patients of traumatic brain injury. <i>Neurological Sciences</i> , 2021, 42, 5065-5070.	0.9	4
24	Usefulness of prioritization systems during the resumption of gastrointestinal endoscopy activity during the COVID-19 pandemic. <i>Endoscopy</i> , 2021, 53, 662-663.	1.0	2
25	Serum Levels of B-cell Lymphoma-2 Anti-Apoptotic Protein and Malignant Middle Cerebral Artery Infarction Mortality. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2021, 30, 105717.	0.7	1
26	DNA and RNA Oxidative Damage and Mortality of Patients With COVID-19. <i>American Journal of the Medical Sciences</i> , 2021, 361, 585-590.	0.4	15
27	Blood concentrations of proapoptotic sFas and antiapoptotic Bcl2 and COVID-19 patient mortality. <i>Expert Review of Molecular Diagnostics</i> , 2021, 21, 837-844.	1.5	8
28	Mortality prediction by serum melatonin levels of patients with spontaneous intracerebral hemorrhage. <i>Neurological Sciences</i> , 2021, , 1.	0.9	1
29	Circulating Bcl-2 concentrations and septic patient mortality. <i>Enfermedades Infecciosas Y Microbiología Clínica (English Ed)</i> , 2021, 39, 330-334.	0.2	1
30	Serum tissue inhibitor of MMP-1 levels at any moment of the first week of spontaneous intracerebral hemorrhage may predict early mortality. <i>Biomarkers in Medicine</i> , 2021, 15, 1245-1251.	0.6	0
31	Serum melatonin levels in predicting mortality in patients with severe traumatic brain injury. <i>Anaesthesia, Critical Care & Pain Medicine</i> , 2021, 40, 100966.	0.6	3
32	Mortality of spontaneous intracerebral haemorrhage patients and high serum caspase-8 concentrations. <i>Anaesthesia, Critical Care & Pain Medicine</i> , 2021, 40, 100969.	0.6	2
33	Association of serum soluble Fas concentrations and mortality of septic patients. <i>Enfermedades Infecciosas Y Microbiología Clínica (English Ed)</i> , 2021, 39, 493-497.	0.2	1
34	Proyecto educativo de investigación clínica en el programa de formación sanitaria especializada del Hospital Universitario de Canarias: evaluación del impacto a 10 años. <i>Educación Médica</i> , 2020, 21, 299-305.	0.3	1
35	Association Between DNA and RNA Oxidative Damage and Mortality of Patients with Traumatic Brain Injury. <i>Neurocritical Care</i> , 2020, 32, 790-795.	1.2	11
36	High Serum Tissue Inhibitor of Matrix Metalloproteinase-1 Levels and Mortality in Patients with Spontaneous Intracerebral Hemorrhage. <i>World Neurosurgery</i> , 2020, 134, e476-e480.	0.7	5

#	ARTICLE	IF	CITATIONS
37	High Serum DNA and RNA Oxidative Damage in Non-surviving Patients with Spontaneous Intracerebral Hemorrhage. <i>Neurocritical Care</i> , 2020, 33, 90-96.	1.2	9
38	Patients with suboptimal hepatitis B virus diagnostic characterization are at risk of liver fibrosis progression. <i>European Journal of Gastroenterology and Hepatology</i> , 2020, 32, 426-432.	0.8	1
39	A Simplified Iohexol-Based Method to Measure Renal Function in Sheep Models of Renal Disease. <i>Biology</i> , 2020, 9, 259.	1.3	3
40	Early Mortality of Brain Infarction Patients and Red Blood Cell Distribution Width. <i>Brain Sciences</i> , 2020, 10, 196.	1.1	14
41	Non-Survivor Ischemic Stroke Patients Maintain High Serum Caspase-Cleaved Cytokeratin-18 Levels. <i>Brain Sciences</i> , 2020, 10, 132.	1.1	1
42	Serum substance P levels and early mortality of spontaneous intracerebral haemorrhage patients. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2020, 29, 104893.	0.7	5
43	Red blood cell distribution width and mortality of spontaneous intracerebral hemorrhage patients. <i>Clinical Neurology and Neurosurgery</i> , 2020, 195, 106066.	0.6	11
44	Risk of liver fibrosis progression in patients with suboptimal diagnosis of hepatitis C virus infection. <i>European Journal of Gastroenterology and Hepatology</i> , 2020, 32, 528-534.	0.8	5
45	Impaired HDL cholesterol efflux capacity in systemic lupus erythematosus patients is related to subclinical carotid atherosclerosis. <i>Rheumatology</i> , 2020, 59, 2847-2856.	0.9	30
46	Traumatic Brain Injury Patients Mortality and Serum Total Antioxidant Capacity. <i>Brain Sciences</i> , 2020, 10, 110.	1.1	7
47	High serum substance P levels and mortality after malignant middle cerebral artery infarction. <i>Journal of Critical Care</i> , 2020, 57, 1-4.	1.0	4
48	Association of serum soluble Fas concentrations and mortality of septic patients. <i>Enfermedades Infecciosas Y Microbiología Clínica</i> , 2020, , .	0.3	2
49	Encuesta de adherencia al tratamiento en enfermedad inflamatoria intestinal. Estudio ENADEII. <i>Gastroenterología Y Hepatología</i> , 2020, 43, 285-292.	0.2	3
50	Association between neutrophil-to-lymphocyte ratio in the first seven days of sepsis and mortality. <i>Enfermedades Infecciosas Y Microbiología Clínica</i> , 2020, , .	0.3	7
51	Mortality prediction of ischemic stroke patients without thrombectomy by blood total antioxidant capacity. <i>Journal of Integrative Neuroscience</i> , 2020, 19, 501.	0.8	0
52	Chronic kidney disease staging with cystatin C or creatinine-based formulas: flipping the coin. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, 287-294.	0.4	22
53	Association between DNA and RNA oxidative damage and mortality in septic patients. <i>Journal of Critical Care</i> , 2019, 54, 94-98.	1.0	10
54	High serum levels of tissue inhibitor of matrix metalloproteinase-1 during the first week of a malignant middle cerebral artery infarction in non-surviving patients. <i>BMC Neurology</i> , 2019, 19, 167.	0.8	6

#	ARTICLE	IF	CITATIONS
55	Persistently High Serum Substance P Levels and Early Mortality in Patients with Severe Traumatic Brain Injury. <i>World Neurosurgery</i> , 2019, 132, e613-e617.	0.7	13
56	High Serum sCD40L Levels During the First Week of Malignant Middle Cerebral Artery Infarction and Mortality. <i>World Neurosurgery</i> , 2019, 132, e630-e636.	0.7	1
57	High Serum Caspase-Cleaved Cytokeratin-18 Levels and Mortality of Traumatic Brain Injury Patients. <i>Brain Sciences</i> , 2019, 9, 269.	1.1	5
58	Non-survivor patients with malignant middle cerebral artery infarction showed persistently high serum malondialdehyde levels. <i>BMC Neurology</i> , 2019, 19, 238.	0.8	2
59	The Serum Melatonin Levels and Mortality of Patients with Spontaneous Intracerebral Hemorrhage. <i>Brain Sciences</i> , 2019, 9, 263.	1.1	11
60	The Error of Estimated GFR in Type 2 Diabetes Mellitus. <i>Journal of Clinical Medicine</i> , 2019, 8, 1543.	1.0	15
61	Reply to "Strengths and limitations of estimated and measured GFR". <i>Nature Reviews Nephrology</i> , 2019, 15, 785-786.	4.1	5
62	Advanced interatrial block is a surrogate for left atrial strain reduction which predicts atrial fibrillation and stroke. <i>Annals of Noninvasive Electrocardiology</i> , 2019, 24, e12632.	0.5	29
63	Serum Caspase-3 Levels and Early Mortality of Patients with Malignant Middle Cerebral Artery Infarction. <i>Neurocritical Care</i> , 2019, 31, 486-493.	1.2	1
64	High serum caspase-3 levels in hepatocellular carcinoma prior to liver transplantation and high mortality risk during the first year after liver transplantation. <i>Expert Review of Molecular Diagnostics</i> , 2019, 19, 635-640.	1.5	2
65	Nonsurviving Patients with Severe Traumatic Brain Injury Have Maintained High Serum sCD40L Levels. <i>World Neurosurgery</i> , 2019, 126, e1537-e1541.	0.7	2
66	Impact of errors of creatinine and cystatin C equations in the selection of living kidney donors. <i>CKJ: Clinical Kidney Journal</i> , 2019, 12, 748-755.	1.4	14
67	Low Serum Melatonin Levels Prior to Liver Transplantation in Patients with Hepatocellular Carcinoma are Associated with Lower Survival after Liver Transplantation. <i>International Journal of Molecular Sciences</i> , 2019, 20, 1696.	1.8	3
68	Facial canal dehiscence in patients with cholesteatoma: concordance between intraoperative inspection, computed tomography and neurophysiological findings. <i>European Archives of Oto-Rhino-Laryngology</i> , 2019, 276, 1915-1920.	0.8	4
69	Impact of a 1-day versus 3-day low-residue diet on bowel cleansing quality before colonoscopy: a randomized controlled trial. <i>Endoscopy</i> , 2019, 51, 628-636.	1.0	40
70	Persistently high circulating tissue inhibitor of matrix metalloproteinase-1 levels in non-survivor brain trauma injury patients. <i>Journal of Critical Care</i> , 2019, 51, 117-121.	1.0	5
71	Maintained high sustained serum malondialdehyde levels after severe brain trauma injury in non-survivor patients. <i>BMC Research Notes</i> , 2019, 12, 789.	0.6	6
72	Higher Serum Melatonin Levels during the First Week of Malignant Middle Cerebral Artery Infarction in Non-Surviving Patients. <i>Brain Sciences</i> , 2019, 9, 346.	1.1	2

#	ARTICLE	IF	CITATIONS
73	Estimated GFR: time for a critical appraisal. <i>Nature Reviews Nephrology</i> , 2019, 15, 177-190.	4.1	187
74	Relationship Between Insulin Sensitivity and β -Cell Secretion in Nondiabetic Subjects with Rheumatoid Arthritis. <i>Journal of Rheumatology</i> , 2019, 46, 229-236.	1.0	10
75	Delirium in Patients With Heart Failure: Influence of Attendance Times in an Emergency Department. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2019, 72, 82-84.	0.4	1
76	Iohexol plasma clearance simplified by dried blood spot testing. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, 1597-1603.	0.4	29
77	Serum Malondialdehyde Levels and Mortality in Patients with Spontaneous Intracerebral Hemorrhage. <i>World Neurosurgery</i> , 2018, 113, e542-e547.	0.7	17
78	Serum melatonin levels during the first seven days of severe sepsis diagnosis are associated with sepsis severity and mortality. <i>Enfermedades Infecciosas Y Microbiología Clínica</i> , 2018, 36, 544-549.	0.3	9
79	Association between serum levels of caspase-cleaved cytokeratin-18 and early mortality in patients with severe spontaneous intracerebral hemorrhage. <i>BMC Neuroscience</i> , 2018, 19, 23.	0.8	6
80	Lower mitochondrial dysfunction in survivor septic patients with mitochondrial DNA haplogroup JT. <i>Enfermedades Infecciosas Y Microbiología Clínica</i> , 2018, 36, 539-543.	0.3	3
81	Response to Letter "High serum soluble CD40L levels previously to liver transplantation in patients with hepatocellular carcinoma are associated with mortality at one year: Methodological issues". <i>Journal of Critical Care</i> , 2018, 43, 371.	1.0	0
82	High serum soluble CD40L levels previously to liver transplantation in patients with hepatocellular carcinoma are associated with mortality at one year. <i>Journal of Critical Care</i> , 2018, 43, 316-320.	1.0	7
83	Predictive ability of the EVARUCI scale and COMHON index for pressure injury risk in critically ill patients: A diagnostic accuracy study. <i>Australian Critical Care</i> , 2018, 31, 355-361.	0.6	9
84	In Reply to "Predictors of Outcome in Spontaneous Intracerebral Hemorrhage" Role of Oxidative Stress Biomarkers. <i>World Neurosurgery</i> , 2018, 120, 602.	0.7	1
85	Serum total antioxidant capacity during the first week of sepsis and mortality. <i>Journal of Critical Care</i> , 2018, 47, 139-144.	1.0	15
86	A Simple Method to Measure Renal Function in Swine by the Plasma Clearance of Iohexol. <i>International Journal of Molecular Sciences</i> , 2018, 19, 232.	1.8	10
87	High serum levels of caspase-cleaved cytokeratin-18 are associated with malignant middle cerebral artery infarction patient mortality. <i>BMC Neurology</i> , 2018, 18, 32.	0.8	10
88	Interatrial block and atrial remodeling assessed using speckle tracking echocardiography. <i>BMC Cardiovascular Disorders</i> , 2018, 18, 38.	0.7	36
89	Serum melatonin levels are associated with mortality in patients with malignant middle cerebral artery infarction. <i>Journal of International Medical Research</i> , 2018, 46, 3268-3277.	0.4	14
90	Impact of bendopnea on postoperative outcomes in patients with severe aortic stenosis undergoing aortic valve replacement. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2018, 27, 808-812.	0.5	2

#	ARTICLE	IF	CITATIONS
91	Serum total antioxidant capacity prior to liver transplantation for hepatocellular carcinoma is associated with 1-year liver transplantation survival. <i>Journal of International Medical Research</i> , 2018, 46, 2641-2649.	0.4	3
92	Patients with high serum substance P levels previously to liver transplantation for hepatocellular carcinoma have higher risk of one-year liver transplantation mortality. <i>Oncotarget</i> , 2018, 9, 21552-21559.	0.8	3
93	Non-survivor septic patients have persistently higher serum sCD40L levels than survivors. <i>Journal of Critical Care</i> , 2017, 41, 177-182.	1.0	11
94	Comparison of Two Intensive Bowel Cleansing Regimens in Patients With Previous Poor Bowel Preparation: A Randomized Controlled Study. <i>American Journal of Gastroenterology</i> , 2017, 112, 951-958.	0.2	51
95	Sustained Low Serum Substance P Levels in Non-Surviving Septic Patients. <i>International Journal of Molecular Sciences</i> , 2017, 18, 1531.	1.8	7
96	Serum melatonin levels in survivor and non-survivor patients with traumatic brain injury. <i>BMC Neurology</i> , 2017, 17, 138.	0.8	17
97	Set of Quality Indicators of Pediatric Intensive Care in Spain: Delphi Method Selection. <i>Pediatric Quality & Safety</i> , 2017, 2, e009.	0.4	9
98	Association between Pre-Transplant Serum Malondialdehyde Levels and Survival One Year after Liver Transplantation for Hepatocellular Carcinoma. <i>International Journal of Molecular Sciences</i> , 2016, 17, 500.	1.8	23
99	Prognostic Value of Serum Caspase-Cleaved Cytokeratin-18 Levels before Liver Transplantation for One-Year Survival of Patients with Hepatocellular Carcinoma. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1524.	1.8	16
100	Association between Interleukin-6 Promoter Polymorphism (-174 G/C), Serum Interleukin-6 Levels and Mortality in Severe Septic Patients. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1861.	1.8	26
101	Serum caspase 3 levels are associated with early mortality in severe septic patients. <i>Journal of Critical Care</i> , 2016, 34, 103-106.	1.0	14
102	Chlorhexidine-silver sulfadiazine-impregnated venous catheters are efficient even at subclavian sites without tracheostomy. <i>American Journal of Infection Control</i> , 2016, 44, 1526-1529.	1.1	4
103	Association between total antioxidant capacity and mortality in ischemic stroke patients. <i>Annals of Intensive Care</i> , 2016, 6, 39.	2.2	30
104	Plasma matrix metalloproteinase 9 as an early surrogate biomarker of advanced colorectal neoplasia. <i>Gastroenterology & Hepatology</i> , 2016, 39, 433-441.	0.2	11
105	Association Between Bendopnea and Key Parameters of Cardiopulmonary Exercise Testing in Patients With Advanced Heart Failure. <i>Journal of Cardiac Failure</i> , 2016, 22, 163-165.	0.7	16
106	Septic patients with mitochondrial DNA haplogroup JT have higher respiratory complex IV activity and survival rate. <i>Journal of Critical Care</i> , 2016, 33, 95-99.	1.0	8
107	Chlorhexidine-silver sulfadiazine or rifampicin-miconazole impregnated venous catheters decrease the risk of catheter-related bloodstream infection similarly. <i>American Journal of Infection Control</i> , 2016, 44, 50-53.	1.1	18
108	Serum caspase-3 levels and mortality are associated in patients with severe traumatic brain injury. <i>BMC Neurology</i> , 2015, 15, 228.	0.8	34

#	ARTICLE	IF	CITATIONS
109	Serum substance P levels are associated with severity and mortality in patients with severe traumatic brain injury. <i>Critical Care</i> , 2015, 19, 192.	2.5	41
110	Levosimendan Improves Hemodynamic Status in Critically Ill Patients with Severe Aortic Stenosis and Left Ventricular Dysfunction: An Interventional Study. <i>Cardiovascular Therapeutics</i> , 2015, 33, 193-199.	1.1	13
111	Association between Serum Soluble CD154 Levels and Mortality in Patients with Malignant Middle Cerebral Artery Infarction. <i>International Journal of Molecular Sciences</i> , 2015, 16, 12147-12158.	1.8	12
112	The 4G/4G Genotype of PAI-1 Polymorphism Is Associated with Higher Plasma PAI-1 Concentrations and Mortality in Patients with Severe Sepsis. <i>PLoS ONE</i> , 2015, 10, e0129565.	1.1	14
113	Serum Malondialdehyde Levels in Patients with Malignant Middle Cerebral Artery Infarction Are Associated with Mortality. <i>PLoS ONE</i> , 2015, 10, e0125893.	1.1	33
114	The impact of frailty in older patients with non-ischaemic cardiomyopathy after implantation of cardiac resynchronization therapy defibrillator. <i>Europace</i> , 2015, 17, 598-602.	0.7	47
115	Consumption of "gofio", a roasted cereal flour from the Canary Islands, is associated with exercise capacity and risk of coronary heart disease in the elderly. <i>International Journal of Cardiology</i> , 2015, 187, 41-43.	0.8	2
116	Total antioxidant capacity is associated with mortality of patients with severe traumatic brain injury. <i>BMC Neurology</i> , 2015, 15, 115.	0.8	20
117	Efficiency of chlorhexidine-silver sulfadiazine-impregnated venous catheters at subclavian sites. <i>American Journal of Infection Control</i> , 2015, 43, 711-714.	1.1	14
118	Serum tissue inhibitor of matrix metalloproteinase-1 levels are associated with mortality in patients with malignant middle cerebral artery infarction. <i>BMC Neurology</i> , 2015, 15, 111.	0.8	11
119	Decrease of oxidative phosphorylation system function in severe septic patients. <i>Journal of Critical Care</i> , 2015, 30, 935-939.	1.0	25
120	Serum melatonin levels are associated with mortality in severe septic patients. <i>Journal of Critical Care</i> , 2015, 30, 860.e1-860.e6.	1.0	18
121	Association between serum substance P levels and mortality in patients with severe sepsis. <i>Journal of Critical Care</i> , 2015, 30, 924-928.	1.0	20
122	Association between serum total antioxidant capacity and mortality in severe septic patients. <i>Journal of Critical Care</i> , 2015, 30, 217.e7-217.e12.	1.0	28
123	Serum Levels of Caspase-Cleaved Cytokeratin-18 in Patients with Severe Traumatic Brain Injury Are Associated with Mortality: A Pilot Study. <i>PLoS ONE</i> , 2015, 10, e0121739.	1.1	29
124	Association of Sepsis-Related Mortality with Early Increase of TIMP-1/MMP-9 Ratio. <i>PLoS ONE</i> , 2014, 9, e94318.	1.1	60
125	Association between Serum Tissue Inhibitor of Matrix Metalloproteinase-1 Levels and Mortality in Patients with Severe Brain Trauma Injury. <i>PLoS ONE</i> , 2014, 9, e94370.	1.1	34
126	Red Blood Cell Distribution Width during the First Week Is Associated with Severity and Mortality in Septic Patients. <i>PLoS ONE</i> , 2014, 9, e105436.	1.1	68

#	ARTICLE	IF	CITATIONS
127	Serum Levels of Caspase-Cleaved Cytokeratin-18 and Mortality Are Associated in Severe Septic Patients: Pilot Study. PLoS ONE, 2014, 9, e109618.	1.1	28
128	Cost/benefit analysis of chlorhexidine-silver sulfadiazine-impregnated venous catheters for femoral access. American Journal of Infection Control, 2014, 42, 1130-1132.	1.1	10
129	Subglottic secretion drainage and continuous control of cuff pressure used together save health care costs. American Journal of Infection Control, 2014, 42, 1101-1105.	1.1	14
130	Chlorhexidine-silver sulfadiazine-impregnated venous catheters save costs. American Journal of Infection Control, 2014, 42, 321-324.	1.1	24
131	Serum soluble CD40 Ligand levels are associated with severity and mortality of brain trauma injury patients. Thrombosis Research, 2014, 134, 832-836.	0.8	23
132	One-day versus two-day cleansing for colon capsule endoscopy: a prospective randomized pilot study. Gastroenterology and Hepatology, 2014, 37, 101-106.	0.2	15
133	Sustained high plasma plasminogen activator inhibitor-1 levels are associated with severity and mortality in septic patients. Thrombosis Research, 2014, 134, 182-186.	0.8	31
134	Measurement of glomerular filtration rate: Internal and external validations of the iohexol plasma clearance technique by HPLC. Clinica Chimica Acta, 2014, 430, 84-85.	0.5	16
135	The 372 T/C genetic polymorphism of TIMP-1 is associated with serum levels of TIMP-1 and survival in patients with severe sepsis. Critical Care, 2013, 17, R94.	2.5	31
136	Effects of Colon Capsule Endoscopy on Medical Decision Making in Patients With Incomplete Colonoscopies. Clinical Gastroenterology and Hepatology, 2013, 11, 534-540.e1.	2.4	64
137	Prognostic Value of Malondialdehyde Serum Levels in Severe Sepsis: A Multicenter Study. PLoS ONE, 2013, 8, e53741.	1.1	49
138	Severe Septic Patients with Mitochondrial DNA Haplogroup JT Show Higher Survival Rates: A Prospective, Multicenter, Observational Study. PLoS ONE, 2013, 8, e73320.	1.1	17
139	Survival and mitochondrial function in septic patients according to mitochondrial DNA haplogroup. Critical Care, 2012, 16, R10.	2.5	22
140	Immunogenicity and Acceptance of Influenza A (H1N1) Vaccine in a Cohort of Chronic Hepatitis C Patients Receiving Pegylated-Interferon Treatment. PLoS ONE, 2012, 7, e48610.	1.1	2
141	Lower associated costs using rifampicin-miconazole-impregnated catheters compared with standard catheters. American Journal of Infection Control, 2011, 39, 895-897.	1.1	16
142	Association between serum soluble CD40 ligand levels and mortality in patients with severe sepsis. Critical Care, 2011, 15, R97.	2.5	53
143	Low-Dose, Low-Concentration Levobupivacaine Plus Fentanyl Selective Spinal Anesthesia for Knee Arthroscopy. Anesthesia and Analgesia, 2011, 112, 477-480.	1.1	13
144	Platelet cytochrome c oxidase activity and quantity in septic patients*. Critical Care Medicine, 2011, 39, 1289-1294.	0.4	30

#	ARTICLE	IF	CITATIONS
145	Lower incidence of catheter-related bloodstream infection in cubital than in femoral artery access. <i>Scandinavian Journal of Infectious Diseases</i> , 2011, 43, 814-817.	1.5	3
146	Endoscopic submucosal dissection training with pig models in a Western country. <i>World Journal of Gastroenterology</i> , 2010, 16, 2895.	1.4	88
147	Diagnostic accuracy of immunochemical versus guaiac faecal occult blood tests for colorectal cancer screening. <i>Journal of Gastroenterology</i> , 2010, 45, 703-712.	2.3	105
148	Lower arterial catheter-related infection in brachial than in femoral access. <i>American Journal of Infection Control</i> , 2010, 38, e40-e42.	1.1	6
149	Matrix metalloproteinase-9, -10, and tissue inhibitor of matrix metalloproteinases-1 blood levels as biomarkers of severity and mortality in sepsis. <i>Critical Care</i> , 2009, 13, R158.	2.5	105
150	Colorectal cancer screening in a Spanish population. <i>Medicina Clínica</i> , 2009, 133, 736-740.	0.3	12
151	Clinical cure of ventilator-associated pneumonia treated with piperacillin/tazobactam administered by continuous or intermittent infusion. <i>International Journal of Antimicrobial Agents</i> , 2009, 33, 464-468.	1.1	120
152	Low-Dose 3 mg Levobupivacaine Plus 10 µg Fentanyl Selective Spinal Anesthesia for Gynecological Outpatient Laparoscopy. <i>Anesthesia and Analgesia</i> , 2009, 109, 1456-1461.	1.1	22
153	Validation of Fujinon intelligent chromoendoscopy with high definition endoscopes in colonoscopy. <i>World Journal of Gastroenterology</i> , 2009, 15, 5266.	1.4	34
154	Factors influencing pregnancy outcome in women with type 2 versus type 1 diabetes mellitus. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2008, 87, 43-49.	1.3	40
155	The Use of Rifampicin-Miconazole Impregnated Catheters Reduces the Incidence of Femoral and Jugular Catheter-Related Bacteremia. <i>Clinical Infectious Diseases</i> , 2008, 47, 1171-1175.	2.9	46
156	Influence of an Endotracheal Tube with Polyurethane Cuff and Subglottic Secretion Drainage on Pneumonia. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2007, 176, 1079-1083.	2.5	242
157	Microorganisms responsible for intravascular catheter-related bloodstream infection according to the catheter site*. <i>Critical Care Medicine</i> , 2007, 35, 2424-2427.	0.4	119
158	Comparison of clinical cure rates in adults with ventilator-associated pneumonia treated with intravenous ceftazidime administered by continuous or intermittent infusion: A retrospective, nonrandomized, open-label, historical chart review. <i>Clinical Therapeutics</i> , 2007, 29, 2433-2439.	1.1	73
159	Internal jugular venous catheter-related bacteremia according to central and posterior accesses. <i>Intensive Care Medicine</i> , 2007, 33, 1071-1075.	3.9	18
160	Risk for high-grade dysplasia or invasive carcinoma in colorectal flat adenomas in a Spanish population. <i>Gastroenterología Y Hepatología</i> , 2006, 29, 602-609.	0.2	26
161	Meropenem by Continuous Versus Intermittent Infusion in Ventilator-Associated Pneumonia due to Gram-Negative Bacilli. <i>Annals of Pharmacotherapy</i> , 2006, 40, 219-223.	0.9	168
162	Arterial catheter-related infection of 2,949 catheters. <i>Critical Care</i> , 2006, 10, R83.	2.5	60

#	ARTICLE	IF	CITATIONS
163	Ventilator-associated pneumonia using a heated humidifier or a heat and moisture exchanger: a randomized controlled trial [ISRCTN88724583]. <i>Critical Care</i> , 2006, 10, R116.	2.5	55
164	New founding mutation in MSH2 associated with hereditary nonpolyposis colorectal cancer syndrome on the Island of Tenerife. <i>Cancer Letters</i> , 2006, 244, 268-273.	3.2	19
165	Up-regulation of gelatinases in the colorectal adenoma-carcinoma sequence. <i>European Journal of Cancer</i> , 2006, 42, 3246-3252.	1.3	33
166	Tracheal suction by closed system without daily change versus open system. <i>Intensive Care Medicine</i> , 2006, 32, 538-544.	3.9	68
167	Central venous catheter-related infection in a prospective and observational study of 2,595 catheters. <i>Critical Care</i> , 2005, 9, R631.	2.5	257
168	Catheter-related infection in critically ill patients. <i>Intensive Care Medicine</i> , 2004, 30, 1681-4.	3.9	58
169	Accidental catheter removal in critically ill patients: a prospective and observational study. <i>Critical Care</i> , 2004, 8, R229.	2.5	55
170	High prevalence of overweight in a stable spanish hemodialysis population: A cross sectional study. , 2003, 13, 52-59.		17
171	Impact of physician specialty on the cost of nonvariceal upper GI bleeding care. <i>American Journal of Gastroenterology</i> , 2002, 97, 1535-1542.	0.2	24
172	Relationship between serum magnesium and parathyroid hormone levels in hemodialysis patients. <i>American Journal of Kidney Diseases</i> , 1999, 34, 43-48.	2.1	93