

He-Gen Zheng

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1803469/he-gen-zheng-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

143
papers

6,240
citations

44
h-index

74
g-index

146
ext. papers

7,028
ext. citations

5
avg, IF

6.34
L-index

#	Paper	IF	Citations
143	Novel MOF-derived hollow CoFe alloy coupled with N-doped Ketjen Black as boosted bifunctional oxygen catalysts for Zn air batteries. <i>Chemical Engineering Journal</i> , 2022 , 427, 131614	14.7	7
142	Energetic MOF-derived hollow carbon tubes with interconnected channels and encapsulated nickel-cobalt alloy sites as bifunctional catalysts for Zn air batteries with stable cycling over 600 cycles. <i>Applied Surface Science</i> , 2022 , 591, 153070	6.7	0
141	A Water-Stable Tb-MOF As a Rapid, Accurate, and Highly Sensitive Ratiometric Luminescent Sensor for the Discriminative Sensing of Antibiotics and DO in HO. <i>Inorganic Chemistry</i> , 2021 , 60, 10513-10521	5.1	16
140	MOF-derived Co-MOF,O-doped carbon as trifunctional electrocatalysts to enable highly efficient Zn air batteries and water-splitting. <i>Journal of Energy Chemistry</i> , 2021 , 56, 290-298	12	41
139	Fluorescence recognition of adenosine triphosphate and uric acid by two Eu-based metal organic frameworks. <i>Journal of Materials Chemistry C</i> , 2021 , 9, 6051-6061	7.1	13
138	Mixed matrix membranes containing fluorescent coordination polymers for detecting CrO with high sensitivity, stability and recyclability. <i>Dalton Transactions</i> , 2021 , 50, 7944-7948	4.3	5
137	Molecular engineering in a family of pillared-layered metal-organic frameworks for tuning gas adsorption behavior. <i>Dalton Transactions</i> , 2021 , 50, 7409-7416	4.3	1
136	Energetic MOF-derived cobalt/iron nitrides embedded into N, S-codoped carbon nanotubes as superior bifunctional oxygen catalysts for Zn air batteries. <i>Applied Surface Science</i> , 2021 , 569, 151030	6.7	1
135	The difference in the CO adsorption capacities of different functionalized pillar-layered metal-organic frameworks (MOFs). <i>Dalton Transactions</i> , 2021 , 50, 9310-9316	4.3	2
134	MOF-derived CoNi,CoO,NiO@N-C bifunctional oxygen electrocatalysts for liquid and all-solid-state Zn-air batteries. <i>Nanoscale</i> , 2021 , 13, 17655-17662	7.7	3
133	Improving the Stability and Visualizing the Structural Transformation of the Stimuli-Responsive Metal-Organic Frameworks (MOFs). <i>Inorganic Chemistry</i> , 2020 , 59, 5093-5098	5.1	4
132	Bifunctional electrocatalysts for Zn air batteries: recent developments and future perspectives. <i>Journal of Materials Chemistry A</i> , 2020 , 8, 6144-6182	13	81
131	Three metal-organic framework isomers of different pore sizes for selective CO adsorption and isomerization studies. <i>Dalton Transactions</i> , 2020 , 49, 5618-5624	4.3	7
130	Syntheses, crystal structures, dye degradation and luminescence sensing properties of four coordination polymers. <i>CrystEngComm</i> , 2020 , 22, 2327-2335	3.3	16
129	A Europium-based MOF Fluorescent Probe for Efficiently Detecting Malachite Green and Uric Acid. <i>Inorganic Chemistry</i> , 2020 , 59, 7181-7187	5.1	49
128	Four New Luminescent Metal Organic Frameworks as Multifunctional Sensors for Detecting Fe ³⁺ , Cr ^{2O7²⁻} and Nitromethane. <i>Crystal Growth and Design</i> , 2020 , 20, 1898-1904	3.5	24
127	Trimetal-based N-doped carbon nanotubes arrays on Ni foams as self-supported electrodes for hydrogen/oxygen evolution reactions and water splitting. <i>Journal of Power Sources</i> , 2020 , 480, 228866	8.9	22

126	MOF-derived Fe,Co@NC bifunctional oxygen electrocatalysts for Zn-air batteries. <i>Journal of Materials Chemistry A</i> , 2020 , 8, 9355-9363	13	77
125	An excellent example illustrating the fluorescence sensing property of cobalt-organic frameworks. <i>Dalton Transactions</i> , 2019 , 48, 2285-2289	4-3	14
124	Three Anionic Indium-Organic Frameworks for Highly Efficient and Selective Dye Adsorption, Lanthanide Adsorption, and Luminescence Regulation. <i>Inorganic Chemistry</i> , 2019 , 58, 8396-8407	5-1	29
123	Effective adsorption of Congo red by a MOF-based magnetic material. <i>Dalton Transactions</i> , 2019 , 48, 4650-4656	4-3	61
122	Two bifunctional photoluminescent Zn (II) coordination polymers for detection of Fe ³⁺ ion and nitrobenzene. <i>Inorganic Chemistry Communication</i> , 2019 , 107, 107479	3-1	6
121	Three Cd(II) MOFs with Different Functional Groups: Selective CO Capture and Metal Ions Detection. <i>Inorganic Chemistry</i> , 2018 , 57, 5232-5239	5-1	59
120	A Highly Solvent-Stable Metal-Organic Framework Nanosheet: Morphology Control, Exfoliation, and Luminescent Property. <i>Small</i> , 2018 , 14, e1703873	11	59
119	Novel MOF-Derived Co@N-C Bifunctional Catalysts for Highly Efficient Zn-Air Batteries and Water Splitting. <i>Advanced Materials</i> , 2018 , 30, 1705431	24	514
118	A bifunctional photoluminescent metal-organic framework for detection of Fe ³⁺ ion and nitroaromatics. <i>Inorganic Chemistry Communication</i> , 2018 , 89, 68-72	3-1	14
117	Three Zn(ii)-based MOFs for luminescence sensing of Fe and CrO ions. <i>Dalton Transactions</i> , 2018 , 47, 3298-3302	4-3	43
116	A triphenylamine-functionalized luminescent sensor for efficient p-nitroaniline detection. <i>Dalton Transactions</i> , 2018 , 47, 7222-7228	4-3	33
115	Selective separation of methyl orange from water using magnetic ZIF-67 composites. <i>Chemical Engineering Journal</i> , 2018 , 333, 49-57	14-7	197
114	The Mutation in the Single-Crystal Structural Transformation Process, Induced by the Combined Stimuli of Temperature and Solvent. <i>Chemistry - A European Journal</i> , 2018 , 24, 327-331	4-8	5
113	Two MOFs as dual-responsive photoluminescence sensors for metal and inorganic ion detection. <i>Dalton Transactions</i> , 2018 , 47, 8257-8263	4-3	33
112	Cd-Based metal-organic frameworks from solvothermal reactions involving in situ aldimine condensation and the highly sensitive detection of Fe ions. <i>Dalton Transactions</i> , 2017 , 46, 2332-2338	4-3	39
111	Exploring the Detection of Metal Ions by Tailoring the Coordination Mode of V-Shaped Thienylpyridyl Ligand in Three MOFs. <i>Inorganic Chemistry</i> , 2017 , 56, 2936-2940	5-1	51
110	Six isostructural lanthanide-containing MOFs built on a semi-rigid tripodal organic ligand. <i>Inorganic Chemistry Communication</i> , 2017 , 78, 1-4	3-1	16
109	Photodegradation of Some Organic Dyes over Two Metal-Organic Frameworks with Especially High Efficiency for Safranin T. <i>Crystal Growth and Design</i> , 2017 , 17, 1293-1298	3-5	64

108	Assembly of Zr-MOF crystals onto magnetic beads as a highly adsorbent for recycling nitrophenol. <i>Chemical Engineering Journal</i> , 2017 , 323, 74-83	14.7	55
107	Two New Luminescent Cd(II)-Metal-Organic Frameworks as Bifunctional Chemosensors for Detection of Cations Fe ³⁺ , Anions CrO ₄ ²⁻ and Cr ₂ O ₇ ²⁻ in Aqueous Solution. <i>Crystal Growth and Design</i> , 2017 , 17, 67-72	3.5	242
106	Structures and applications of metal-organic frameworks featuring metal clusters. <i>CrystEngComm</i> , 2017 , 19, 745-757	3.3	20
105	Five New Transition Metal Coordination Polymers Based on V-Shaped Bis-triazole Ligand with Aromatic Dicarboxylates: Syntheses, Structures, and Properties. <i>Crystal Growth and Design</i> , 2017 , 17, 2757-2766	3.5	29
104	Construction of a novel Cd(II) coordination polymer based on a flexible tripodal carboxylic acid and bimid coligands. <i>Inorganic Chemistry Communication</i> , 2017 , 79, 17-20	3.1	5
103	A second-order nonlinear optical material with a 5-fold interpenetrating diamondoid framework based on two achiral precursors: spontaneous resolution to absolute chiral induction. <i>Dalton Transactions</i> , 2017 , 46, 4589-4594	4.3	22
102	Two new Zn(II)/Cu(II) complexes based on bi- and tritopic 1,2,4-triazole derivatives with glutaric acid: Syntheses, structures, luminescent and magnetic properties. <i>Inorganic Chemistry Communication</i> , 2017 , 79, 21-24	3.1	11
101	Two Lanthanide Metal-Organic Frameworks as Remarkably Selective and Sensitive Bifunctional Luminescence Sensor for Metal Ions and Small Organic Molecules. <i>ACS Applied Materials & Interfaces</i> , 2017 , 9, 1629-1634	9.5	282
100	A pair of 3D enantiotopic zinc(ii) complexes based on two asymmetric achiral ligands. <i>Dalton Transactions</i> , 2017 , 46, 14779-14784	4.3	7
99	One 2D anionic coordination polymer with {[Co(H ₂ O) ₆]} ²⁺ cationic guest for fast and selective adsorption of cationic dyes. <i>Inorganic Chemistry Communication</i> , 2017 , 85, 89-91	3.1	2
98	The impact of adjusting auxiliary donors on the performance of dye-sensitized solar cells based on phenothiazine D-D- π A sensitizers. <i>Dyes and Pigments</i> , 2017 , 146, 127-135	4.6	21
97	Two stable 3D porous metal-organic frameworks with high selectivity for detection of PA and metal ions. <i>Dyes and Pigments</i> , 2017 , 136, 515-521	4.6	46
96	Diverse structures of metal-organic frameworks via a side chain adjustment: interpenetration and gas adsorption. <i>Dalton Transactions</i> , 2016 , 45, 16205-16210	4.3	7
95	Three Highly Stable Cobalt MOFs Based on "Y"-Shaped Carboxylic Acid: Synthesis and Absorption of Anionic Dyes. <i>Inorganic Chemistry</i> , 2016 , 55, 8816-21	5.1	63
94	A new five-coordinated copper compound for efficient degradation of methyl orange and Congo red in the absence of UV-visible radiation. <i>Dalton Transactions</i> , 2016 , 45, 18566-18571	4.3	29
93	H-Bonding Interactions Induced Two Isostructural Cd(II) Metal-Organic Frameworks Showing Different Selective Detection of Nitroaromatic Explosives. <i>Inorganic Chemistry</i> , 2016 , 55, 10999-11005	5.1	88
92	Enhanced performance of dye-sensitized solar cells with Y-shaped organic dyes containing di-anchoring groups. <i>New Journal of Chemistry</i> , 2016 , 40, 2799-2805	3.6	23
91	Two new luminescent Cd(II)/Zn(II) metal-organic frameworks for exceptionally selective detection of picric acid explosives. <i>Inorganic Chemistry Communication</i> , 2016 , 66, 51-54	3.1	9

90	Four coordination polymers derived from a one-pot reaction and their controlled synthesis. <i>Dalton Transactions</i> , 2016 , 45, 6418-23	4.3	6
89	Syntheses, Structures, and Properties of Four Metal-Organic Frameworks Based on a N-Centered Multidentate Pyridine-Carboxylate Bifunctional Ligand. <i>Crystal Growth and Design</i> , 2016 , 16, 4711-4719	3.5	15
88	Effects of structural optimization on the performance of dye-sensitized solar cells: spirobifluorene as a promising building block to enhance Voc. <i>Journal of Materials Chemistry A</i> , 2016 , 4, 11782-11788	13	31
87	Dicarboxylate-dependent structural diversity in amino-functionalized complexes: From mononuclear to multinuclear coordination polymer. <i>Inorganic Chemistry Communication</i> , 2016 , 69, 4-6	3.1	5
86	Syntheses, Characterization, and Luminescence Properties of Four Metal-Organic Frameworks Based on a Linear-Shaped Rigid Pyridine Ligand. <i>Crystal Growth and Design</i> , 2016 , 16, 2496-2503	3.5	48
85	Zn(II)/Cd(II) Terephthalate Coordination Polymers Incorporating Bi-, Tri-, and Tetraptopic Phenylamine Derivatives: Crystal Structures and Photoluminescent Properties. <i>Crystal Growth and Design</i> , 2016 , 16, 2747-2755	3.5	48
84	Cyclopentaneteracarboxylic Metal-Organic Frameworks: Tuning the Distance between Layers and Pore Structures with N-Ligands. <i>Inorganic Chemistry</i> , 2016 , 55, 4951-7	5.1	14
83	Insight into the effects of modifying μ bridges on the performance of dye-sensitized solar cells containing triphenylamine dyes. <i>Physical Chemistry Chemical Physics</i> , 2016 , 18, 29555-29560	3.6	14
82	Syntheses, characterization, and magnetic properties of novel divalent Co/Ni coordination polymers based on a V-shaped pyridine ligand and dicarboxylate acids. <i>RSC Advances</i> , 2015 , 5, 64514-64519	3.7	9
81	Porous and single crystalline Co ₃ O ₄ nanospheres for pseudocapacitors with enhanced performance. <i>RSC Advances</i> , 2015 , 5, 27266-27272	3.7	7
80	Three different metal-organic frameworks derived from a one-pot crystallization and their controllable synthesis. <i>Chemical Communications</i> , 2015 , 51, 8338-41	5.8	15
79	Two luminescent Zn(II) metal-organic frameworks for exceptionally selective detection of picric acid explosives. <i>Chemical Communications</i> , 2015 , 51, 8300-3	5.8	199
78	Syntheses, structures, and properties of six cobalt(II) complexes based on a tripodal tris(4-(1H-1,2,4-triazol-1-yl)phenyl)amine ligand. <i>Dalton Transactions</i> , 2015 , 44, 16854-64	4.3	19
77	One non-interpenetrated chiral porous multifunctional metal-organic framework and its applications for sensing small solvent molecules and adsorption. <i>Chemical Communications</i> , 2015 , 51, 2447-9	5.8	57
76	Improvement of dye-sensitized solar cells performance through introducing different heterocyclic groups to triarylamine dyes. <i>RSC Advances</i> , 2015 , 5, 3720-3727	3.7	11
75	Critical factors influencing the structures and properties of metal-organic frameworks. <i>CrystEngComm</i> , 2015 , 17, 981-991	3.3	28
74	Effects of heterocycles containing different atoms as μ bridges on the performance of dye-sensitized solar cells. <i>Physical Chemistry Chemical Physics</i> , 2015 , 17, 16334-40	3.6	26
73	Picolinic acid as an efficient tridentate anchoring group adsorbing at Lewis acid sites and Brønsted acid sites of the TiO ₂ surface in dye-sensitized solar cells. <i>Journal of Materials Chemistry A</i> , 2015 , 3, 14805-14815	13	25

72	A rare three-coordinated zinc cluster-organic framework with two types of secondary building units. <i>Chemical Communications</i> , 2015 , 51, 2899-902	5.8	21
71	Syntheses, Characterizations, Luminescent Properties, and Controlling Interpenetration of Five Metal-Organic Frameworks Based on Bis(4-(pyridine-4-yl)phenyl)amine. <i>Crystal Growth and Design</i> , 2015 , 15, 1303-1310	3.5	29
70	Assembly of various degrees of interpenetration of Co-MOFs based on mononuclear or dinuclear cluster units: magnetic properties and gas adsorption. <i>Dalton Transactions</i> , 2015 , 44, 4751-8	4.3	26
69	Diverse structures of metal-organic frameworks based on different metal ions: luminescence and gas adsorption properties. <i>Dalton Transactions</i> , 2015 , 44, 4238-45	4.3	20
68	Four new metal-organic frameworks based on a rigid linear ligand: synthesis, optical properties and structural investigation. <i>CrystEngComm</i> , 2014 , 16, 5662-5671	3.3	17
67	Improvement of photovoltaic performance of DSSCs by modifying panchromatic zinc porphyrin dyes with heterocyclic units. <i>Journal of Materials Chemistry A</i> , 2014 , 2, 20841-20848	13	11
66	Syntheses, structures, and photoluminescent properties of a series of metal-organic frameworks constructed by 5,5'-bis(1H-imidazol-1-yl)-2,2'-bithiophene and various carboxylate ligands. <i>CrystEngComm</i> , 2014 , 16, 900-909	3.3	21
65	Two pairs of isomorphism and two 3D metal-organic frameworks based on a star-like ligand tri(4-pyridylphenyl)amine. <i>CrystEngComm</i> , 2014 , 16, 698-706	3.3	23
64	Syntheses, characterization, and properties of five coordination compounds based on the ligand tetrakis(4-pyridyloxymethylene)methane. <i>CrystEngComm</i> , 2014 , 16, 3917-3925	3.3	10
63	Interpenetrated Metal-Organic Framework with Selective Gas Adsorption and Luminescent Properties. <i>Crystal Growth and Design</i> , 2014 , 14, 2742-2746	3.5	32
62	Crystal Structures and Spectroscopic Properties of Metal-Organic Frameworks Based on Rigid Ligands with Flexible Functional Groups. <i>Crystal Growth and Design</i> , 2014 , 14, 491-499	3.5	57
61	Promising alkoxy-wrapped porphyrins with novel push-pull moieties for dye-sensitized solar cells. <i>Journal of Materials Chemistry A</i> , 2014 , 2, 14883-14889	13	16
60	One rutile Co(II) coordinated polymer with bifunctional ligand. <i>Inorganic Chemistry Communication</i> , 2014 , 46, 191-193	3.1	2
59	Molecular Tectonics of Four-Connected Network Topologies by Regulating the Ratios of Tetrahedral and Square-Planar Building Units. <i>Crystal Growth and Design</i> , 2014 , 14, 6607-6612	3.5	12
58	Application of WO ₆ -based secondary building units in functional metal-organic frameworks. <i>CrystEngComm</i> , 2013 , 15, 9265	3.3	9
57	Syntheses, characterizations and properties of five new metal-organic complexes based on flexible ligand 4,4'-(phenylazanediy) dibenzoic acid. <i>CrystEngComm</i> , 2013 , 15, 616-627	3.3	23
56	A second-order nonlinear optical material with a hydrated homochiral helix obtained via spontaneous symmetric breaking crystallization from an achiral ligand. <i>Chemical Communications</i> , 2013 , 49, 3585-7	5.8	46
55	The synthesis, structure and third-order nonlinear optical effect of a new 2D cluster polymer based on a [W ₄ Cu ₄] ²⁺ SBU and 1,2-di(pyridin-4-yl)ethane. <i>CrystEngComm</i> , 2013 , 15, 7354	3.3	14

54	Chiral 3D/3D hetero-interpenetrating framework with six kinds of helices, 3D polyrotaxane and 2D network via one-pot reaction. <i>CrystEngComm</i> , 2013 , 15, 227-230	3.3	28
53	Syntheses, Structures, Photochemical and Magnetic Properties of Novel Divalent Cd/Mn Coordination Polymers Based on a Semirigid Tripodal Carboxylate Ligand. <i>Crystal Growth and Design</i> , 2013 , 13, 1694-1702	3.5	26
52	Tuning Structural Topologies of a Series of Metal-Organic Frameworks: Different Bent Dicarboxylates. <i>Crystal Growth and Design</i> , 2013 , 13, 2111-2117	3.5	27
51	Series of Metal-Organic Frameworks Including Novel Architectural Features Based on a Star-like Tri(4-pyridylphenyl)amine Ligand. <i>Crystal Growth and Design</i> , 2013 , 13, 1961-1969	3.5	66
50	Anion-selectivity of cationic cluster-organic nanospheres based on a nest-shaped [MS ₄ Cu ₃ X ₃] cluster monomer with a ditopic ligand. <i>CrystEngComm</i> , 2013 , 15, 5016	3.3	14
49	A porous metal-organic framework based on Zn ₆ O ₂ clusters: chemical stability, gas adsorption properties and solvatochromic behavior. <i>Chemical Communications</i> , 2013 , 49, 555-7	5.8	106
48	Three 2D/2D-j2D or 3D Coordination Polymers: Parallel Stacked, Interpenetration, and Polycatenated. <i>Crystal Growth and Design</i> , 2013 , 13, 5045-5049	3.5	29
47	Metal-organic frameworks constructed from versatile [WS ₄ Cu(x)](x-2) units: micropores in highly interpenetrated systems. <i>Chemistry - A European Journal</i> , 2012 , 18, 2812-24	4.8	53
46	Metal-organic frameworks constructed from flexible V-shaped ligands: adjustment of the topology, interpenetration and porosity via a solvent system. <i>Chemical Communications</i> , 2012 , 48, 10016-8	5.8	92
45	A microporous metal-organic framework with FeS(2) topology based on [Zn ₆ (β-O)] cluster for reversible sensing of small molecules. <i>Chemical Communications</i> , 2012 , 48, 7967-9	5.8	82
44	Metal-Organic Frameworks Based on Flexible V-Shaped Polycarboxylate Acids: Hydrogen Bondings, Non-Interpenetrated and Polycatenated. <i>Crystal Growth and Design</i> , 2012 , 12, 4072-4082	3.5	65
43	Controlled Synthesis of Three-Fold Dendrites of Ce(OH)CO ₃ with Multilayer Caltrop and Their Thermal Conversion to CeO ₂ . <i>Crystal Growth and Design</i> , 2012 , 12, 271-280	3.5	28
42	Construction of Metal-Organic Frameworks Based on Two Neutral Tetradentate Ligands. <i>Crystal Growth and Design</i> , 2012 , 12, 4911-4918	3.5	24
41	Structure-property relationship of homochiral and achiral supramolecular isomers obtained by one-pot synthesis. <i>Chemical Communications</i> , 2012 , 48, 10757-9	5.8	42
40	Structural Diversity and Properties of Six 2D or 3D Metal-Organic Frameworks Based on Thiophene-Containing Ligand. <i>Crystal Growth and Design</i> , 2012 , 12, 5783-5791	3.5	21
39	Construction of a series of metal-organic frameworks with a neutral tetradentate ligand and rigid carboxylate co-ligands. <i>CrystEngComm</i> , 2012 , 14, 8274	3.3	12
38	Diverse Structures of Metal-Organic Frameworks Based on a New Star-Like Tri(4-pyridylphenyl)amine Ligand. <i>Crystal Growth and Design</i> , 2012 , 12, 3957-3963	3.5	50
37	Three self-penetrated, interlocked, and polycatenated supramolecular isomers via one-pot synthesis and crystallization. <i>Chemical Communications</i> , 2012 , 48, 681-3	5.8	75

36	Effect of Carboxylate Coligands with Different Rigidity on Supramolecular Architectures Based on One Rigid Didentate Linear Ligand. <i>Crystal Growth and Design</i> , 2012 , 12, 403-413	3.5	77
35	Five Novel Coordination Polymers Based on a C-Centered Triangular Flexible Ligand. <i>Crystal Growth and Design</i> , 2012 , 12, 1022-1031	3.5	36
34	Synthesis and properties of five unexpected copper complexes with ring-cleavage of 3,6-di-2-pyridyl-1,2,4,5-tetrazine by one pot in situ hydrothermal reaction. <i>CrystEngComm</i> , 2012 , 14, 2258-2263	3.3	21
33	Syntheses, Structures, and Characteristics of Four New Metal-Organic Frameworks Based on Flexible Tetrapyridines and Aromatic Polycarboxylate Acids. <i>Crystal Growth and Design</i> , 2012 , 12, 3426-3435	3.5	70
32	Solvatochromic behavior of a nanotubular metal-organic framework for sensing small molecules. <i>Journal of the American Chemical Society</i> , 2011 , 133, 4172-4	16.4	609
31	The rational synthesis of (10,3)-type MOFs based on tetranuclear [W(Mo)OS ₃ Cu ₃] ⁺ secondary building units. <i>Chemical Communications</i> , 2011 , 47, 10049-51	5.8	64
30	Syntheses, structures, photoluminescence and magnetic properties of four new metal-organic frameworks based on imidazole ligands and aromatic polycarboxylate acids. <i>CrystEngComm</i> , 2011 , 13, 857-865	3.3	46
29	Six new metal-organic frameworks based on polycarboxylate acids and V-shaped imidazole-based synthon: syntheses, crystal structures, and properties. <i>Inorganic Chemistry</i> , 2011 , 50, 2404-14	5.1	88
28	Syntheses, structures, magnetic and photoluminescence properties of metal-organic frameworks based on aromatic polycarboxylate acids. <i>CrystEngComm</i> , 2011 , 13, 1617-1624	3.3	35
27	Solvothermal synthesis, structures and physical properties of four new complexes constructed from multi-variant tricarboxylate ligand and pyridyl-based ligands. <i>CrystEngComm</i> , 2011 , 13, 459-466	3.3	47
26	[WS ₄ Cu ₃ I ₂] ⁻ and [WS ₄ Cu ₄] ²⁺ secondary building units formed a metal-organic framework: large tubes in a highly interpenetrated system. <i>Chemical Communications</i> , 2011 , 47, 2919-21	5.8	70
25	Three New Coordination Polymers Based on One Reduced Symmetry Tripodal Linker. <i>Crystal Growth and Design</i> , 2011 , 11, 3115-3121	3.5	64
24	Syntheses, Structures, and Photoluminescence of Five New Metal-Organic Frameworks Based on Flexible Tetrapyridines and Aromatic Polycarboxylate Acids. <i>Crystal Growth and Design</i> , 2010 , 10, 2676-2684	3.5	102
23	Syntheses, Characterizations, and Properties of Six Metal-Organic Complexes Based on Flexible Ligand 5-(4-Pyridyl)-methoxyl Isophthalic Acid. <i>Crystal Growth and Design</i> , 2010 , 10, 4176-4183	3.5	83
22	Syntheses, structures, photoluminescence and magnetic properties of five compounds with 1,3,5-benzenetricarboxylate acid and imidazole ligands. <i>CrystEngComm</i> , 2010 , 12, 612-619	3.3	58
21	Organic-organic hybrid coordination polymers based on the 5-oxycetate isophthalic acid (H ₃ OABDC) ligand: syntheses, structures, magnetic and luminescent properties. <i>CrystEngComm</i> , 2010 , 12, 4424	3.3	19
20	Eight new complexes based on flexible multicarboxylate ligands: synthesis, structures and properties. <i>CrystEngComm</i> , 2010 , 12, 3183	3.3	10
19	Three new heterothiometallic cluster polymers with fascinating topologies. <i>Inorganic Chemistry</i> , 2009 , 48, 5772-8	5.1	69

18	Unusual three-dimensional coordination networks with [WS ₄ Cu ₆] cluster nodes and EC ₃ N ₄ topology. <i>CrystEngComm</i> , 2009 , 11, 605-609	3.3	18
17	Studies on the Thermodynamic and Kinetic Properties of Reactions of Bo(Bs) with H ₂ . <i>Progress in Reaction Kinetics and Mechanism</i> , 2006 , 31, 1-9	0.5	3
16	Theoretical study of the insertion reaction of singlet phosphinidene with hydrogen sulfide. <i>Journal of Chemical Research</i> , 2006 , 2006, 303-305	0.6	3
15	Quinoxalines Incorporating Triarylaminines: Dipolar Electroluminescent Materials with Tunable Emission Characteristics. <i>Journal of the Chinese Chemical Society</i> , 2006 , 53, 233-242	1.5	4
14	Structures and stabilities of the donor-acceptor complexes HXPY (X=Al, B; Y=H, F, OH). <i>Molecular Physics</i> , 2006 , 104, 447-452	1.7	3
13	Reactions of singlet phosphinidene and its hydroxy derivative with polar molecule hydrogen fluoride. <i>Molecular Physics</i> , 2006 , 104, 599-605	1.7	6
12	Organic electroluminescent derivatives containing dibenzothiophene and diarylamine segments. <i>Journal of Materials Chemistry</i> , 2005 , 15, 3233		19
11	Synthesis, crystal structure and nonlinear optical properties of a cluster compound containing the bipy ligand. <i>Transition Metal Chemistry</i> , 2004 , 29, 185-188	2.1	8
10	Self-assembly of interpenetrating coordination nets formed from interpenetrating cationic and anionic three-dimensional diamondoid cluster coordination polymers. <i>Angewandte Chemie - International Edition</i> , 2004 , 43, 5776-9	16.4	171
9	Crystal Structure and Excited Optical Nonlinearity of a 1D Polymeric [W ₂ O ₂ S ₆ Cu ₄ (NCMe) ₄] _n Cluster. <i>European Journal of Inorganic Chemistry</i> , 2004 , 2004, 2754-2758	2.3	9
8	Synthesis, crystal structure and nonlinear optical properties of a new cluster complex: WCu ₃ OS ₃ (PPh ₃) ₃ {S ₂ P(OPri) ₂ }. <i>Transition Metal Chemistry</i> , 2003 , 28, 137-141	2.1	2
7	Synthesis, crystal structure and non-linear optical properties of the heterobimetallic polymeric compound {[n-Bu ₄ N][W ₂ Ag ₃ S ₈]} _n . <i>CrystEngComm</i> , 2003 , 5, 62-64	3.3	12
6	Synthesis, Crystal Structure and Nonlinear Optical Properties of a new cluster compound: MoS ₄ Cu ₃ (PyPPH ₂) ₃ Cl. <i>Journal of Coordination Chemistry</i> , 2003 , 56, 595-601	1.6	2
5	Synthesis, Structural Characterization of a Novel 4,4'-Bipyridyl Based HgI ₂ Adduct. <i>Synthesis and Reactivity in Inorganic, Metal Organic, and Nano Metal Chemistry</i> , 2003 , 33, 1-10		13
4	Synthesis and Crystal Structures of Two Nest-Shaped Cluster Compounds, [MoOS ₃ Cu ₃ (SCN)py ₅] and [WOS ₃ Cu ₃ (SCN)py ₅]. <i>Synthesis and Reactivity in Inorganic, Metal Organic, and Nano Metal Chemistry</i> , 2000 , 30, 761-775		2
3	Syntheses, crystal structures and non-linear optical properties of two novel windmill-shaped clusters: [M ₂ Pd ₄ S ₈ (dppm) ₂][DMF] (M = W or Mo). <i>Dalton Transactions RSC</i> , 2000 , 2145-2149		22
2	Metal-organic frameworks constructed from an [MS ₄ Cu _x] _x (M = W, Mo) unit: isomerization of the cluster unit induced by temperature. <i>CrystEngComm</i> ,	3.3	
1	A novel and efficient method of MOF-derived electrocatalyst for HER performance through doping organic ligands. <i>Materials Chemistry Frontiers</i> ,	7.8	2

